

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 August 2001 (02.08.2001)

PCT

(10) International Publication Number
WO 01/55411 A2

(51) International Patent Classification⁷: **C12N 15/55**,
5/10, 9/16, C07K 16/40, G01N 33/53, C12Q 1/68, A61K
38/46

(74) Agents: **KRON, Eric, J. et al.**; Alston & Bird LLP, P.O.
Drawer 34009, Charlotte, NC 28234-4009 (US).

(21) International Application Number: **PCT/US01/03266**

(22) International Filing Date: 31 January 2001 (31.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/495,823 31 January 2000 (31.01.2000) US

(71) Applicant (*for all designated States except US*): **MIL-
LENNIUM PHARMACEUTICALS, INC.** [US/US]; 75
Sidney Street, Cambridge, MA 02139 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GLUCKSMANN, Maria, Alexandra** [AR/US]; 33 Summit Road, Lexington, MA 02173 (US). **WILLIAMSON, Mark** [US/US]; 15 Stonecrest Drive, Saugus, MA 01906 (US). **RUDOLPH-OWEN, Laura, A.** [US/US]; 186 Arborway, #1, Jamaica Plain, MA 02130 (US). **TSAI, Fong-Ying** [—/US]; 15 Montclair Road, Newton, MA 02468 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 22438, 23553, 25278, AND 26212 NOVEL HUMAN SULFATASES

(57) Abstract: The present invention relates to newly identified human sulfatases. In particular, the invention relates to sulfatase polypeptides and polynucleotides, methods of detecting the sulfatase polypeptides and polynucleotides, and methods of diagnosing and treating sulfatase-related disorders. Also provided are vectors, host cells, and recombinant methods for making and using the novel molecules.

WO 01/55411 A2

22438, 23553, 25278, AND 26212 NOVEL HUMAN SULFATASES

FIELD OF THE INVENTION

The present invention relates to newly identified human sulfatases. In particular, the invention relates to sulfatase polypeptides and polynucleotides, methods of detecting the sulfatase polypeptides and polynucleotides, and methods of diagnosing and treating sulfatase-related disorders. Also provided are vectors, host cells, and recombinant
5 methods for making and using the novel molecules.

BACKGROUND OF THE INVENTION

The biology and functions of the reversible sulfation pathway catalyzed by
10 human sulfotransferases and sulfatases has been reviewed by Coughtrie *et al.* (*Chemico-Biological Interactions* 109: 3-27 (1998)). This review, summarized below, focuses on the sulfation of small molecules carried out by cytosolic sulfotransferases rather than the sulfation of macromolecules and lipids catalyzed by membrane-associated sulfotransferases.

15 Sulfation functions in the metabolism of xenobiotic compounds, steroid biosynthesis, and modulating the biological activity and inactivation and elimination of potent endogenous chemicals such as thyroid hormones, steroids and catechols. This pathway is reversible, comprising the sulfotransferase enzymes that cause the sulfation and the sulfatases that hydrolyze the sulfate esters formed by the action of the
20 sulfotransferases. Accordingly, the interplay between these families regulates the availability and biological activity of xenobiotic and endogenous chemicals. The sulfatases, including the arylsulfatases (ARS), are located in lysosomes or endoplasmic reticulum.

The presence of sulfated components depends upon the availability of key
25 members of the sulfate pathway, i.e., substrate and activated sulfate donor molecule (co-substrate) and the balance between sulfation and sulfate conjugate hydrolysis that depends upon the activity and localization of the sulfotransferases and the sulfatases.

Essentially, divalent sulfate is converted to adenosine 5' phosphosulfate (PAPS) by hydrolysis of ATP. This compound is in turn converted to 3' phosphoadenosine 5' phosphosulfate by hydrolysis of ATP to ADP. This compound is then converted to adenosine 3' 5' biphosphate concurrently with the formation of 4-nitrophenolsulfate from 4-nitrophenol. An ARS would then cleave the monovalent sulfate from the 4-nitrophenolsulfate to produce the original 4-nitrophenol. This forms the basis for the sulfation system in humans. Over- or under-production of any of these key molecules can result in sulfate-related disorders. For example, the brachymorphic mouse has a connective tissue disorder that results from a defect in PAPS formation that causes undersulfated cartilage proteoglycans.

ARS enzymes and their genes have been associated with specific genetic diseases. ARSA is located in the lysosomes and removes sulfate from sulfated glycolipids. A deficiency of ARSA has been associated with metachromatic leukodystrophy and multiple sulfatase deficiency (MSD). ARSB is located in lysosomes and has, as an endogenous substrate, dermatan sulfate and chondroitin sulfate. A deficiency of ARSB is associated with Maroteaux-Lamy syndrome and MSD. ARSC is located in the endoplasmic reticulum and has, as its endogenous substrate, cholesterol sulfate and steroid sulfates. A deficiency of ARSC is associated with X-linked ichthyosis and MSD. ARSD may be associated with MSD. ARSE has been associated with chondrodysplasia punctata and MSD. ARSF may be associated with MSD. ARSC hydrolyses sulfate esters on a wide range of steroids and cholesterol. ARSs also hydrolyse sulfate conjugates of xenobiotics.

MSD results from an inability to perform a co- or post-translational modification of a cysteine residue to serine semialdehyde (2-oxo-3-propionic acid). This residue is conserved in all eukaryotic sulfatases described by Coughtrie *et al.* ARSC may have a very broad specificity, extending to iodothyronine sulfates and a number of sulfate conjugates of xenobiotic phenols.

The kinetic and catalytic properties of ARS enzymes in isolation, important for understanding substrate specificity and the physical and chemical properties of enzymes and substrates that allow substrate preference, have been characterized recently based on recombinant enzyme systems. For the expression of the human sulfotransferases, COS and V79 cells have been used. Coughtrie *et al.* have constructed and characterized V79

cell lines stably expressing ARSA, ARSB, and ARSC. These cell lines exhibited the expected substrate preferences of the three enzymes among the substrates 4-nitrocatechol sulfate, estrone sulfate, and dehydroepiandrosterone sulfate(DHEAS).

5 The sulfation of small molecules can be broadly divided into the areas of chemical defense, hormone biosynthesis, and bioactivation. It was originally viewed that sulfation protected against the toxic effects of xenobiotics in that sulfate conjugates are more readily excreted in urine or bile and generally exhibit reduced pharmacological/biological activity relative to the parent compound. Many drugs and other xenobiotics are conjugated with sulfate. Many phenolic metabolites of the
10 cytochrome P450 mono-oxygenase system are excreted as sulfate conjugates.

Further, potent endogenous chemicals, such as steroids and catecholamines are found at high levels as circulating sulfate conjugates. For example, greater than 90% of circulating dopamine exists as the sulfated form. Sulfation is also suggested to play a role in the inactivation of potent steroids such as estrogens and androgens. Accordingly,
15 sulfation is important in metabolism and homeostasis of such compounds in humans.

DHEAS is the major circulating steroid in humans and estrone sulfate is the major estrogen. These chemicals act as precursors of estrogens and androgens. Extremely large quantities of such steroids or estrogens may occur during various stages of development, such as pregnancy. Estrone sulfate is a precursor for β -estradiol
20 synthesis. In breast cancer cells it is hydrolysed by steroid sulfatase (ARSC) to estrone which is then converted to β -estradiol by action of another enzyme. Accordingly, ARSC is important for maintaining active estrogen. It is thus an important therapeutic target for the treatment of breast cancer.

Cholesterol sulfate, synthesized in the skin epidermis, may have a role in
25 keratinocyte differentiation. Accordingly, hydrolysis of cholesterol sulfate by steroid sulfatase may be important in skin formation and differentiation. This is the major organ affected in X-linked ichthyosis caused by mutations in ARSC.

Although sulfation may widely serve to detoxify potent compounds, some sulfate conjugates are more biologically active than the corresponding parent compound.
30 Minoxidil and cicletanine are activated upon sulfation. Further, an inhibitor of ARSC was shown to potentiate the memory enhancing effect of DHEAS. This suggests a role for sulfates and sulfation in the central nervous system.

An important example of bioactivation by means of sulfation, however, occurs with dietary and environmental mutagens and carcinogens. For a large number of these, sulfation is the terminal step in the pathway to metabolic activation. Examples of such chemicals include aromatic amines (including heterocyclic amines) and benzylic
5 alcohols of chemicals such as polycyclic aromatic hydrocarbons, safrole, and estragole.

The sulfatase gene family has been reviewed in Parenti *et al.* (*Current Opinion in Genetics and Development* 7:386-391 (1997)), summarized below.

The sulfatase family of enzymes is functionally and structurally similar. Nevertheless, these enzymes catalyze the hydrolysis of sulfate ester bonds from a wide
10 variety of substrates ranging from complex molecules such as glycosaminoglycans and sulfolipids to steroid sulfates (see also Coughtrie *et al.*, above). Several human genetic disorders result from the accumulation of intermediate sulfate compounds that result from a deficiency of single or multiple sulfatase activities. A subset of sulfatase, ARS, is characterized by the ability to hydrolyze sulfate esters of chromogenic or fluorogenic
15 aromatic compounds such as *p*-nitrocatechol sulfate and 4-methylumbelliferyl sulfate. Desulfation is required to degrade glycosaminoglycans, heparan sulfate, chondroitin sulfate and dermatan sulfate and sulfolipids. Steroid sulfatase differs from other members of the family with respect to subcellular localization. It is localized in the microsomes rather than in lysosomes. Further, ARSD, ARSE, and ARSF are also non-
20 lysosomal, being localized in the endoplasmic reticulum or Golgi compartment.

The natural substrate of ARSA is cerebroside sulfate. Associated diseases are MLD and MSD. The natural substrate of ARSB is dermatan sulfate. The disease associated with this enzyme is MPSVI and MSD. The natural substrate of ARSC/STS is sulfated steroids. Diseases associated with this enzyme are XLI and MSD. The natural
25 substrates of ARSD-F are unknown. The natural substrates of iduronate-2-sulfatase (IDS) are dermatan sulfate and heparan sulfate. Diseases associated with this enzyme are MPSII and MSD. The natural substrate of galactose 6-sulfatase is keratan sulfate and chondroitin 6-sulfate. Diseases associated with this enzyme include MPSIVA and MSD. The natural substrate of glucosamine-6-sulfatase is heparan sulfate
30 and keratan sulfate. A disease associated with this enzyme is MPSIIID and MSD. The natural substrate of glucuronate-2-sulfatase is heparan sulfate. The natural substrate of glucosamine-3-sulfatase is heparan sulfate.

Sulfatases are activated through conversion of a cysteine residue as described above. The conversion is required for catalytic activity and is defective in MSD. It is likely that all sulfatases undergo the same modification. The substitution of this cysteine was shown to destroy the enzymatic activity of N-acetyl galactosamine-4-sulfatase (ARSB). It has been shown that the modified residue and a metal ion are located at the base of a substrate binding pocket.

Nine human sulfatase genes are known and murine rat, goat, or avian orthologs for some of these have been identified. A high degree of similarity occurs particularly in the amino terminal region which contains accordingly a potential consensus sulfatase signature.

Sulfatases, as discussed above, are associated with human disease. Most sulfatase deficiencies cause lysosomal storage disorders. The mucopolysaccharidoses contain various associations of mental retardation, facial dysmorphisms, skeletal deformities, hepatosplenomegaly, and deformities of soft tissues caused by deficiencies of sulfatases acting on glycosaminoglycans. In metachromatic leukodystrophy, a deficiency of ARSA causes the storage of sulfolipids in the central and peripheral nervous systems, leading to neurologic deterioration. X-linked ichthyosis is caused by STS deficiency leading to increased cholesterol sulfate levels. MSD, a disorder in which all sulfatase activities are simultaneously defective, was shown to result from a defect in the co- or post-translational processing of sulfatases.

Accordingly, sulfatases are a major target for drug action and development. Therefore, it is valuable to the field of pharmaceutical development to identify and characterize previously unknown sulfatases. The present invention advances the state of the art by providing previously unidentified human sulfatases.

25

SUMMARY OF THE INVENTION

Novel sulfatase nucleotide sequences, and the deduced sulfatase polypeptides are described herein. Accordingly, the invention provides isolated sulfatase nucleic acid molecules having the sequences shown in SEQ ID NOS:2, 4, 6, and 8 or in the cDNA deposited with ATCC as Patent Deposit Number _____, PTA-1639, PTA-1846, or _____, respectively ("the deposited cDNA"), and variants and fragments thereof.

30

It is also an object of the invention to provide nucleic acid molecules encoding the sulfatase polypeptides, and variants and fragments thereof. Such nucleic acid molecules are useful as targets and reagents in sulfatase expression assays, are applicable to treatment and diagnosis of sulfatase-related disorders and are useful for producing
5 novel sulfatase polypeptides by recombinant methods.

The invention thus further provides nucleic acid constructs comprising the nucleic acid molecules described herein. In a preferred embodiment, the nucleic acid molecules of the invention are operatively linked to a regulatory sequence. The invention also provides vectors and host cells for expressing the sulfatase nucleic acid
10 molecules and polypeptides, and particularly recombinant vectors and host cells.

In another aspect, it is an object of the invention to provide isolated sulfatase polypeptides and fragments and variants thereof, including a polypeptide having the amino acid sequence shown in SEQ ID NOS:1, 3, 5 or 7 or the amino acid sequences encoded by the deposited cDNAs. The disclosed sulfatase polypeptides are useful as
15 reagents or targets in sulfatase assays and are applicable to treatment and diagnosis of sulfatase-related disorders.

The invention also provides assays for determining the activity of or the presence or absence of the sulfatase polypeptides or nucleic acid molecules in a biological sample, including for disease diagnosis. In addition, the invention provides assays for
20 determining the presence of a mutation in the polypeptides or nucleic acid molecules, including for disease diagnosis.

A further object of the invention is to provide compounds that modulate expression of the sulfatase for treatment and diagnosis of sulfatase-related disorders. Such compounds may be used to treat conditions related to aberrant activity or
25 expression of the sulfatase polypeptides or nucleic acids.

The disclosed invention further relates to methods and compositions for the study, modulation, diagnosis and treatment of sulfatase related disorders. The compositions include sulfatase polypeptides, nucleic acids, vectors, transformed cells and related variants thereof. In particular, the invention relates to the diagnosis and
30 treatment of sulfatase-related disorders including, but not limited to disorders as described in the background above, further herein, or involving a tissue shown in the figures herein.

In yet another aspect, the invention provides antibodies or antigen-binding fragments thereof that selectively bind the sulfatase polypeptides and fragments. Such antibodies and antigen binding fragments have use in the detection of the sulfatase polypeptide, and in the prevention, diagnosis and treatment of sulfatase related disorders.

5 The sulfatases disclosed herein are designated as follows: 22438, 23553, 25278, and 26212.

DESCRIPTION OF THE DRAWINGS

10 Figure 1 shows the 22438 sulfatase cDNA sequence (SEQ ID NO:2) and the deduced amino acid sequence (SEQ ID NO:1). The 22438 sulfatase coding sequence is set forth in SEQ ID NO:11.

15 Figure 2 shows a 22438 sulfatase hydrophobicity plot. Relative hydrophobic residues are shown above the dashed horizontal line, and relative hydrophilic residues are below the dashed horizontal line. The cysteine residues (cys) and N glycosylation site (Ngly) are indicated by short vertical lines just below the hydropathy trace. The numbers corresponding to the amino acid sequence (shown in SEQ ID NO:1) of 22438 sulfatase are indicated. Polypeptides of the invention include fragments which
20 include: all or a part of a hydrophobic sequence (a sequence above the dashed line); or all or part of a hydrophilic fragment (a sequence below the dashed line). Other fragments include a cysteine residue or as N-glycosylation site.

25 Figure 3 shows an analysis of the 22438 sulfatase amino acid sequence: α turn and coil regions; hydrophilicity; amphipathic regions; flexible regions; antigenic index; and surface probability plot.

30 Figure 4 shows an analysis of the 22438 sulfatase open reading frame for amino acids corresponding to specific functional sites. For the N-glycosylation sites, the actual modified residue is the first amino acid. For cAMP- and cGMP-dependent protein kinase phosphorylation sites, the actual modified residue is the last amino acid. For protein kinase C phosphorylation sites, the actual modified residue is the

first amino acid. For casein kinase II phosphorylation sites, the actual modified residue is the first amino acid. For N-myristoylation sites, the actual modified residue is the first amino acid. In addition, an amidation site is found from about amino acids 56-59, an EGF-like domain cysteine pattern signature found from about amino acids 260-271, and a sulfatase signature is found from about amino acids 129-138.

Figure 5 shows the 23553 sulfatase cDNA sequence (SEQ ID NO:4) and the deduced amino acid sequence (SEQ ID NO:3). The 23553 sulfatase coding sequence is set forth in SEQ ID NO:12.

10

Figure 6 shows a 23553 sulfatase hydrophobicity plot. Relative hydrophobic residues are shown above the dashed horizontal line, and relative hydrophilic residues are below the dashed horizontal line. The cysteine residues (cys) and N glycosylation site (Ngly) are indicated by short vertical lines just below the hydropathy trace. The numbers corresponding to the amino acid sequence (shown in SEQ ID NO:3) of 23553 sulfatase are indicated. Polypeptides of the invention include fragments which include: all or a part of a hydrophobic sequence (a sequence above the dashed line); or all or part of a hydrophilic fragment (a sequence below the dashed line). Other fragments include a cysteine residue or as N-glycosylation site.

20

Figure 7 shows an analysis of the 23553 sulfatase amino acid sequence: α turn and coil regions; hydrophilicity; amphipathic regions; flexible regions; antigenic index; and surface probability plot.

25

Figure 8 shows an analysis of the 23553 sulfatase open reading frame for amino acids corresponding to specific functional sites. For the N-glycosylation sites, the actual modified residue is the first amino acid. For protein kinase C phosphorylation sites, the actual modified residue is the first amino acid. For casein kinase II phosphorylation sites, the actual modified residue is the first amino acid. For the tyrosine kinase phosphorylation site, the actual modified residue is the last amino acid residue. For N-myristoylation sites, the actual modified residue is the first amino acid. In addition, a sulfatase signature is found from about amino acids 85-97.

30

Figure 9 shows relative expression of the 23553 sulfatase mRNA in normal and cancerous human tissues.

5 Figure 10 shows the 25278 sulfatase cDNA sequence (SEQ ID NO:6) and the deduced amino acid sequence (SEQ ID NO:5). The 25278 sulfatase coding sequence is set forth in SEQ ID NO:13.

10 Figure 11 shows a 25278 sulfatase hydrophobicity plot. Relative hydrophobic residues are shown above the dashed horizontal line, and relative hydrophilic residues are below the dashed horizontal line. The cysteine residues (cys) and N glycosylation site (Ngly) are indicated by short vertical lines just below the hydropathy trace. The numbers corresponding to the amino acid sequence (shown in SEQ ID NO:5) of 25278 sulfatase are indicated. Polypeptides of the invention include fragments which
15 include: all or a part of a hydrophobic sequence (a sequence above the dashed line); or all or part of a hydrophilic fragment (a sequence below the dashed line). Other fragments include a cysteine residue or as N-glycosylation site.

 Figure 12 shows an analysis of the 25278 sulfatase amino acid sequence:
20 α turn and coil regions; hydrophilicity; amphipathic regions; flexible regions; antigenic index; and surface probability plot.

 Figure 13 shows an analysis of the 25278 sulfatase open reading frame for amino acids corresponding to specific functional sites. For the N-glycosylation sites,
25 the actual modified residue is the first amino acid. For cAMP- and cGMP-dependent protein kinase phosphorylation sites, the actual modified residue is the last amino acid. For protein kinase C phosphorylation sites, the actual modified residue is the first amino acid. For casein kinase II phosphorylation sites, the actual modified residue is the first amino acid. For the tyrosine kinase phosphorylation site, the actual
30 modified residue is the last amino acid residue. For N-myristoylation sites, the actual modified residue is the first amino acid. In addition, amidation sites are found from

about amino acids 312-315 and 541-544, and sulfatase signatures are found from about amino acids 139-148 and 91-103.

Figure 14 shows relative expression of 25278 sulfatase mRNA in normal and cancerous human tissues.

Figure 15 shows the 26212 sulfatase cDNA sequence (SEQ ID NO:8) and the deduced amino acid sequence (SEQ ID NO:7). The 26212 sulfatase coding sequence is set forth in SEQ ID NO:14.

Figure 16 shows a 26212 sulfatase hydrophobicity plot. Relative hydrophobic residues are shown above the dashed horizontal line, and relative hydrophilic residues are below the dashed horizontal line. The cysteine residues (cys) and N glycosylation site (Ngly) are indicated by short vertical lines just below the hydropathy trace. The numbers corresponding to the amino acid sequence (shown in SEQ ID NO:7) of 26212 sulfatase are indicated. Polypeptides of the invention include fragments which include: all or a part of a hydrophobic sequence (a sequence above the dashed line); or all or part of a hydrophilic fragment (a sequence below the dashed line). Other fragments include a cysteine residue or as N-glycosylation site.

Figure 17 shows an analysis of the 26212 sulfatase amino acid sequence: α turn and coil regions; hydrophilicity; amphipathic regions; flexible regions; antigenic index; and surface probability plot.

Figure 18 shows an analysis of the 26212 sulfatase open reading frame for amino acids corresponding to specific functional sites. For the N-glycosylation sites, the actual modified residue is the first amino acid. For cAMP- and cGMP-dependent protein kinase phosphorylation sites, the actual modified residue is the last amino acid. For protein kinase C phosphorylation sites, the actual modified residue is the first amino acid. For casein kinase II phosphorylation sites, the actual modified residue is the first amino acid. For the tyrosine kinase phosphorylation site, the actual modified residue is the last amino acid residue. For N-myristoylation sites, the actual

modified residue is the first amino acid. In addition, sulfatase signature sites are found from about amino acids 168-177 and 120-132.

Figure 19 depicts an alignment of the 22438 sulfatase domain with a
5 consensus amino acid sequence derived from a hidden Markov model. The upper sequence is the consensus amino acid sequence (SEQ ID NO:9), while the lower amino acid sequence corresponds to amino acids 36 to 462 of SEQ ID NO:1.

Figure 20 depicts an alignment of the 23553 sulfatase domain with a
10 consensus amino acid sequence derived from a hidden Markov model. The upper sequence is the consensus amino acid sequence (SEQ ID NO:9), while the lower amino acid sequence corresponds to amino acids 43 to 467 of SEQ ID NO:3.

Figure 21 shows the expression of 23553 in the following human carcinoma
15 cell lines: breast cancer cell lines MCF-7, ZR75, T47D, MDA231, and MDA435; colon cancer cell lines DLD-1, SW480, SW620, HCT116, HT29, and Colo205; lung cancer cell lines NCIH125, NCIH69, NCIH322, NCIH460, and A549. Expression levels were determined by reverse transcriptase(RT) quantitative PCR (Taqman® brand quantitative PCR kit, Applied Biosystems). The quantitative PCR reactions
20 were performed according to the kit manufacturer's instructions.

Figure 22 shows the expression of 23553 in clinical samples of normal human breast tissue and the following human breast tumor tissues: ductal in situ carcinoma (DCIS), invasive ductal carcinoma (IDC), and invasive lobular carcinoma (ILC).
25 Expression levels were determined as described in the description of Figure 21.

Figure 23 shows the expression of 23553 in human clinical samples of normal colon, colon tumor; metastatic liver, and normal liver tissue. Expression levels were determined as described in the description of Figure 21.

30

Figure 24 shows the expression of 23553 in normal human lung and adenocarcinoma (AC) and squamous cell carcinoma (SCC) lung tumor tissue. Expression levels were determined as described in the description of Figure 21.

5 Figure 25 shows the expression of 23553 in the following normal human tissues: prostate (column 1), liver (columns 2 and 3), breast (columns 4 and 5), skeletal muscle (column 6), brain (columns 7 and 8), colon (columns 9 and 10), heart (columns 11 and 12), ovary (columns 13 and 14), kidney (columns 15 and 16), lung (columns 17 and 18), vein (columns 19 and 20), trachea (column 21), adipose
10 (columns 22 and 23), small intestine (column 24), thyroid (columns 25 and 26), skin (columns 27 and 28), testes (column 29), placenta (column 30), fetal liver (columns 31 and 32), fetal heart (columns 33 and 34), osteoblasts (undifferentiated, column 35 and primary culture, column 36), fetal spinal cord (column 38), cervix (column 39), spleen (column 40), spinal cord (column 41), thymus (column 42), tonsil (column 43),
15 lymph node (column 44), and aorta (column 45). 23553 was expressed at high levels in trachea, vein, osteoblast, kidney, and testes tissue; significant expression of 23553 was noted in adipose, colon, skeletal muscle, thyroid, and prostate tissues. Expression levels were determined as described in the description of Figure 21.

20 Figure 26 shows the expression of 23553 in the following human tissues: normal brain (column 1), glioblastoma (columns 2-5), normal breast (column 6), breast tumor (columns 7-9), normal colon (column 10), colon tumor (columns 11-13), normal liver (column 14), metastatic colon (columns 15 and 16), normal lung (column 17), lung tumor (columns 18-20), placenta (column 21), fetal adrenal gland (column
25 22), normal skin (columns 23 and 24), and adipose (column 25). 23553 was detectable in all tissues tested, with evidence of increased expression levels in breast, colon, and lung tumors. In addition, 23553 was expressed at an elevated level in glioblastoma tissue, as compared to normal brain tissue. Expression levels were determined as described in the description of Figure 21.

30

Figure 27 depicts an alignment of the 25278 sulfatase domain with a consensus amino acid sequence derived from a hidden Markov model. The upper

sequence is the consensus amino acid sequence (SEQ ID NO:9), while the lower amino acid sequence corresponds to amino acids 47 to 471 of SEQ ID NO:5.

Figure 28 shows the relative expression of 25278 in various human tissues, as follows. Row 1, NDR 19, breast, DCIS (ductal in situ carcinoma); Row 2, MDA 138, breast, normal; Row 3, NDR 01, breast, IDC (invasive ductal carcinoma); Row 4, NDR 15, breast, DC (ductal carcinoma); Row 5, NDR 133, breast, ILC (invasive lobular carcinoma); Row 6, MDA 161, breast, IDC; Row 7, MDA 155, breast, IDC/DCIS; Row 8, PIT 270, lung, normal; Row 9, CHT 427, lung, normal; Row 10, PIT 241, lung, normal; Row 11, PIT 298, lung, normal; Row 12, CHT 800, lung, AC (adenocarcinoma); Row 13, CHT 335, lung, SCC (squamous cell carcinoma); Row 14, CHT447, lung, AC; Row 15, CHT 752, lung, AC; Row 16, CHT 799, lung, AC; Row 17, CHT 369, lung, SCC; Row 18, CHT 369, lung, SCC; Row 19, CHT 371, colon, normal; Row 20, CHT 396, colon, normal; Row 21, CHT 398, colon, normal; Row 22, NDR 104, colon, normal; Row 23, CHT 520, colon, adenocarcinoma; Row 24, CHT 122, colon, adenocarcinoma; Row 25, CHT 536, colon, adenocarcinoma; Row 26, CHT 528, colon, adenocarcinoma; Row 27, CHT 386, colon, adenocarcinoma; Row 28, CHT 372, colon, adenocarcinoma; Row 29, CHT 532, colon, adenocarcinoma; Row 30, CHT 77, liver, metastatic; Row 31, CHT 321, liver, metastatic; Row 32, CHT 84, liver, metastatic; Row 33, NDR 100, liver, metastatic; Row 34, NDR 154, liver, normal; Row 35, CHT 322, liver, normal; Row 36, PIT 51, liver, normal; Row 37, CHT 339, liver, normal; Row 38, PIT 265, breast, normal; Row 39, MDA 335, breast, normal; Row 40, NDR 132, breast, DCIS; Row 41, NDR 13, breast, normal; Row 42, NDR 56, breast, normal.

25

Figure 29 depicts an alignment of the 26212 sulfatase domain with a consensus amino acid sequence derived from a hidden Markov model. The upper sequence is the consensus amino acid sequence (SEQ ID NO:10), while the lower amino acid sequence corresponds to amino acids 76 to 502 of SEQ ID NO:7.

30

Figure 30 shows the expression of 26212 in various human endothelial cells, as follows. Proliferating human umbilical vein endothelial cells (HUVEC) (column 1);

arresting HUVEC (column 2); HUVEC minus growth factor (column 3); proliferating cardiac human microvascular endothelial cells (HMVEC) (columns 4 and 6); arresting cardiac HMVEC (columns 5 and 7); proliferating lung HMVEC (columns 8, 11, and 13); arresting lung HMVEC (columns 9, 12, and 14); and lung HMVEC minus growth factor (columns 10 and 15); HEK 293 (non-endothelial) cells (column 16). In six of six independent experiments, 26212 is up-regulated in proliferating endothelial cells as compared to arrested endothelial cells. Further, 26212 expression levels are higher in proliferating endothelial cells than in HEK 293 (non-endothelial) cells. Expression levels were determined as described in the description of Figure 21.

10

Figure 31 shows the expression of 26212 in the following human tissues. Figure 31A: normal breast (columns 1 and 2), breast tumor (columns 3-9), normal ovary (columns 10 and 11), ovary tumor (columns 12-19), normal lung (columns 20-23), lung tumor (columns 24-31). Figure 31B: normal colon (columns 1-4), colon tumor (columns 5-12), liver metastases (columns 13-16), normal liver (columns 17-18), normal brain (columns 19-20), astrocyte (column 21), brain tumor (columns 22-25), arresting human microvascular endothelial cells (column 26), proliferating human microvascular endothelial cells (column 27), placenta (column 28), fetal adrenal tissue (columns 29-30), and fetal liver (column 31). Expression levels were determined as described in the description of Figure 21.

20

Figure 32 shows 26212 expression in normal human clinical breast samples (columns 1 and 2) and human clinical breast tumor samples (columns 3-9). Expression levels were determined as described in the description of Figure 21.

25

Figure 33 shows 26212 expression in normal human clinical lung samples (columns 1-4) and human clinical lung tumor samples (columns 5-12). Expression levels were determined as described in the description of Figure 21.

Figure 34 shows the temporal expression of 26212 in human normal and breast cancer epithelial cell lines (MCF10A and MCF3B, respectively) after treatment with epidermal growth factor (EGF). MCF10A cells are shown 0, 0.5, 1, 2, 4, and 8 hours

30

after treatment with EGF (columns 1-6, respectively). Similarly, MCF3B cells are shown 0, 0.5, 1, 2, 4, and 8 hours after treatment with EGF (columns 7-12, respectively). 26212 is up-regulated in both cell lines. Expression levels were determined as described in the description of Figure 21.

5

Figure 35 shows expression of 26212 in human hemangiomas and other angiogenic tissues: hemangioma (ONC 101; column 1); hemangioma (ONC 102; column 2); hemangioma (ONC 103; column 3); skin (NDR 295; column 4); fetal heart (BWH4; column 5); normal heart (MPI 849; column 6); spinal cord (CKN 746; column 7); uterine adenocarcinoma (CHT 1424; column 8); and endometrial polyps (CLN 944; column 9). Expression levels were determined as described in the description of Figure 21.

Figure 36 shows expression of 26212 in the following human tissues: normal artery (column 1), normal vein (column 2), aortic smooth muscle cells (SMC), early (column 3), coronary SMC (column 4), static human umbilical vein endothelial cells (HUVEC) (column 5), shear HUVEC (column 6), normal heart (column 7), heart, congestive heart failure (CHF) (column 8), kidney (column 9), skeletal muscle (column 10), normal adipose (column 11), pancreas (column 12), primary osteoblasts (column 13), osteoclasts, differentiated (column 14), normal skin (column 15), normal spinal cord (column 16), normal brain cortex (column 17), normal brain hypothalamus (column 18), nerve (column 19), dorsal root ganglion (DRG) (column 20), glial cells (astrocytes) (column 21), glioblastoma (column 22), normal breast (column 23), breast tumor (column 24), normal ovary (column 25), ovary tumor (column 26), normal prostate (column 27), prostate tumor (column 28), prostate epithelial cells (column 29), normal colon (column 30), colon tumor (column 31), normal lung (column 32), lung tumor (column 33), lung, chronic obstructive pulmonary disease (COPD) (column 34), colon, inflammatory bowel disease (IBD) (column 35), normal liver (column 36), liver fibrosis (column 37), dermal cells, fibroblasts (column 38), normal spleen (column 39), normal tonsil (column 40), lymph node (column 41), small intestine (column 42), skin, decubitus (column 43), synovium (column 44), bone marrow mononuclear cells (BM-MNC) (column 45), and activated peripheral blood mononuclear cells (PBMC) (column

46). The expression levels of 26212 are higher in endothelial and glial cells than in other tissues and cells. Expression levels were determined as described in the description of Figure 21.

5

DETAILED DESCRIPTION OF THE INVENTION

Sulfatase Polypeptides

The invention is based on the identification of the novel human 22438 sulfatase. *In situ* hybridization experiments showed that this sulfatase is expressed in the following monkey tissues: sub-populations of DRG neurons (mainly in small and medium sized neurons), in spinal cord (interneurons and motor neurons), and in the brain. The sulfatase is also expressed in human brain. The sulfatase cDNA was identified based on consensus motifs or protein domains characteristic of sulfatases and, in particular, arylsulfatase. BLAST analysis has shown homology with human arylsulfatase E, a human iduronate-2-sulfatase, human N-acetylgalactosamine-6-sulfatase, murine arylsulfatase A, and human arylsulfatase A. However, some homology has also been found with other arylsulfatases from various mammalian species, including, but not limited to, human arylsulfatase D, E, F, and B.

The invention is also based on the identification of the novel human 23553 sulfatase. Taqman analysis has shown positive differential expression in breast and colon cancer and in colonic metastases to the liver (Figure 9). This sulfatase has been identified as a glucosamine-6-sulfatase based on ProDom matches and BLAST analysis. Some homology has also been found to human arylsulfatase A, human N-acetylglucosamine-6-sulfatase, and human iduronate-2-sulfatase.

The invention is also based on the identification of the novel human 25278 sulfatase. The sulfatase is differentially expressed in human colon cancer and in colonic metastases to the liver, as determined by Taqman analysis. This sulfatase has been identified as a N-acetylgalactosamine-4-sulfatase by ProDom matching and BLAST homology alignment. Further, based on BLAST analysis, some homology has also been shown to arylsulfatase B and arylsulfatase A.

The invention is also based on the identification of the novel human 26212 sulfatase. This sulfatase has been identified as an arylsulfatase by ProDom matching

and BLAST sequence alignment. Homology has been shown to arylsulfatase B. Some homology has also been found with arylsulfatase F, E, D, and A, as well as with iduronate 2 sulfatase. Arylsulfatase B is also known as N-acetylgalactosamine-4-sulfatase.

- 5 Specifically, newly-identified human genes, termed 22438, 23553, 25278, and 26212 sulfatases are provided. These sequences, and other nucleotide sequences encoding the sulfatase proteins or fragments and variants thereof, are referred to as "22438, 23553, 25278, and 26212 sulfatase sequences."

- Plasmids containing the sulfatase cDNA inserts were deposited with the Patent
10 Depository of the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Virginia, on ___, April 5, 2000, May 9, 2000, or ___, and assigned Patent Deposit Numbers ___, PTA-1639, PTA-1846, or ___, respectively. The deposits will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of
15 Patent Procedure. The deposits were made merely as a convenience for those of skill in the art and is not an admission that a deposit is required under 35 U.S.C. §112.

- The sulfatase cDNA was identified in human cDNA libraries. Specifically, expressed sequence tags (EST) found in human cDNA libraries, were selected based on homology to known sulfatase sequences. Based on such EST sequences, primers were
20 designed to identify a full length clone from a human cDNA library. Positive clones were sequenced and the overlapping fragments were assembled. The 22438, 23553, 25278, and 26212 sulfatase amino acid sequences are shown in Figures 1, 5, 10, and 15, respectively, and SEQ ID NOS:1, 3, 5, and 7. The 22438, 23553, 25278, and 26212 sulfatase cDNA sequences are shown in Figures 1, 5, 10, and 15 and SEQ ID NOS:2, 4,
25 6, and 8.

- Analysis of the assembled sequences revealed that the cloned cDNA molecules encoded sulfatase-like polypeptides. BLAST analysis indicated that the 23553 sulfatase is a glucosamine-6-sulfatase, that the 25278 sulfatase is an N-acetylgalactosamine-4-sulfatase, that the 22438 is an arylsulfatase with highest
30 homology to arylsulfatase A and E genes and that the 26212 sulfatase is an arylsulfatase with highest homology to the arylsulfatase B gene (N-acetylgalactosamine-4-sulfatase).

The sulfatase sequences of the invention belong to the sulfatase family of molecules having conserved functional features. The term "family" when referring to the proteins and nucleic acid molecules of the invention is intended to mean two or more proteins or nucleic acid molecules having sufficient amino acid or nucleotide
5 sequence identity as defined herein to provide a specific function. Such family members can be naturally-occurring and can be from either the same or different species. For example, a family can contain a first protein of murine origin and an ortholog of that protein of human origin, as well as a second, distinct protein of human origin and a murine ortholog of that protein.

10 The 22438 sulfatase gene encodes an approximately 2175 nucleotide mRNA transcript having the corresponding cDNA set forth in SEQ ID NO:2. This transcript has an open reading frame which encodes a 525 amino acid protein (SEQ ID NO:1).

The 23553 sulfatase gene encodes an approximately 4321 nucleotide mRNA transcript having the corresponding cDNA set forth in SEQ ID NO:4. This transcript has
15 an open reading frame which encodes an 871 amino acid protein (SEQ ID NO:3).

The 25278 sulfatase gene encodes an approximately 2940 nucleotide mRNA transcript having the corresponding cDNA set forth in SEQ ID NO:6. This transcript has an open reading frame which encodes a 569 amino acid protein (SEQ ID NO:5).

The 26212 sulfatase gene encodes an approximately 2253 nucleotide mRNA
20 transcript having the corresponding cDNA set forth in SEQ ID NO:8. This transcript has an open reading frame which encodes a 599 amino acid protein (SEQ ID NO:7).

Prosites program analysis was used to predict various sites within the 22438 sulfatase protein as shown in Figure 4.

Prosites program analysis was used to predict various sites within the 23553
25 sulfatase protein as shown in Figure 8.

Prosites program analysis was used to predict various sites within the 25278 sulfatase protein as shown in Figure 13.

Prosites program analysis was used to predict various sites within the 26212 sulfatase protein as shown in Figure 18.

30 *In situ* hybridization experiments showed that 22438 is expressed in subpopulations of DRG neurons, spinal cord, and brain, as disclosed hereinabove.

Expression of the 22438 sulfatase mRNA in the above cells and tissues indicates that the sulfatase is likely to be involved in the proper function of and in disorders involving these tissues. Accordingly, the disclosed invention further relates to methods and compositions for the study, modulation, diagnosis and treatment of sulfatase related disorders, especially disorders of these tissues that include, but are not limited to those disclosed herein.

The 23553 sulfatase is differentially expressed in breast and colon cancer and in colonic metastases to the liver. Accordingly, the disclosed invention further relates to methods and compositions for the study, modulation, diagnosis and treatment in these tissues (normal and tumor).

The 25278 sulfatase is differentially expressed in colon tumors and colonic metastases to the liver. Accordingly, the disclosed invention further relates to methods and compositions for the study, modulation, diagnosis and treatment in these normal and tumor tissues.

The 26212 sulfatase is differentially expressed in colon metastases and lung tumors. Accordingly, the disclosed invention further relates to methods and compositions for the study, modulation, diagnosis and treatment in these normal and tumor tissues.

The compositions include sulfatase polypeptides, nucleic acids, vectors, transformed cells and related variants and fragments thereof, as well as agents that modulate expression of the polypeptides and polynucleotides. In particular, the invention relates to the modulation, diagnosis and treatment of sulfatase related disorders as described herein.

Treatment is defined as the application or administration of a therapeutic agent to a patient, or application or administration of a therapeutic agent to an isolated tissue or cell line from a patient, who has a disease, a symptom of disease or a predisposition toward a disease, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve or affect the disease, the symptoms of disease or the predisposition toward disease. "Subject, as used herein, can refer to a mammal, *e.g.* a human, or to an experimental or animal or disease model. The subject can also be a non-human animal, *e.g.* a horse, cow, goat, or other domestic animal. A therapeutic

agent includes, but is not limited to, small molecules, peptides, antibodies, ribozymes and antisense oligonucleotides.

Disorders involving the brain include, but are not limited to, disorders involving neurons, and disorders involving glia, such as astrocytes, oligodendrocytes, 5 ependymal cells, and microglia; cerebral edema, raised intracranial pressure and herniation, and hydrocephalus; malformations and developmental diseases, such as neural tube defects, forebrain anomalies, posterior fossa anomalies, and syringomyelia and hydromyelia; perinatal brain injury; cerebrovascular diseases, such as those related to hypoxia, ischemia, and infarction, including hypotension, hypoperfusion, 10 and low-flow states--global cerebral ischemia and focal cerebral ischemia--infarction from obstruction of local blood supply, intracranial hemorrhage, including intracerebral (intraparenchymal) hemorrhage, subarachnoid hemorrhage and ruptured berry aneurysms, and vascular malformations, hypertensive cerebrovascular disease, including lacunar infarcts, slit hemorrhages, and hypertensive encephalopathy; 15 infections, such as acute meningitis, including acute pyogenic (bacterial) meningitis and acute aseptic (viral) meningitis, acute focal suppurative infections, including brain abscess, subdural empyema, and extradural abscess, chronic bacterial meningoencephalitis, including tuberculosis and mycobacterioses, neurosyphilis, and neuroborreliosis (Lyme disease), viral meningoencephalitis, including arthropod- 20 borne (Arbo) viral encephalitis, *Herpes simplex* virus Type 1, *Herpes simplex* virus Type 2, *Varicella-zoster* virus (*Herpes zoster*), cytomegalovirus, poliomyelitis, rabies, and human immunodeficiency virus 1, including HIV-1 meningoencephalitis (subacute encephalitis), vacuolar myelopathy, AIDS-associated myopathy, peripheral neuropathy, and AIDS in children, progressive multifocal leukoencephalopathy, 25 subacute sclerosing panencephalitis, fungal meningoencephalitis, other infectious diseases of the nervous system; transmissible spongiform encephalopathies (prion diseases); demyelinating diseases, including multiple sclerosis, multiple sclerosis variants, acute disseminated encephalomyelitis and acute necrotizing hemorrhagic encephalomyelitis, and other diseases with demyelination; degenerative diseases, such 30 as degenerative diseases affecting the cerebral cortex, including Alzheimer disease and Pick disease, degenerative diseases of basal ganglia and brain stem, including Parkinsonism, idiopathic Parkinson disease (paralysis agitans), progressive

supranuclear palsy, corticobasal degeneration, multiple system atrophy, including striatonigral degeneration, Shy-Drager syndrome, and olivopontocerebellar atrophy, and Huntington disease; spinocerebellar degenerations, including spinocerebellar ataxias, including Friedreich ataxia, and ataxia-telangiectasia, degenerative diseases affecting motor neurons, including amyotrophic lateral sclerosis (motor neuron disease), bulbospinal atrophy (Kennedy syndrome), and spinal muscular atrophy; inborn errors of metabolism, such as leukodystrophies, including Krabbe disease, metachromatic leukodystrophy, adrenoleukodystrophy, Pelizaeus-Merzbacher disease, and Canavan disease, mitochondrial encephalomyopathies, including Leigh disease and other mitochondrial encephalomyopathies; toxic and acquired metabolic diseases, including vitamin deficiencies such as thiamine (vitamin B₁) deficiency and vitamin B₁₂ deficiency, neurologic sequelae of metabolic disturbances, including hypoglycemia, hyperglycemia, and hepatic encephatopathy, toxic disorders, including carbon monoxide, methanol, ethanol, and radiation, including combined methotrexate and radiation-induced injury; tumors, such as gliomas, including astrocytoma, including fibrillary (diffuse) astrocytoma and glioblastoma multiforme, pilocytic astrocytoma, pleomorphic xanthoastrocytoma, and brain stem glioma, oligodendroglioma, and ependymoma and related paraventricular mass lesions, neuronal tumors, poorly differentiated neoplasms, including medulloblastoma, other parenchymal tumors, including primary brain lymphoma, germ cell tumors, and pineal parenchymal tumors, meningiomas, metastatic tumors, paraneoplastic syndromes, peripheral nerve sheath tumors, including schwannoma, neurofibroma, and malignant peripheral nerve sheath tumor (malignant schwannoma), and neurocutaneous syndromes (phakomatoses), including neurofibromatosis, including Type 1 neurofibromatosis (NF1) and TYPE 2 neurofibromatosis (NF2), tuberous sclerosis, and Von Hippel-Lindau disease.

Furthermore, as disclosed in the background hereinabove, specific disorders have been associated with function of the various sulfatases. Accordingly, the sulfatases disclosed herein, having homology to specific sulfatases as disclosed herein, are useful for diagnosis and treatment of the disorders associated with sulfatase dysfunction as disclosed herein and to modulation of gene expression in the affected tissues.

The sequences of the invention find use in diagnosis of disorders involving an increase or decrease in sulfatase expression relative to normal expression, such as a proliferative disorder, a differentiative disorder, or a developmental disorder. The sequences also find use in modulating sulfatase-related responses. By "modulating" is intended the upregulating or downregulating of a response. That is, the compositions of the invention affect the targeted activity in either a positive or negative fashion.

The invention relates to novel sulfatases, having the deduced amino acid sequence shown in Figures 1, 5, 10, and 15 (SEQ ID NOS:1, 3, 5, and 7) or having the amino acid sequences encoded by the deposited cDNAs, Patent Deposit Numbers _____, PTA-1639, PTA-1846, or _____. The deposited sequences, as well as the polypeptides encoded by the sequences, are incorporated herein by reference and control in the event of any conflict, such as a sequencing error, with description in this application.

Thus, the present invention provides an isolated or purified sulfatase polypeptides and variants and fragments thereof. "Sulfatase polypeptide" or "sulfatase protein" refers to the polypeptide in SEQ ID NOS:1, 3, 5, or 7 or encoded by the deposited cDNAs. The term "sulfatase protein" or "sulfatase polypeptide," however, further includes the numerous variants described herein, as well as fragments derived from the full-length sulfatase and variants.

Sulfatase polypeptides can be purified to homogeneity. It is understood, however, that preparations in which the polypeptide is not purified to homogeneity are useful and considered to contain an isolated form of the polypeptide. The critical feature is that the preparation allows for the desired function of the polypeptide, even in the presence of considerable amounts of other components. Thus, the invention encompasses various degrees of purity.

As used herein, a polypeptide is said to be "isolated" or "purified" when it is substantially free of cellular material when it is isolated from recombinant and non-recombinant cells, or free of chemical precursors or other chemicals when it is chemically synthesized. A polypeptide, however, can be joined to another polypeptide with which it is not normally associated in a cell and still be considered "isolated" or "purified."

In one embodiment, the language "substantially free of cellular material" includes preparations of sulfatase having less than about 30% (by dry weight) other

proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins. When the polypeptide is recombinantly produced, it can also be substantially free of culture medium, i.e., culture medium represents less than about 20%, less than about 10%, or less than about 5% of the volume of the protein preparation.

The sulfatase polypeptide is also considered to be isolated when it is part of a membrane preparation or is purified and then reconstituted with membrane vesicles or liposomes.

The language "substantially free of chemical precursors or other chemicals" includes preparations of the sulfatase polypeptide in which it is separated from chemical precursors or other chemicals that are involved in its synthesis. The language "substantially free of chemical precursors or other chemicals" includes, but is not limited to, preparations of the polypeptide having less than about 30% (by dry weight) chemical precursors or other chemicals, less than about 20% chemical precursors or other chemicals, less than about 10% chemical precursors or other chemicals, or less than about 5% chemical precursors or other chemicals.

In one embodiment, the sulfatase polypeptide comprises the amino acid sequence shown in SEQ ID NOS:1, 3, 5, or 7. However, the invention also encompasses sequence variants. By "variants" is intended proteins or polypeptides having an amino acid sequence that is at least about 45%, 55%, 65%, preferably about 75%, 85%, 95%, or 98% identical to the amino acid sequence of SEQ ID NOS:1,3,5, or 7. Variants also include polypeptides encoded by the cDNA insert of the plasmid deposited with ATCC as Patent Deposit Numbers ____, PTA-1639, PTA-1846, or ____, or polypeptides encoded by a nucleic acid molecule that hybridizes to the nucleic acid molecule of SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14, or a complement thereof, under stringent conditions. In another embodiment, a variant of an isolated polypeptide of the present invention differs, by at least 1, but less than 5, 10, 20, 50, or 100 amino acid residues from the sequence shown in SEQ ID NO:1, 3, 5, or 7. If alignment is needed for this comparison the sequences should be aligned for maximum identity. "Looped" out sequences from deletions or insertions, or mismatches, are considered differences. Such variants generally retain the functional activity of the 22438-like, 23553-like, 25278-like, or 26212-like proteins of the invention. Variants include

polypeptides that differ in amino acid sequence due to natural allelic variation or mutagenesis.

5 Variants include a substantially homologous protein encoded by the same genetic locus in an organism, i.e., an allelic variant. Variants also encompass proteins derived from other genetic loci in an organism, but having substantial homology to the sulfatase of SEQ ID NOS:1, 3, 5, or 7. Variants also include proteins substantially homologous to the sulfatase but derived from another organism, i.e., an ortholog. Variants also include proteins that are substantially homologous to the sulfatase that are produced by chemical synthesis. Variants also include proteins that are substantially homologous to the
10 sulfatase that are produced by recombinant methods. Variants retain the biological activity (for example, sulfatase activity) of the polypeptide set forth by the reference sequence (SEQ ID NOS: 1, 3, 5, or 7). It is understood, however, that variants exclude any amino acid sequences disclosed prior to the invention.

Preferred sulfatase polypeptides of the present invention have an amino acid
15 sequence sufficiently identical to the amino acid sequence of SEQ ID NOS:1, 3, 5, or 7. The term "sufficiently identical" is used herein to refer to a first amino acid or nucleotide sequence that contains a sufficient or minimum number of identical or equivalent (e.g., with a similar side chain) amino acid residues or nucleotides to a second amino acid or nucleotide sequence such that the first and second amino acid or
20 nucleotide sequences have a common structural domain and/or common functional activity. For example, amino acid or nucleotide sequences that contain a common structural domain having at least about 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identity are defined herein as sufficiently identical.

In one embodiment, a variant of the 23553 sulfatase is greater than 92%
25 homologous. In another embodiment, a variant of the 25278 sulfatase is greater than 50% identical. In another embodiment, the 26212 sulfatase is greater than 50% identical.

To determine the percent identity of two amino acid sequences, or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes
30 (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, the length of a

reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, and even more preferably at least 70%, 80%, 90%, 100% of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (1970) *J. Mol. Biol.* 48:444-453 algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. A particularly preferred set of parameters (and the one that should be used if the practitioner is uncertain about what parameters should be applied to determine if a molecule is within a sequence identity or homology limitation of the invention) is using a Blossum 62 scoring matrix with a gap open penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5.

The percent identity between two amino acid or nucleotide sequences can be determined using the algorithm of E. Meyers and W. Miller (1989) *CABIOS* 4:11-17 which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences described herein can be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.* (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

The invention also encompasses polypeptides having a lower degree of identity but having sufficient similarity so as to perform one or more of the same functions performed by the sulfatase. Similarity is determined by conservative amino acid substitution, as shown in Table 1. Such substitutions are those that substitute a given amino acid in a polypeptide by another amino acid of like characteristics. Conservative substitutions are likely to be phenotypically silent. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr, exchange of the acidic residues Asp and Glu, substitution between the amide residues Asn and Gln, exchange of the basic residues Lys and Arg and replacements among the aromatic residues Phe, Tyr. Guidance concerning which amino acid changes are likely to be phenotypically silent are found in Bowie *et al.*, *Science* 247:1306-1310 (1990).

TABLE 1. Conservative Amino Acid Substitutions.

Aromatic	Phenylalanine
	Tryptophan
	Tyrosine
Hydrophobic	Leucine
	Isoleucine
	Valine
Polar	Glutamine
	Asparagine
Basic	Arginine
	Lysine
	Histidine
Acidic	Aspartic Acid
	Glutamic Acid
Small	Alanine
	Serine
	Threonine
	Methionine
	Glycine

A variant polypeptide can differ in amino acid sequence by one or more
5 substitutions, deletions, insertions, inversions, fusions, and truncations or a
combination of any of these. Variant polypeptides can be fully functional or can lack
function in one or more activities. Thus, in the present case, variations can affect the
function, for example, of one or more of regions including a metal (e.g., Ca^{++})-
binding domain, activation domain, sulfatase catalytic domain, the region containing a
10 propeptide, regulatory regions, substrate binding regions, regions involved in
membrane association or subcellular localization, regions involved in post-

translational modification, for example, by phosphorylation, and regions that are important for effector function (i.e., agents that act upon the protein, such as in the conversion of cysteine to 2-amino-3-oxopropionic acid or serine semi-aldehyde).

5 Fully functional variants typically contain only conservative variation or variation in non-critical residues or in non-critical regions. Functional variants can also contain substitution of similar amino acids, which results in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree.

10 Non-functional variants typically contain one or more non-conservative amino acid substitutions, deletions, insertions, inversions, or truncation or a substitution, insertion, inversion, or deletion in a critical residue or critical region.

As indicated, variants can be naturally-occurring or can be made by recombinant means or chemical synthesis to provide useful and novel characteristics for the sulfatase polypeptide. This includes preventing immunogenicity from pharmaceutical
15 formulations by preventing protein aggregation.

Useful variations further include alteration of functional activity. For example, one embodiment involves a variation at the substrate binding site that results in binding but not hydrolysis or more or less hydrolysis of the substrate than wild type. A further useful variation at the same site can result in altered affinity for the substrate. Useful
20 variations also include changes that provide for affinity for another substrate. Useful variations further include the ability to bind an effector molecule with greater or lesser affinity, such as not to bind or to bind but not release it. Further useful variations include alteration in the ability of the propeptide to be cleaved by a cleavage protein, including alteration in the binding or recognition site. Further, the cleavage site can
25 also be modified so that recognition and cleavage are by a different protease. A specific useful variation involves a variation in the ability to be bound or activated by the enzyme that activates the sulfatase by the conversion of cysteine to 2-3-oxopropionic acid or serine semi-aldehyde. Further variation could include a variation in the specificity of metal binding.

30 Another useful variation provides a fusion protein in which one or more domains or subregions are operationally fused to one or more domains, subregions, or motifs from another sulfatase. For example, a transmembrane domain from a protein can be

introduced into the sulfatase such that the protein is anchored in the cell surface.

Other permutations include changing the number of sulfatase domains, and mixing of sulfatase domains from different sulfatase families, so that substrate specificity is altered. Mixing these various domains can allow the formation of novel sulfatase molecules with different host cell, subcellular localization, substrate, and effector molecule (one that acts on the sulfatase) specificity.

The term "substrate" is intended to refer not only to the sulfated substrate that is cleaved by the sulfatase domain, but to refer to any component with which the polypeptide interacts in order to produce an effect on that component or a subsequent biological effect that is a result of interacting with that component. This can include, but is not limited to, for example, interaction with the sulfatase activation enzyme and components involved in the conversion of 3' phosphoadenosine 5' phosphosulfate to adenosine 3' 5' biphosphate.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham *et al.* (1985) *Science* 244:1081-1085). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity, such as peptide bond hydrolysis *in vitro* or related biological activity, such as proliferative activity. Sites that are critical for binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.* (1992) *J. Mol. Biol.* 224:899-904; de Vos *et al.* (1992) *Science* 255:306-312).

The invention thus also includes polypeptide fragments of the sulfatases. Fragments can be derived from the amino acid sequence shown in SEQ ID NOS:1, 3, 5, or 7. However, the invention also encompasses fragments of the variants of the sulfatase polypeptides as described herein. The fragments to which the invention pertains, however, are not to be construed as encompassing fragments that may be disclosed prior to the present invention.

A fragment can comprise at least about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50 or more contiguous amino acids. Fragments can retain one or more of the biological activities of the protein, for example as discussed

above, as well as fragments that can be used as an immunogen to generate sulfatase antibodies.

For example, for the 25278 sulfatase, the invention encompasses amino acid fragments greater than 5 amino acids, particularly from regions up to around
5 nucleotide 450 and beyond around nucleotide 1520. Specific fragments which may be excluded include those that are underlined in Figure 1. However, even in regions between around nucleotide 450 to around nucleotide 1520, fragments include those that are five or greater excluding those which may have been disclosed prior to the present invention.

10 For the 23553 sulfatase, fragments particularly include fragments of 5 amino acids or more up to around nucleotide 670.

For the 26212 sulfatase, for example, fragments containing 5 or more amino acids up to about nucleotide 572 are particularly encompassed by the invention. However, fragments of 5 amino acids or more encoded by around nucleotide 572 to
15 around nucleotide 1985 are also encompassed by the invention with the understanding that such fragments do not encompass those which may have been disclosed prior to the invention. For example, these can include the sections underlined in Figure 15.

Biologically active fragments (peptides which are, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 50, 100 or more amino acids in length) can comprise a
20 functional site. Such sites include but are not limited to those discussed above, such as a catalytic site, regulatory site, site important for substrate recognition or binding, regions containing a sulfatase domain or motif, phosphorylation sites, glycosylation sites, and other functional sites disclosed herein.

Fragments, for example, can extend in one or both directions from the functional
25 site to encompass 5, 10, 15, 20, 30, 40, 50, or up to 100 amino acids. Further, fragments can include sub-fragments of the specific sites or regions disclosed herein, which sub-fragments retain the function of the site or region from which they are derived.

The invention also provides fragments with immunogenic properties. These contain an epitope-bearing portion of the sulfatase polypeptide and variants. These
30 epitope-bearing peptides are useful to raise antibodies that bind specifically to a sulfatase polypeptide or region or fragment. These peptides can contain at least 10, 12, at least 14, or between at least about 15 to about 30 amino acids. The epitope-bearing sulfatase

polypeptides may be produced by any conventional means (Houghten, R.A. (1985) *Proc. Natl. Acad. Sci. USA* 82:5131-5135). Simultaneous multiple peptide synthesis is described in U.S. Patent No. 4,631,211.

Non-limiting examples of antigenic polypeptides that can be used to generate
5 antibodies include but are not limited to peptides derived from extracellular regions. Regions having a high antigenicity index are shown in Figures 3, 7, 12, and 17. However, intracellularly-made antibodies ("intrabodies") are also encompassed, which would recognize intracellular peptide regions.

Fragments can be discrete (not fused to other amino acids or polypeptides) or can
10 be within a larger polypeptide. Further, several fragments can be comprised within a single larger polypeptide. In one embodiment a fragment designed for expression in a host can have heterologous pre- and pro-polypeptide regions fused to the amino terminus of the sulfatase polypeptide fragment and an additional region fused to the carboxyl terminus of the fragment.

15 The invention thus provides chimeric or fusion proteins. These comprise a sulfatase peptide sequence operatively linked to a heterologous peptide having an amino acid sequence not substantially homologous to the sulfatase polypeptide. "Operatively linked" indicates that the sulfatase polypeptide and the heterologous peptide are fused in-frame. The heterologous peptide can be fused to the N-terminus or C-terminus of the
20 sulfatase polypeptide or can be internally located.

In one embodiment the fusion protein does not affect sulfatase function *per se*. For example, the fusion protein can be a GST-fusion protein in which sulfatase sequences are fused to the N- or C-terminus of the GST sequences. Other types of fusion proteins include, but are not limited to, enzymatic fusion proteins, for example
25 beta-galactosidase fusions, yeast two-hybrid GAL4 fusions, poly-His fusions and Ig fusions. Such fusion proteins, particularly poly-His fusions, can facilitate the purification of recombinant sulfatase polypeptide. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence. Therefore, in another embodiment, the fusion protein
30 contains a heterologous signal sequence at its C- or N-terminus.

EP-A-O 464 533 discloses fusion proteins comprising various portions of immunoglobulin constant regions. The Fc is useful in therapy and diagnosis and thus

results, for example, in improved pharmacokinetic properties (EP-A 0232 262). In drug discovery, for example, human proteins have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists (Bennett *et al.* (1995) *J. Mol. Recog.* 8:52-58 (1995) and Johanson *et al. J. Biol. Chem.* 270:9459-9471). Thus, this invention also encompasses soluble fusion proteins containing a sulfatase polypeptide and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclass (IgG, IgM, IgA, IgE). Preferred as immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. For some uses it is desirable to remove the Fc after the fusion protein has been used for its intended purpose, for example when the fusion protein is to be used as antigen for immunizations. In a particular embodiment, the Fc part can be removed in a simple way by a cleavage sequence, which is also incorporated and can be cleaved with factor Xa.

A chimeric or fusion protein can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different protein sequences are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see Ausubel *et al.* (1992) *Current Protocols in Molecular Biology*). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST protein). A sulfatase-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to sulfatase.

Another form of fusion protein is one that directly affects sulfatase functions. Accordingly, a sulfatase polypeptide is encompassed by the present invention in which one or more of the sulfatase regions (or parts thereof) has been replaced by heterologous or homologous regions (or parts thereof) from another sulfatase. Accordingly, various permutations are possible, for example, as discussed above. Thus, chimeric sulfatases can be formed in which one or more of the native domains or subregions has been

1 duplicated, removed, or replaced by another. This includes but is not limited to catalytic
sulfatase or substrate binding domains, and regions involved in activation.

It is understood however that such regions could be derived from a sulfatase that
has not yet been characterized. Moreover, sulfatase function can be derived from
5 peptides that contain these functions but are not in a sulfatase family.

The isolated 22438 sulfatase protein can be purified from cells that naturally
express it, such as DRG neurons, including small and medium sized neurons, spinal
cord, including interneurons and motor neurons, and brain, especially purified from cells
that have been altered to express it (recombinant), or synthesized using known protein
10 synthesis methods.

The isolated 23553 sulfatase protein can be purified from cells that naturally
express it, such as cells from any of the tissues shown in Figures 9 and 21-26, especially
purified from cells that have been altered to express it (recombinant), or synthesized
using known protein synthesis methods.

15 The isolated 25278 sulfatase protein can be purified from cells that naturally
express it, such as cells from any of the tissues shown in Figures 14 and 28, especially
purified from cells that have been altered to express it (recombinant), or synthesized
using known protein synthesis methods.

The isolated 26212 sulfatase protein can be purified from cells that naturally
20 express it, such as cells from any of the tissues shown in Figures 30-36, especially
purified from cells that have been altered to express it (recombinant), or synthesized
using known protein synthesis methods.

In one embodiment, the protein is produced by recombinant DNA techniques.
For example, a nucleic acid molecule encoding the sulfatase polypeptide is cloned into
25 an expression vector, the expression vector introduced into a host cell and the protein
expressed in the host cell. The protein can then be isolated from the cells by an
appropriate purification scheme using standard protein purification techniques.

Polypeptides often contain amino acids other than the 20 amino acids commonly
referred to as the 20 naturally-occurring amino acids. Further, many amino acids,
30 including the terminal amino acids, may be modified by natural processes, such as
processing and other post-translational modifications, or by chemical modification
techniques well known in the art. Common modifications that occur naturally in

polypeptides are described in basic texts, detailed monographs, and the research literature, and they are well known to those of skill in the art.

Accordingly, the polypeptides also encompass derivatives or analogs in which a substituted amino acid residue is not one encoded by the genetic code, in which a
5 substituent group is included, in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or in which the additional amino acids are fused to the mature polypeptide, such as a leader or secretory sequence or a sequence for purification of the mature polypeptide or a pro-protein sequence.

10 Known modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of
15 covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

20 Such modifications are well-known to those of skill in the art and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as *Proteins - Structure and Molecular Properties*, 2nd ed., T.E.
25 Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as by Wold, F., *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York 1-12 (1983); Seifter *et al.* (1990) *Meth. Enzymol.* 182: 626-646) and Rattan *et al.* (1992) *Ann. N.Y. Acad. Sci.* 663:48-62).

30 As is also well known, polypeptides are not always entirely linear. For instance, polypeptides may be branched as a result of ubiquitination, and they may be circular, with or without branching, generally as a result of post-translation events, including

natural processing events and events brought about by human manipulation which do not occur naturally. Circular, branched and branched circular polypeptides may be synthesized by non-translational natural processes and by synthetic methods.

5 Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. Blockage of the amino or carboxyl group in a polypeptide, or both, by a covalent modification, is common in naturally-occurring and synthetic polypeptides. For instance, the aminoterminal residue of polypeptides made in *E. coli*, prior to proteolytic processing, almost invariably will be N-formylmethionine.

10 The modifications can be a function of how the protein is made. For recombinant polypeptides, for example, the modifications will be determined by the host cell posttranslational modification capacity and the modification signals in the polypeptide amino acid sequence. Accordingly, when glycosylation is desired, a polypeptide should be expressed in a glycosylating host, generally a eukaryotic cell.

15 Insect cells often carry out the same posttranslational glycosylations as mammalian cells and, for this reason, insect cell expression systems have been developed to efficiently express mammalian proteins having native patterns of glycosylation. Similar considerations apply to other modifications.

The same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain more than one type of modification.

20

Polypeptide Uses

The protein sequences of the present invention can be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to

25

30 the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped

alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

5 Sulfatase polypeptides are useful for producing antibodies specific for sulfatase, regions, or fragments. Regions having a high antigenicity index score are shown in Figures 3, 7, 12, and 17.

 Sulfatase polypeptides are useful for biological assays related to sulfatases. Such assays involve any of the known sulfatase functions or activities or properties useful for
10 diagnosis and treatment of sulfatase-related conditions, including those in the references cited herein, which are incorporated by reference for these assays, functions, and disorders.

 These assays include, but are not limited to, binding to and/or cleaving specific substrates to produce fragments, steady state levels of sulfated compounds, cysteine
15 modification, and biological assays related to the functions produced by sulfated compounds. Specific substrates useful for assays related to sulfate conjugate hydrolysis include but are not limited to xenobiotics, thyroid hormones, steroids, and catechols. Specific sulfate conjugates include, but are not limited to, 3 α -sulfatolithocholytaurine, sulfate conjugates of estrone, 4-methylumbelliferone, and harmol, sulfated cartilage and
20 proteoglycans, 4-nitrophenol, simple phenols, hydroxyarylamines, iodothyronines, catecholamines, 1-naphthyl, salbutamol, estrogens, ethinylestradiol, equilenin, diethylstilbestrol, androgens, cholesterol bile salts, pregnenolone, benzylic alcohols, glycolipidsulfates, complex carbohydrates such as dermatan and chondrotin sulfate, steroid sulfate, sulfate conjugates of xenobiotics, cholesterol sulfate, xenobiotic phenyls,
25 *o*-cresol, vanillin, eugenol, *m*-cresol, thymol, ethyl-4,4-dihydroxybenzoate, *p*-cresol, sesamol, methyl-2,6-dihydroxy-4-methylbenzyloate, methyl-2,4-dihydroxybenzoate, methyl-3,5-dihydroxybenzoate, tyramine, dopamine, 5 hydroxytryptamine, pyrogallol, 4-nitrocatecholsulfate, estrone sulfate, metabolites of the cytochrome P450 mono-
30 oxygenase system, dihydroepiandrosterone sulfate (DHEAS), minoxidil, cicletanine, sulfated mutagens and carcinogens, such as aromatic amines (including heterocyclic amines), and benzylic alcohols of chemicals such as polycyclic aromatic hydrocarbons, saffrole and estragole, glycosaminoglycans, sulfolipids, betahydroxysteroids, sulfate

esters of chromogenic or fluorogenic aromatic compounds, cerebroside sulfate, keratan sulfate, and heparan sulfate. Substrates also include any in the references cited herein, which are incorporated herein by reference for these substrates. Accordingly the assays include, but are not limited to, these sulfated substrates and biological effects of sulfation or desulfation of these substrates and associated biochemical, cellular, or phenotypic effects of sulfation of desulfation, and any of the other biological or functional properties of these proteins, including, but not limited to, those disclosed herein, and in any reference cited herein which is incorporated herein by reference for the disclosure of these properties and for the assays based on these properties. Further, assays may relate to changes in the protein, *per se*, and on the effects of these changes, for example, activation of the sulfatase by modification of a cysteine residue as disclosed herein, cleavage of the propeptide by a proteinase, induction of expression of the protein *in vivo*, inhibition of function, as well as any other effects on the protein mentioned herein or cited in any reference herein, which are incorporated herein by reference for these effects and for the subsequent biological consequences of these effects.

Sulfatase polypeptides are also useful in drug screening assays, in cell-based or cell-free systems. Cell-based systems can be native, i.e., cells that normally express sulfatase, such as those discussed above, especially tumor cells, as a biopsy, or expanded in cell culture. In one embodiment, however, cell-based assays involve recombinant host cells expressing sulfatase. Accordingly, these drug-screening assays can be based on effects on protein function as described above for biological assays useful for diagnosis and treatment.

Determining the ability of the test compound to interact with a sulfatase can also comprise determining the ability of the test compound to preferentially bind to the polypeptide as compared to the ability of a known binding molecule to bind to the polypeptide.

The polypeptides can be used to identify compounds that modulate sulfatase activity. Such compounds, for example, can increase or decrease affinity or rate of binding to substrate, compete with substrate for binding to sulfatase, or displace substrate bound to sulfatase. Both sulfatase and appropriate variants and fragments can be used in high-throughput screens to assay candidate compounds for the ability to bind to sulfatase. These compounds can be further screened against a functional sulfatase to

determine the effect of the compound on sulfatase activity. Compounds can be identified that activate (agonist) or inactivate (antagonist) sulfatase to a desired degree. Modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject).

5 Sulfatase polypeptides can be used to screen a compound for the ability to stimulate or inhibit interaction between sulfatase protein and a target molecule that normally interacts with the sulfatase, for example, substrate of the sulfatase domain. The assay includes the steps of combining sulfatase protein with a candidate compound under conditions that allow the sulfatase protein or fragment to interact
10 with the target molecule, and to detect the formation of a complex between the sulfatase protein and the target or to detect the biochemical consequence of the interaction with the sulfatase and the target.

Determining the ability of the sulfatase to bind to a target molecule can also be accomplished using a technology such as real-time Bimolecular Interaction Analysis
15 (BIA). Sjolander *et al.* (1991) *Anal. Chem.* 63:2338-2345 and Szabo *et al.* (1995) *Curr. Opin. Struct. Biol.* 5:699-705. As used herein, "BIA" is a technology for studying biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore™). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological
20 molecules.

The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-
25 compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to polypeptide libraries, while the other four approaches are applicable to polypeptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997) *Anticancer Drug Des.* 12:145).

30 Examples of methods for the synthesis of molecular libraries can be found in the art, for example in DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:6909; Erb *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994). *J. Med.*

Chem. 37:2678; Cho *et al.* (1993) *Science* 261:1303; Carell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in Gallop *et al.* (1994) *J. Med. Chem.* 37:1233. Libraries of compounds may be presented in solution (e.g., Houghten (1992) *Biotechniques* 13:412-421), or on
5 beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirla *et al.* (1990) *Proc. Natl. Acad. Sci.* 97:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-
10 310); (Ladner *supra*).

Candidate compounds include, for example, 1) peptides such as soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, e.g., Lam *et al.* (1991) *Nature* 354:82-84; Houghten *et al.* (1991) *Nature* 354:84-86) and combinatorial chemistry-derived molecular libraries made of D- and/or L- configuration
15 amino acids; 2) phosphopeptides (e.g., members of random and partially degenerate, directed phosphopeptide libraries, see, e.g., Songyang *et al.* (1993) *Cell* 72:767-778); 3) antibodies (e.g., polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, and single chain antibodies as well as Fab, F(ab')₂, Fab expression library fragments, and epitope-binding fragments of antibodies); 4) small organic and inorganic molecules (e.g.,
20 molecules obtained from combinatorial and natural product libraries); substrate analogs including, but not limited to, substrates disclosed herein.

One candidate compound is a soluble full-length sulfatase or fragment that competes for substrate. Other candidate compounds include mutant sulfatases or appropriate fragments containing mutations that affect sulfatase function and compete
25 for substrate. Accordingly, a fragment that competes for substrate, for example with a higher affinity, or a fragment that binds substrate but does not process or otherwise affect it, is encompassed by the invention.

The invention provides other end points to identify compounds that modulate (stimulate or inhibit) sulfatase activity. The assays typically involve an assay of cellular
30 events that indicate sulfatase activity. Thus, the expression of genes that are up- or down-regulated in response to sulfatase activity can be assayed. In one embodiment, the regulatory region of such genes can be operably linked to a marker that is easily

detectable, such as luciferase. Alternatively, modification of the sulfatase could also be measured.

Any of the biological or biochemical functions mediated by the sulfatase can be used as an endpoint assay. These include any of the biochemical or
5 biochemical/biological events described herein, in any reference cited herein, incorporated by reference for these endpoint assay targets, and other functions known to those of ordinary skill in the art. Specific end points can include, but are not limited to, the events resulting from expression (or lack thereof) of sulfatase activity. With respect to disorders, this would include, but not be limited to, effects on function,
10 differentiation, and proliferation, which can be assayed, as well as the biological effects of function, such as disorders discussed hereinabove and in the references cited hereinabove which are incorporated herein by reference for the disorders disclosed in those references and other disorders and pathology. In the case of the 22438 sulfatase, models of pain can be used as an end point. In the case of the 23553
15 and 25278 sulfatases, tumor progression can be used as an end point. In the case of the 26212 sulfatase, tumor angiogenesis and/or tumor progression can be used as an end point.

Binding and/or activating compounds can also be screened by using chimeric sulfatase proteins in which one or more regions, segments, sites, and the like, as
20 disclosed herein, or parts thereof, can be replaced by heterologous and homologous counterparts derived from other sulfatases. For example, a catalytic region can be used that interacts with a different substrate specificity and/or affinity than the native sulfatase. Accordingly, a different set of components is available as an end-point assay for activation. As a further alternative, the site of modification by an effector protein, for
25 example, activation or phosphorylation, can be replaced with the site for a different effector protein. Activation can also be detected by a reporter gene containing an easily detectable coding region operably linked to a transcriptional regulatory sequence that is part of the native pathway in which sulfatase is involved.

Sulfatase polypeptides are also useful in competition binding assays in methods
30 designed to discover compounds that interact with the sulfatase. Thus, a compound is exposed to a sulfatase polypeptide under conditions that allow the compound to bind or to otherwise interact with the polypeptide. Soluble sulfatase polypeptide is also added to

the mixture. If the test compound interacts with the soluble sulfatase polypeptide, it decreases the amount of complex formed or activity from the sulfatase target. This type of assay is particularly useful in cases in which compounds are sought that interact with specific regions of the sulfatase. Thus, the soluble polypeptide that competes with the target sulfatase region is designed to contain peptide sequences corresponding to the region of interest.

Another type of competition-binding assay can be used to discover compounds that interact with specific functional sites. As an example, bindable substrate analog and a candidate compound can be added to a sample of the sulfatase. Compounds that interact with the sulfatase at the same site as the substrate or analog will reduce the amount of complex formed between the sulfatase and the substrate or analog. Accordingly, it is possible to discover a compound that specifically prevents interaction between the sulfatase and the component. Another example involves adding a candidate compound to a sample of sulfatase and cleavable substrate. A compound that competes with the substrate will reduce the amount of hydrolysis or binding of the substrate to the sulfatase. Accordingly, compounds can be discovered that directly interact with the sulfatase and compete with the substrate. Such assays can involve any other component that interacts with the sulfatase.

To perform cell free drug screening assays, it is desirable to immobilize either sulfatase, or fragment, or its target molecule to facilitate separation of complexes from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay.

Techniques for immobilizing proteins on matrices can be used in the drug screening assays. In one embodiment, a fusion protein can be provided which adds a domain that allows the protein to be bound to a matrix. For example, glutathione-S-transferase/sulfatase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the cell lysates (e.g., ³⁵S-labeled) and the candidate compound, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads are washed to remove any unbound label, and the matrix immobilized and radiolabel determined directly, or in the supernatant after the complexes is dissociated. Alternatively, the

complexes can be dissociated from the matrix, separated by SDS-PAGE, and the level of sulfatase-binding protein found in the bead fraction quantitated from the gel using standard electrophoretic techniques. For example, either the polypeptide or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin using techniques well known in the art. Alternatively, antibodies reactive with the protein but which do not interfere with binding of the protein to its target molecule can be derivatized to the wells of the plate, and the protein trapped in the wells by antibody conjugation. Preparations of a sulfatase-binding target component, such as substrate or activating enzyme, and a candidate compound are incubated in sulfatase-presenting wells and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the sulfatase target molecule, or which are reactive with the sulfatase and compete with the target molecule; as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the target molecule.

Modulators of sulfatase activity identified according to these drug screening assays can be used to treat a subject with a disorder related to the sulfatase, by treating cells that express the sulfatase. These methods of treatment include the steps of administering the modulators of sulfatase activity in a pharmaceutical composition as described herein, to a subject in need of such treatment.

The 23553, 25278, and 26212 sulfatases are differentially expressed in tumor cells as disclosed herein. Accordingly, these sulfatases are relevant to these disorders and relevant as well to differentiation, function, and growth of the tissues giving rise to the tumors. The 22438 sulfatase is expressed as described above, and accordingly is relevant for disorders involving these tissues. Disorders include, but are not limited to, those discussed hereinabove. Moreover, since the gene is expressed in the central nervous system, this sulfatase is relevant for the treatment of pain.

Sulfatase polypeptides are thus useful for treating a sulfatase-associated disorder characterized by aberrant expression or activity of a sulfatase. "Aberrant expression" or "misexpression", as used herein, refers to a non-wild type pattern of gene expression, at the RNA or protein level. It includes: expression at non-wild type levels, i.e., over or under expression; a pattern of expression that differs from wild

type in terms of the time or stage at which the gene is expressed, e.g., increased or decreased expression (as compared with wild type) at a predetermined developmental period or stage; a pattern of expression that differs from wild type in terms of decreased expression (as compared with wild type) in a predetermined cell type or tissue type; a pattern of expression that differs from wild type in terms of the splicing size, amino acid sequence, post-translational modification, or biological activity of the expressed polypeptide; a pattern of expression that differs from wild type in terms of the effect of an environmental stimulus or extracellular stimulus on expression of the gene, e.g., a pattern of increased or decreased expression (as compared with wild type) in the presence of an increase or decrease in the strength of the stimulus.

In one embodiment, the method involves administering an agent (e.g., an agent identified by a screening assay described herein), or combination of agents that modulates (e.g., upregulates or downregulates) expression or activity of the protein. In another embodiment, the method involves administering sulfatase as therapy to compensate for reduced or aberrant expression or activity of the protein.

Methods for treatment include but are not limited to the use of soluble sulfatase or fragments of sulfatase protein that compete for substrate or any other component that directly interacts with sulfatase, or any of the enzymes that modify the sulfatase. These sulfatases or fragments can have a higher affinity for the target so as to provide effective competition.

Stimulation of activity is desirable in situations in which the protein is abnormally downregulated and/or in which increased activity is likely to have a beneficial effect. Likewise, inhibition of activity is desirable in situations in which the protein is abnormally upregulated and/or in which decreased activity is likely to have a beneficial effect. In one example of such a situation, a subject has a disorder characterized by aberrant development or cellular differentiation. In another example, the subject has a disorder characterized by an aberrant hematopoietic response. In another example, it is desirable to achieve tissue regeneration in a subject.

In yet another aspect of the invention, the proteins of the invention can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; Iwabuchi *et*

al. (1993) *Oncogene* 8:1693-1696; and Brent WO 94/10300), to identify other proteins (captured proteins) which bind to or interact with the proteins of the invention and modulate their activity.

Sulfatase polypeptides also are useful to provide a target for diagnosing a disease or predisposition to disease mediated by the sulfatase, including, but not limited to, those diseases disclosed herein, in the references cited herein, and as disclosed above in the background. Accordingly, methods are provided for detecting the presence, or levels of the sulfatase in a cell, tissue, or organism. The method involves contacting a biological sample with a compound capable of interacting with the sulfatase such that the interaction can be detected. One agent for detecting a sulfatase is an antibody capable of selectively binding to the sulfatase. A biological sample includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject.

The sulfatase also provides a target for diagnosing active disease, or predisposition to disease, in a patient having a variant sulfatase. Thus, sulfatase can be isolated from a biological sample and assayed for the presence of a genetic mutation that results in an aberrant protein. This includes amino acid substitution, deletion, insertion, rearrangement, (as the result of aberrant splicing events), and inappropriate post-translational modification. Analytic methods include altered electrophoretic mobility, altered tryptic peptide digest, altered sulfatase activity in cell-based or cell-free assays, such as by alteration in substrate binding or degradation, or ability to be activated by the activation enzyme, or antibody-binding pattern, altered isoelectric point, direct amino acid sequencing, and any other of the known assay techniques useful for detecting mutations in a protein in general or in a sulfatase specifically, such as are disclosed herein.

In vitro techniques for detection of sulfatase include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. Alternatively, the protein can be detected *in vivo* in a subject by introducing into the subject a labeled anti-sulfatase antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods, which detect

the allelic variant of sulfatase expressed in a subject, and methods, which detect fragments of sulfatase in a sample.

Sulfatase polypeptides are also useful in pharmacogenomic analysis.

Pharmacogenomics deal with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, 5 e.g., Eichelbaum, M. (1996) *Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985, and Linder, M.W. (1997) *Clin. Chem.* 43(2):254-266. The clinical outcomes of these variations result in severe toxicity of therapeutic drugs in certain individuals or therapeutic failure of drugs in certain individuals as a result of individual variation in 10 metabolism. Thus, the genotype of the individual can determine the way a therapeutic compound acts on the body or the way the body metabolizes the compound. Further, the activity of drug metabolizing enzymes affects both the intensity and duration of drug action. Thus, the pharmacogenomics of the individual permit the selection of effective compounds and effective dosages of such compounds for prophylactic or therapeutic 15 treatment based on the individual's genotype. The discovery of genetic polymorphisms in some drug metabolizing enzymes has explained why some patients do not obtain the expected drug effects, show an exaggerated drug effect, or experience serious toxicity from standard drug dosages. Polymorphisms can be expressed in the phenotype of the extensive metabolizer and the phenotype of the poor metabolizer. Accordingly, genetic 20 polymorphism may lead to allelic protein variants of sulfatase in which one or more of sulfatase functions in one population is different from those in another population. The polypeptides thus allow a target to ascertain a genetic predisposition that can affect treatment modality. Thus, in a peptide-based treatment, polymorphism may give rise to catalytic regions that are more or less active. Accordingly, dosage would necessarily be 25 modified to maximize the therapeutic effect within a given population containing the polymorphism. As an alternative to genotyping, specific polymorphic polypeptides could be identified.

Sulfatase polypeptides are also useful for monitoring therapeutic effects during clinical trials and other treatment. Thus, the therapeutic effectiveness of an agent that is 30 designed to increase or decrease gene expression, protein levels or sulfatase activity can be monitored over the course of treatment using sulfatase polypeptides as an end-point target. The monitoring can be, for example, as follows: (i) obtaining a pre-

administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression or activity of the protein in the pre-administration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the protein in the post-administration samples; (v) comparing the level of expression or activity of the protein in the pre-administration sample with the protein in the post-administration sample or samples; and (vi) increasing or decreasing the administration of the agent to the subject accordingly.

10 Antibodies

The invention also provides antibodies that selectively bind to the sulfatase and its variants and fragments. An antibody is considered to selectively bind, even if it also binds to other proteins that are not substantially homologous with the sulfatase. These other proteins share homology with a fragment or domain of sulfatase. This conservation in specific regions gives rise to antibodies that bind to both proteins by virtue of the homologous sequence. In this case, it would be understood that antibody binding to the sulfatase is still selective.

Antibodies can be polyclonal or monoclonal. An intact antibody, or a fragment thereof (e.g. Fab or F(ab')₂) can be used. An appropriate immunogenic preparation can be derived from native, recombinantly expressed, or chemically synthesized peptides.

To generate antibodies, an isolated sulfatase polypeptide is used as an immunogen to generate antibodies using standard techniques for polyclonal and monoclonal antibody preparation. Either the full-length protein or antigenic peptide fragment can be used. Regions having a high antigenicity index are disclosed hereinabove.

Antibodies are preferably prepared from these regions or from discrete fragments in these regions. However, antibodies can be prepared from any region of the peptide as described herein. A preferred fragment produces an antibody that diminishes or completely prevents substrate hydrolysis or binding. Antibodies can be developed against the entire sulfatase or domains of the sulfatase as described herein, for example, the substrate binding region, sulfatase motif, or subregions thereof.

Antibodies can also be developed against other specific functional sites as disclosed herein.

The antigenic peptide can comprise a contiguous sequence of at least 12, 14, 15-20, 20-25, or 25-30 or more amino acid residues. In one embodiment, fragments
5 correspond to regions that are located on the surface of the protein, e.g., hydrophilic regions. These fragments are not to be construed, however, as encompassing any fragments, which may be disclosed prior to the invention.

Detection can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes,
10 prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein,
15 fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

20 Antibody Uses

The antibodies can be used to isolate a sulfatase by standard techniques, such as affinity chromatography or immunoprecipitation. The antibodies can facilitate the purification of the natural sulfatase from cells and recombinantly produced sulfatase expressed in host cells.

25 The antibodies are useful to detect the presence of a sulfatase in cells or tissues to determine the pattern of expression of the sulfatase among various tissues in an organism and over the course of normal development. The antibodies can be used to detect a sulfatase *in situ*, *in vitro*, or in a cell lysate or supernatant in order to evaluate the abundance and pattern of expression. Antibody detection of circulating fragments of the
30 full length sulfatase can be used to identify sulfatase turnover. In addition, the antibodies can be used to assess abnormal tissue distribution or abnormal expression during development.

Further, the antibodies can be used to assess sulfatase expression in disease states such as in active stages of the disease or in an individual with a predisposition toward disease related to sulfatase function. When a disorder is caused by an inappropriate tissue distribution, developmental expression, or level of expression of sulfatase protein, the antibody can be prepared against the normal sulfatase protein. If a disorder is characterized by a specific mutation in sulfatase, antibodies specific for this mutant protein can be used to assay for the presence of the specific mutant sulfatase. However, intracellularly-made antibodies ("intrabodies") are also encompassed, which would recognize intracellular sulfatase peptide regions.

The antibodies can also be used to assess normal and aberrant subcellular localization of cells in the various tissues in an organism. Antibodies can be developed against the whole sulfatase or portions of the sulfatase.

The diagnostic uses can be applied, not only in genetic testing, but also in monitoring a treatment modality. Accordingly, where treatment is ultimately aimed at correcting sulfatase expression level or the presence of aberrant sulfatases and aberrant tissue distribution or developmental expression, antibodies directed against the sulfatase or relevant fragments can be used to monitor therapeutic efficacy.

Additionally, antibodies are useful in pharmacogenomic analysis. Thus, antibodies prepared against polymorphic sulfatase can be used to identify individuals that require modified treatment modalities.

The antibodies are also useful as diagnostic tools as an immunological marker for aberrant sulfatase analyzed by electrophoretic mobility, isoelectric point, tryptic peptide digest, and other physical assays known to those in the art.

The antibodies are also useful for tissue typing. Thus, where a specific sulfatase has been correlated with expression in a specific tissue, antibodies that are specific for this sulfatase can be used to identify a tissue type.

The antibodies are also useful in forensic identification. Accordingly, where an individual has been correlated with a specific genetic polymorphism resulting in a specific polymorphic protein, an antibody specific for the polymorphic protein can be used as an aid in identification.

The antibodies are also useful for inhibiting sulfatase function, for example, substrate binding, or sulfatase activity. For example, sulfatase activity may be measured

by the ability to form a binding complex with a sulfated conjugate, such as disclosed herein.

These uses can also be applied in a therapeutic context in which treatment involves inhibiting sulfatase function. An antibody can be used, for example, to block
5 substrate binding. Antibodies can be prepared against specific fragments containing sites required for function or against intact sulfatase associated with a cell.

Completely human antibodies are particularly desirable for therapeutic treatment of human patients. For an overview of this technology for producing human antibodies, see Lonberg *et al.* (1995) *Int. Rev. Immunol.* 13:65-93. For a detailed
10 discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, e.g., U.S. Patent 5,625,126; U.S. Patent 5,633,425; U.S. Patent 5,569,825; U.S. Patent 5,661,016; and U.S. Patent 5,545,806.

The invention also encompasses kits for using antibodies to detect the presence
15 of a sulfatase protein in a biological sample. The kit can comprise antibodies such as a labeled or labelable antibody and a compound or agent for detecting the sulfatase in a biological sample; means for determining the amount of sulfatase in the sample; and means for comparing the amount of sulfatase in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further
20 comprise instructions for using the kit to detect the sulfatase.

Polynucleotides

The nucleotide sequences in SEQ ID NOS:2, 4, 6, and 8 were obtained by sequencing the deposited human cDNAs. Accordingly, the sequences of the deposited
25 clones are controlling as to any discrepancies between the two and any reference to a sequence of SEQ ID NOS:2, 4, 6, or 8, includes reference to the sequence of the deposited cDNA.

The specifically disclosed cDNA comprises the coding region and 5' and 3' untranslated sequences in SEQ ID NOS:2, 4, 6, or 8. The coding sequences of the
30 cDNA's are set forth in SEQ ID NOS:11, 12, 13, and 14.

The invention provides isolated polynucleotides encoding the novel sulfatases. The term "sulfatase polynucleotide" or "sulfatase nucleic acid" refers to the sequences

shown in SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14, or in the deposited cDNAs. The term "sulfatase polynucleotide" or "sulfatase nucleic acid" further includes variants and fragments of sulfatase polynucleotides.

Generally, nucleotide sequence variants of the invention will have at least
5 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to one of the nucleotide sequences disclosed herein.

An "isolated" sulfatase nucleic acid is one that is separated from other nucleic acid present in the natural source of sulfatase nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank sulfatase nucleic acid (i.e.,
10 sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. However, there can be some flanking nucleotide sequences, for example up to about 5KB. The important point is that the sulfatase nucleic acid is isolated from flanking sequences such that it can be subjected to the specific manipulations described herein, such as recombinant expression, preparation
15 of probes and primers, and other uses specific to the sulfatase nucleic acid sequences. In one embodiment, the sulfatase nucleic acid comprises only the coding region.

Moreover, an "isolated" nucleic acid molecule, such as a cDNA or RNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when
20 chemically synthesized. However, the nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered isolated.

In some instances, the isolated material will form part of a composition (for example, a crude extract containing other substances), buffer system or reagent mix. In other circumstances, the material may be purified to essential homogeneity, for
25 example as determined by PAGE or column chromatography such as HPLC. Preferably, an isolated nucleic acid comprises at least about 50, 80 or 90% (on a molar basis) of all macromolecular species present.

For example, recombinant DNA molecules contained in a vector are considered isolated. Further examples of isolated DNA molecules include recombinant DNA
30 molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the isolated DNA molecules of the present invention. Isolated nucleic acid

molecules according to the present invention further include such molecules produced synthetically.

In some instances, the isolated material will form part of a composition (or example, a crude extract containing other substances), buffer system or reagent mix. In other circumstances, the material may be purified to essential homogeneity, for example as determined by PAGE or column chromatography such as HPLC. Preferably, an isolated nucleic acid comprises at least about 50, 80 or 90% (on a molar basis) of all macromolecular species present.

Sulfatase polynucleotides can encode the mature protein plus additional amino or carboxyterminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, facilitate protein trafficking, prolong or shorten protein half-life or facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in situ*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

Sulfatase polynucleotides include, but are not limited to, the sequence encoding the mature polypeptide alone, the sequence encoding the mature polypeptide and additional coding sequences, such as a leader or secretory sequence (e.g., a pre-pro or pro-protein sequence), the sequence encoding the mature polypeptide, with or without the additional coding sequences, plus additional non-coding sequences, for example introns and non-coding 5' and 3' sequences such as transcribed but non-translated sequences that play a role in transcription, mRNA processing (including splicing and polyadenylation signals), ribosome binding and stability of mRNA. In addition, the polynucleotide may be fused to a marker sequence encoding, for example, a peptide that facilitates purification.

Sulfatase polynucleotides can be in the form of RNA, such as mRNA, or in the form DNA, including cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The nucleic acid, especially DNA, can be double-stranded or single-stranded. Single-stranded nucleic acid can be the coding strand (sense strand) or the non-coding strand (anti-sense strand).

The invention further provides variant sulfatase polynucleotides, and fragments thereof, that differ from the nucleotide sequence shown in SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14 due to degeneracy of the genetic code and thus encode the same protein as that encoded by a nucleotide sequence shown in SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14.

Alternatively, a nucleic acid molecule that is a fragment of a 22438-like nucleotide sequence of the present invention comprises a nucleotide sequence consisting of nucleotides 1-100, 100-200, 200-300, 300-400, 400-500, 500-600, 600-700, 700-900, 900-1000, 1000-1100, 1100-1200, 1200-1300, 1300-1400, 1400-1500, 1500-1600, 1600-1700, 1700-1800, 1800-1900, 1900-2000, 2000-2100, 2100-2175 of SEQ ID NO:2.

A nucleic acid molecule that is a fragment of a 23553-like nucleotide sequence of the present invention comprises a nucleotide sequence consisting of nucleotides 1-100, 100-200, 200-300, 300-400, 400-500, 500-600, 600-700, 700-900, 900-1000, 1000-1100, 1100-1200, 1200-1300, 1300-1400, 1400-1500, 1500-1600, 1600-1700, 1700-1800, 1800-1900, 1900-2000, 2000-2100, 2100-2200, 2200-2300, 2300-2400, 2400-2500, 2500-2600, 2600-2700, 2700-2800, 2800-2900, 2900-3000, 3000-3100, 3100-3200, 3200-3300, 3300-3400, 3400-3500, 3500-3600, 3600-3700, 3700-3800, 3800-3900, 3900-4000, 4000-4100, 4100-4200, 4200-4300, 4300-4321 of SEQ ID NO:4.

A nucleic acid molecule that is a fragment of a 25278-like nucleotide sequence of the present invention comprises a nucleotide sequence consisting of nucleotides 1-100, 100-200, 200-300, 300-400, 400-500, 500-600, 600-700, 700-900, 900-1000, 1000-1100, 1100-1200, 1200-1300, 1300-1400, 1400-1500, 1500-1600, 1600-1700, 1700-1800, 1800-1900, 1900-2000, 2000-2100, 2100-2200, 2200-2300, 2300-2400, 2400-2500, 2500-2600, 2600-2700, 2700-2800, 2800-2900, 2900-2940 of SEQ ID NO:6.

A nucleic acid molecule that is a fragment of a 26212-like nucleotide sequence of the present invention comprises a nucleotide sequence consisting of nucleotides 1-100, 100-200, 200-300, 300-400, 400-500, 500-600, 600-700, 700-900, 900-1000, 1000-1100, 1100-1200, 1200-1300, 1300-1400, 1400-1500, 1500-1600, 1600-1700, 1700-1800, 1800-1900, 1900-2000, 2000-2100, 2100-2200, 2200-2253 of SEQ ID NO:8.

The invention also provides sulfatase nucleic acid molecules encoding the variant polypeptides described herein. Such polynucleotides may be naturally occurring, such as allelic variants (same locus), homologs (different locus), and orthologs (different organism), or may be constructed by recombinant DNA methods or by chemical
5 synthesis. Such non-naturally occurring variants may be made by mutagenesis techniques, including those applied to polynucleotides, cells, or organisms. Accordingly, as discussed above, the variants can contain nucleotide substitutions, deletions, inversions and insertions.

Typically, variants have a substantial identity with a nucleic acid molecules of
10 SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14, and the complements thereof. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions.

Orthologs, homologs, and allelic variants can be identified using methods well known in the art. These variants comprise a nucleotide sequence encoding a sulfatase
15 that is typically at least about 40-45%, 45-50%, 50-55%, 55-60%, 60-65%, 65-70%, 70-75%, more typically at least about 75-80% or 80-85%, and most typically at least about 85-90% or 90-95% or more homologous to the nucleotide sequence shown in SEQ ID NOS:2, 4, 6 or 8, or a fragment of this sequence. Such nucleic acid molecules can readily be identified as being able to hybridize under stringent conditions, to the
20 nucleotide sequence shown in SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14, or a fragment of the sequence.

In the case of the 23553 sulfatase, in one embodiment, a variant is greater than 65% homologous with respect to nucleotide sequence. For the 25278 sulfatase, in one embodiment, a variant is greater than 50-60% homologous with respect to nucleotide
25 sequence. With respect to the 26212 sulfatase, in one embodiment, a variant is greater than about 65-75% homologous with respect to nucleotide sequence.

It is understood that stringent hybridization does not indicate substantial homology where it is due to general homology, such as polyA⁺ sequences, or sequences common to all or most proteins, sulfatases, arylsulfatases, glucosamine-6-sulfatases, N-acetylgalactosamine-4-sulfatases, or any of the sulfatases to which the sulfatases of the
30 present invention have shown homology by BLAST analysis, for example, regions to arylsulfatases A, B, C, D, E, F, IDS, and the like. Moreover, it is understood that

variants do not include any of the nucleic acid sequences that may have been disclosed prior to the invention.

As used herein, the term "hybridizes under stringent conditions" describes conditions for hybridization and washing. Stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology* John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Aqueous and nonaqueous methods are described in that reference and either can be used. A preferred, example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 50°C. Another example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 55°C. A further example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 60°C. Preferably, stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 65°C. Particularly preferred stringency conditions (and the conditions that should be used if the practitioner is uncertain about what conditions should be applied to determine if a molecule is within a hybridization limitation of the invention) are 0.5M Sodium Phosphate, 7% SDS at 65°C, followed by one or more washes at 0.2X SSC, 1% SDS at 65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14 corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (e.g., encodes a natural protein).

The present invention also provides isolated nucleic acids that contain a single or double stranded fragment or portion that hybridizes under stringent conditions to the nucleotide sequence of SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14, or the complements of SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14. In one embodiment, the nucleic acid consists of a portion of a nucleotide sequence of SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14 and the complements. The nucleic acid fragments of the invention

are at least about 10-15, preferably at least about 15-20 or 20-25 contiguous nucleotides, and can be 30, 33, 35, 40, 50, 60, 70, 75, 80, 90, 100, 200, 500 or more nucleotides in length. Longer fragments, for example, 600 or more nucleotides in length, which encode antigenic proteins or polypeptides described herein are also
5 useful.

In the case of the 23553 sulfatase, in one embodiment, fragments are derived from nucleotide 1 to about nucleotide 670 and comprise 5-10 and 10-20 contiguous base pairs, and particularly greater than 18. For this sulfatase, in another
10 embodiment, a fragment is derived from around nucleotide 3008 to 3514 and comprises around 5-10 and 10-20 contiguous nucleotides. In other embodiments for this sulfatase, a fragment is derived from around nucleotide 3994 to 4321 and is about 5-10 or 10-20 contiguous nucleotides. For the 25278, in one embodiment, a fragment is derived from around nucleotide 130 to around nucleotide 454 and comprises a
15 contiguous sequence of about 5-10 or 10-20 nucleotides. In another embodiment, the fragment is derived from around nucleotide 454 to around nucleotide 1400 and comprises around 5-10 or 10-20 contiguous nucleotides, especially a fragment greater than 17 nucleotides. In another embodiment the fragment is derived from around nucleotide 1400 to around nucleotide 1850 and comprises a continuous sequence of
20 around 5-10, 10-20, or 20-25 nucleotides, especially a fragment greater than 23 nucleotides. In another embodiment, a fragment is derived from about nucleotide 1933 to about nucleotide 2421. Such a fragment comprises around 5-10 or 10-20 contiguous nucleotides. For the 26212 sulfatase, in one embodiment, a fragment is derived from around nucleotide 272 to around nucleotide 538 and comprises a
25 contiguous sequence of around 5-10 or 10-20 nucleotides, especially a fragment greater than 17 nucleotides. In another embodiment, the fragment is derived from around nucleotide 538 to around nucleotide 751 and comprises a contiguous sequence of at least 5-10 or 10-20 nucleotides, especially greater than 12 nucleotides. In
another embodiment, the fragment is derived from around nucleotide 1074 to around 1551 and comprises a contiguous nucleotide sequence of around 5-10, 10-20, or 20-
30 30, especially greater than 20 nucleotides. In a further embodiment, the fragment is derived from around nucleotide 2052 to 2251 and comprises a contiguous sequence of 5-10 and 10-20 nucleotides, especially fragments greater than 18 nucleotides.

The fragment can comprise DNA or RNA and can be derived from either the coding or the non-coding sequence.

In another embodiment an isolated sulfatase nucleic acid encodes the entire coding region. In another embodiment the isolated sulfatase nucleic acid encodes a sequence corresponding to the mature protein. Other fragments include nucleotide sequences encoding the amino acid fragments described herein.

Thus, sulfatase nucleic acid fragments further include sequences corresponding to the regions described herein, subregions also described, and specific functional sites. Sulfatase nucleic acid fragments also include combinations of the regions, segments, motifs, and other functional sites described above. It is understood that a sulfatase fragment includes any nucleic acid sequence that does not include the entire gene. A person of ordinary skill in the art would be aware of the many permutations that are possible. Nucleic acid fragments, according to the present invention, are not to be construed as encompassing those fragments that may have been disclosed prior to the invention.

Where the location of the regions or sites have been predicted by computer analysis, one of ordinary skill would appreciate that the amino acid residues constituting these regions can vary depending on the criteria used to define the regions.

20 Polynucleotide Uses

The nucleotide sequences of the present invention can be used as a "query sequence" to perform a search against public databases, for example, to identify other family members or related sequences. For more information about public databases, see page 26, above.

25 The nucleic acid fragments of the invention provide probes or primers in assays such as those described below. "Probes" are oligonucleotides that hybridize in a base-specific manner to a complementary strand of nucleic acid. Such probes include polypeptide nucleic acids, as described in Nielsen *et al.* (1991) *Science* 254:1497-1500. Typically, a probe comprises a region of nucleotide sequence that hybridizes under highly stringent conditions to at least about 15, typically about 20-25, and more typically about 30, 40, 50 or 75 consecutive nucleotides of the nucleic acid sequence shown in SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14, and the

complements thereof. More typically, the probe further comprises a label, e.g., radioisotope, fluorescent compound, enzyme, or enzyme co-factor.

As used herein, the term "primer" refers to a single-stranded oligonucleotide which acts as a point of initiation of template-directed DNA synthesis using well-known methods (e.g., PCR, LCR) including, but not limited to those described herein. The appropriate length of the primer depends on the particular use, but typically ranges from about 15 to 30 nucleotides. The term "primer site" refers to the area of the target DNA to which a primer hybridizes. The term "primer pair" refers to a set of primers including a 5' (upstream) primer that hybridizes with the 5' end of the nucleic acid sequence to be amplified and a 3' (downstream) primer that hybridizes with the complement of the sequence to be amplified.

Sulfatase polynucleotides are thus useful for probes, primers, and in biological assays. Where the polynucleotides are used to assess sulfatase properties or functions, such as in the assays described herein, all or less than all of the entire cDNA can be useful. Assays specifically directed to sulfatase functions, such as assessing agonist or antagonist activity, encompass the use of known fragments. Further, diagnostic methods for assessing sulfatase function can also be practiced with any fragment, including those fragments that may have been known prior to the invention. Similarly, in methods involving treatment of sulfatase dysfunction, all fragments are encompassed including those, which may have been known in the art.

Sulfatase polynucleotides are useful as a hybridization probe for cDNA and genomic DNA to isolate a full-length cDNA and genomic clones encoding the polypeptides described in SEQ ID NOS:1, 3, 5, or 7, and to isolate cDNA and genomic clones that correspond to variants producing the same polypeptides shown in SEQ ID NOS:1, 3, 5, or 7, or the other variants described herein. Variants can be isolated from the same tissue and organism from which a polypeptide shown in SEQ ID NOS:1, 3, 5, or 7 was isolated, different tissues from the same organism, or from different organisms. This method is useful for isolating genes and cDNA that are developmentally-controlled and therefore may be expressed in the same tissue or different tissues at different points in the development of an organism.

The probe can correspond to any sequence along the entire length of the gene encoding the sulfatase polypeptide. Accordingly, it could be derived from 5' noncoding regions, the coding region, and 3' noncoding regions.

The nucleic acid probe can be, for example, the full-length cDNA of SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14 or a fragment thereof, such as an oligonucleotide of at least 5, 10, 15, 20, 25, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to mRNA or DNA.

Fragments of the polynucleotides described herein are also useful to synthesize larger fragments or full-length polynucleotides described herein, ribozymes or antisense molecules. For example, a fragment can be hybridized to any portion of an mRNA and a larger or full-length cDNA can be produced.

Antisense nucleic acids of the invention can be designed using the nucleotide sequences of SEQ ID NOS:2, 4, 6, 8, 11, 12, 13, or 14 and constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-

diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest).

5 Additionally, the nucleic acid molecules of the invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, e.g., the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids (see Hyrup *et al.* (1996) *Bioorganic & Medicinal Chemistry* 4:5). As used herein, the
10 terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, e.g., DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed
15 using standard solid phase peptide synthesis protocols as described in Hyrup *et al.* (1996), *supra*; Perry-O'Keefe *et al.* (1996) *Proc. Natl. Acad. Sci. USA* 93:14670. PNAs can be further modified, e.g., to enhance their stability, specificity or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery
20 known in the art. The synthesis of PNA-DNA chimeras can be performed as described in Hyrup (1996), *supra*, Finn *et al.* (1996) *Nucleic Acids Res.* 24(17):3357-63, Mag *et al.* (1989) *Nucleic Acids Res.* 17:5973, and Peterser *et al.* (1975) *Bioorganic Med. Chem. Lett.* 5:1119.

 The nucleic acid molecules and fragments of the invention can also include
25 other appended groups such as peptides (e.g., for targeting host cell sulfatases *in vivo*), or agents facilitating transport across the cell membrane (see, e.g., Letsinger *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:6553-6556; Lemaitre *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; PCT Publication No. WO 88/0918) or the blood brain barrier (see, e.g., PCT Publication No. WO 89/10134). In addition,
30 oligonucleotides can be modified with hybridization-triggered cleavage agents (see, e.g., Krol *et al.* (1988) *Bio-Techniques* 6:958-976) or intercalating agents (see, e.g., Zon (1988) *Pharm Res.* 5:539-549).

Sulfatase polynucleotides are also useful as primers for PCR to amplify any given region of a sulfatase polynucleotide.

Sulfatase polynucleotides are also useful for constructing recombinant vectors. Such vectors include expression vectors that express a portion of, or all of, the sulfatase polypeptides. Vectors also include insertion vectors, used to integrate into another polynucleotide sequence, such as into the cellular genome, to alter *in situ* expression of sulfatase genes and gene products. For example, an endogenous sulfatase coding sequence can be replaced via homologous recombination with all or part of the coding region containing one or more specifically introduced mutations.

Sulfatase polynucleotides are also useful for expressing antigenic portions of sulfatase proteins.

Sulfatase polynucleotides are also useful as probes for determining the chromosomal positions of sulfatase polynucleotides by means of *in situ* hybridization methods, such as FISH. (For a review of this technique, see Verma *et al.* (1988) *Human Chromosomes: A Manual of Basic Techniques* (Pergamon Press, New York), and PCR mapping of somatic cell hybrids. The mapping of the sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. (Such data are found, for example, in V. McKusick, *Mendelian Inheritance in Man*, available on-line through Johns Hopkins University Welch Medical Library). The relationship between a gene and a disease mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, for example, Egeland *et al.* ((1987) *Nature* 325:783-787).

Moreover, differences in the DNA sequences between individuals affected and unaffected with a disease associated with a specified gene, can be determined. If a

mutation is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease. Comparison of affected and unaffected individuals generally involves first looking for structural alterations in the chromosomes, such as deletions or translocations, that are visible from chromosome spreads, or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of genes from several individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

Sulfatase polynucleotide probes are also useful to determine patterns of the presence of the gene encoding sulfatases and their variants with respect to tissue distribution, for example, whether gene duplication has occurred and whether the duplication occurs in all or only a subset of tissues. The genes can be naturally occurring or can have been introduced into a cell, tissue, or organism exogenously.

Sulfatase polynucleotides are also useful for designing ribozymes corresponding to all, or a part, of the mRNA produced from genes encoding the polynucleotides described herein.

Sulfatase polynucleotides are also useful for constructing host cells expressing a part, or all, of a sulfatase polynucleotide or polypeptide.

Sulfatase polynucleotides are also useful for constructing transgenic animals expressing all, or a part, of a sulfatase polynucleotide or polypeptide.

Sulfatase polynucleotides are also useful for making vectors that express part, or all, of a sulfatase polypeptide.

Sulfatase polynucleotides are also useful as hybridization probes for determining the level of sulfatase nucleic acid expression. Accordingly, the probes can be used to detect the presence of, or to determine levels of, sulfatase nucleic acid in cells, tissues, and in organisms. The nucleic acid whose level is determined can be DNA or RNA. Accordingly, probes corresponding to the polypeptides described herein can be used to assess gene copy number in a given cell, tissue, or organism. This is particularly relevant in cases in which there has been an amplification of a sulfatase gene.

Alternatively, the probe can be used in an *in situ* hybridization context to assess the position of extra copies of a sulfatase gene, as on extrachromosomal elements or as

integrated into chromosomes in which the sulfatase gene is not normally found, for example, as a homogeneously staining region.

These uses are relevant for diagnosis of disorders involving an increase or decrease in sulfatase expression relative to normal, such as a proliferative disorder, a
5 differentiative or developmental disorder, or a hematopoietic disorder. Disorders in which sulfatase expression is relevant include, but are not limited to, those disclosed herein above.

Disorders in which 22438 sulfatase expression is relevant include, but are not limited to, those involving the tissues as disclosed herein and those associated with
10 pain.

Disorders in which 23553 sulfatase expression is relevant include, but are not limited to, breast and colon carcinoma.

Disorders in which 25278 sulfatase expression is relevant include, but are not limited to, colon carcinoma.

15 Disorders in which 26212 sulfatase expression is relevant include, but are not limited to, hemangioma and uterine adenocarcinoma.

Thus, the present invention provides a method for identifying a disease or disorder associated with aberrant expression or activity of a sulfatase nucleic acid, in which a test sample is obtained from a subject and nucleic acid (e.g., mRNA, genomic
20 DNA) is detected, wherein the presence of the nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant expression or activity of the nucleic acid.

One aspect of the invention relates to diagnostic assays for determining nucleic acid expression as well as activity in the context of a biological sample (e.g.,
25 blood, serum, cells, tissue) to determine whether an individual has a disease or disorder, or is at risk of developing a disease or disorder, associated with aberrant nucleic acid expression or activity. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with expression or activity of the nucleic
30 acid molecules.

In vitro techniques for detection of mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detecting DNA includes Southern hybridizations and *in situ* hybridization.

Probes can be used as a part of a diagnostic test kit for identifying cells or tissues
5 that express a sulfatase, such as by measuring the level of a sulfatase-encoding nucleic acid in a sample of cells from a subject e.g., mRNA or genomic DNA, or determining if the sulfatase gene has been mutated.

Nucleic acid expression assays are useful for drug screening to identify compounds that modulate sulfatase nucleic acid expression (e.g., antisense,
10 polypeptides, peptidomimetics, small molecules or other drugs). A cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of the mRNA in the presence of the candidate compound is compared to the level of expression of the mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of nucleic acid expression
15 based on this comparison and be used, for example to treat a disorder characterized by aberrant nucleic acid expression. The modulator can bind to the nucleic acid or indirectly modulate expression, such as by interacting with other cellular components that affect nucleic acid expression.

Modulatory methods can be performed *in vitro* (e.g., by culturing the cell with
20 the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject) in patients or in transgenic animals. The invention thus provides a method for identifying a compound that can be used to treat a disorder associated with nucleic acid expression of a sulfatase gene. The method typically includes assaying the ability of the compound to modulate the expression of the sulfatase nucleic acid and thus identifying a compound
25 that can be used to treat a disorder characterized by undesired sulfatase nucleic acid expression.

The assays can be performed in cell-based and cell-free systems. Cell-based assays include cells naturally expressing the sulfatase nucleic acid or recombinant cells genetically engineered to express specific nucleic acid sequences. Alternatively,
30 candidate compounds can be assayed *in vivo* in patients or in transgenic animals.

The assay for sulfatase nucleic acid expression can involve direct assay of nucleic acid levels, such as mRNA levels, or on collateral compounds (such as substrate

hydrolysis). Further, the expression of genes that are up- or down-regulated in response to sulfatase activity can also be assayed. In this embodiment the regulatory regions of these genes can be operably linked to a reporter gene such as luciferase.

Thus, modulators of sulfatase gene expression can be identified in a method
5 wherein a cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of sulfatase mRNA in the presence of the candidate compound is compared to the level of expression of sulfatase mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of nucleic acid expression based on this comparison and be used, for example to treat a
10 disorder characterized by aberrant nucleic acid expression. When expression of mRNA is statistically significantly greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of nucleic acid expression. When nucleic acid expression is statistically significantly less in the presence of the candidate compound than in its absence, the candidate compound is identified as an
15 inhibitor of nucleic acid expression.

Accordingly, the invention provides methods of treatment, with the nucleic acid as a target, using a compound identified through drug screening as a gene modulator to modulate sulfatase nucleic acid expression. Modulation includes both up-regulation (i.e. activation or agonization) or down-regulation (suppression or antagonization) or effects
20 on nucleic acid activity (e.g. when nucleic acid is mutated or improperly modified). Treatment is of disorders characterized by aberrant expression or activity of the nucleic acid.

Alternatively, a modulator for sulfatase nucleic acid expression can be a small molecule or drug identified using the screening assays described herein as long as the
25 drug or small molecule inhibits sulfatase nucleic acid expression.

Sulfatase polynucleotides are also useful for monitoring the effectiveness of modulating compounds on the expression or activity of a sulfatase gene in clinical trials or in a treatment regimen. Thus, the gene expression pattern can serve as a barometer for the continuing effectiveness of treatment with the compound, particularly with
30 compounds to which a patient can develop resistance. The gene expression pattern can also serve as a marker indicative of a physiological response of the affected cells to the compound. Accordingly, such monitoring would allow either increased administration

of the compound or the administration of alternative compounds to which the patient has not become resistant. Similarly, if the level of nucleic acid expression falls below a desirable level, administration of the compound could be commensurately decreased.

Monitoring can be, for example, as follows: (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of a specified mRNA or genomic DNA of the invention in the pre-administration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the mRNA or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the mRNA or genomic DNA in the pre-administration sample with the mRNA or genomic DNA in the post-administration sample or samples; and (vi) increasing or decreasing the administration of the agent to the subject accordingly.

Sulfatase polynucleotides are also useful in diagnostic assays for qualitative changes in sulfatase nucleic acid, and particularly in qualitative changes that lead to pathology. The polynucleotides can be used to detect mutations in sulfatase genes and gene expression products such as mRNA. The polynucleotides can be used as hybridization probes to detect naturally-occurring genetic mutations in a sulfatase gene and thereby to determine whether a subject with the mutation is at risk for a disorder caused by the mutation. Mutations include deletion, addition, or substitution of one or more nucleotides in the gene, chromosomal rearrangement, such as inversion or transposition, modification of genomic DNA, such as aberrant methylation patterns or changes in gene copy number, such as amplification. Detection of a mutated form of a sulfatase gene associated with a dysfunction provides a diagnostic tool for an active disease or susceptibility to disease when the disease results from overexpression, underexpression, or altered expression of a sulfatase.

Mutations in a sulfatase gene can be detected at the nucleic acid level by a variety of techniques. Genomic DNA can be analyzed directly or can be amplified by using PCR prior to analysis. RNA or cDNA can be used in the same way.

In certain embodiments, detection of the mutation involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.* (1988) *Science* 241:1077-1080; and

Nakazawa *et al.* (1994) *PNAS* 91:360-364), the latter of which can be particularly useful for detecting point mutations in the gene (see Abravaya *et al.* (1995) *Nucleic Acids Res.* 23:675-682). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a gene under conditions such that hybridization and amplification of the gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. Deletions and insertions can be detected by a change in size of the amplified product compared to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to normal RNA or antisense DNA sequences.

It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

Alternative amplification methods include: self sustained sequence replication (Guatelli *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi *et al.* (1988) *Bio/Technology* 6:1197), or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well-known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

Alternatively, mutations in a sulfatase gene can be directly identified, for example, by alterations in restriction enzyme digestion patterns determined by gel electrophoresis.

Further, sequence-specific ribozymes (U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

Perfectly matched sequences can be distinguished from mismatched sequences by nuclease cleavage digestion assays or by differences in melting temperature.

Sequence changes at specific locations can also be assessed by nuclease protection assays such as RNase and S1 protection or the chemical cleavage method.

Furthermore, sequence differences between a mutant sulfatase gene and a wild-type gene can be determined by direct DNA sequencing. A variety of automated sequencing procedures can be utilized when performing the diagnostic assays ((1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.* (1996) *Adv. Chromatogr.* 36:127-162; and Griffin *et al.* (1993) *Appl. Biochem. Biotechnol.* 38:147-159).

Other methods for detecting mutations in the gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA duplexes (Myers *et al.* (1985) *Science* 230:1242); Cotton *et al.* (1988) *PNAS* 85:4397; Saleeba *et al.* (1992) *Meth. Enzymol.* 217:286-295), electrophoretic mobility of mutant and wild type nucleic acid is compared (Orita *et al.* (1989) *PNAS* 86:2766; Cotton *et al.* (1993) *Mutat. Res.* 285:125-144; and Hayashi *et al.* (1992) *Genet. Anal. Tech. Appl.* 9:73-79), and movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (Myers *et al.* (1985) *Nature* 313:495). The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In one embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen *et al.* (1991) *Trends Genet.* 7:5). Examples of other techniques for detecting point mutations include, selective oligonucleotide hybridization, selective amplification, and selective primer extension.

In other embodiments, genetic mutations can be identified by hybridizing a sample and control nucleic acids, e.g., DNA or RNA, to high density arrays containing hundreds or thousands of oligonucleotide probes (Cronin *et al.* (1996) *Human Mutation* 7:244-255; Kozal *et al.* (1996) *Nature Medicine* 2:753-759). For example, genetic mutations can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin *et al. supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the

characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

5 Sulfatase polynucleotides are also useful for testing an individual for a genotype that while not necessarily causing the disease, nevertheless affects the treatment modality. Thus, the polynucleotides can be used to study the relationship between an individual's genotype and the individual's response to a compound used for treatment (pharmacogenomic relationship). In the present case, for example, a mutation in the
10 sulfatase gene that results in altered affinity for a substrate-related compound could result in an excessive or decreased drug effect with standard concentrations of the compound. Accordingly, the sulfatase polynucleotides described herein can be used to assess the mutation content of the gene in an individual in order to select an appropriate compound or dosage regimen for treatment.

15 Thus polynucleotides displaying genetic variations that affect treatment provide a diagnostic target that can be used to tailor treatment in an individual. Accordingly, the production of recombinant cells and animals containing these polymorphisms allow effective clinical design of treatment compounds and dosage regimens.

 The methods can involve obtaining a control biological sample from a control
20 subject, contacting the control sample with a compound or agent capable of detecting mRNA, or genomic DNA, such that the presence of mRNA or genomic DNA is detected in the biological sample, and comparing the presence of mRNA or genomic DNA in the control sample with the presence of mRNA or genomic DNA in the test sample.

25 Sulfatase polynucleotides are also useful for chromosome identification when the sequence is identified with an individual chromosome and to a particular location on the chromosome. First, the DNA sequence is matched to the chromosome by *in situ* or other chromosome-specific hybridization. Sequences can also be correlated to specific chromosomes by preparing PCR primers that can be used for PCR screening of somatic
30 cell hybrids containing individual chromosomes from the desired species. Only hybrids containing the chromosome containing the gene homologous to the primer will yield an amplified fragment. Sublocalization can be achieved using chromosomal fragments.

Other strategies include prescreening with labeled flow-sorted chromosomes and preselection by hybridization to chromosome-specific libraries. Further mapping strategies include fluorescence *in situ* hybridization, which allows hybridization with probes shorter than those traditionally used. Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on the chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

Sulfatase polynucleotides can also be used to identify individuals from small biological samples. This can be done for example using restriction fragment-length polymorphism (RFLP) to identify an individual. Thus, the polynucleotides described herein are useful as DNA markers for RFLP (See U.S. Patent No. 5,272,057).

Furthermore, the sulfatase sequences can be used to provide an alternative technique, which determines the actual DNA sequence of selected fragments in the genome of an individual. Thus, the sulfatase sequences described herein can be used to prepare two PCR primers from the 5' and 3' ends of the sequences. These primers can then be used to amplify DNA from an individual for subsequent sequencing.

Panels of corresponding DNA sequences from individuals prepared in this manner can provide unique individual identifications, as each individual will have a unique set of such DNA sequences. It is estimated that allelic variation in humans occurs with a frequency of about once per each 500 bases. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. Sulfatase sequences can be used to obtain such identification sequences from individuals and from tissue. The sequences represent unique fragments of the human genome. Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes.

If a panel of reagents from the sequences is used to generate a unique identification database for an individual, those same reagents can later be used to identify tissue from that individual. Using the unique identification database, positive

identification of the individual, living or dead, can be made from extremely small tissue samples.

Sulfatase polynucleotides can also be used in forensic identification procedures. PCR technology can be used to amplify DNA sequences taken from very small
5 biological samples, such as a single hair follicle, body fluids (e.g. blood, saliva, or semen). The amplified sequence can then be compared to a standard allowing identification of the origin of the sample.

Sulfatase polynucleotides can thus be used to provide polynucleotide reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can enhance the
10 reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (i.e. another DNA sequence that is unique to a particular individual). As described above, actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments. Sequences targeted to the noncoding region are particularly useful
15 since greater polymorphism occurs in the noncoding regions, making it easier to differentiate individuals using this technique.

Sulfatase polynucleotides can further be used to provide polynucleotide reagents, e.g., labeled or labelable probes which can be used in, for example, an *in situ* hybridization technique, to identify a specific tissue. This is useful in cases in which a
20 forensic pathologist is presented with a tissue of unknown origin. Panels of sulfatase probes can be used to identify tissue by species and/or by organ type.

In a similar fashion, these primers and probes can be used to screen tissue culture for contamination (i.e. screen for the presence of a mixture of different types of cells in a culture).

25 Alternatively, sulfatase polynucleotides can be used directly to block transcription or translation of sulfatase gene sequences by means of antisense or ribozyme constructs. Thus, in a disorder characterized by abnormally high or undesirable sulfatase gene expression, nucleic acids can be directly used for treatment.

Sulfatase polynucleotides are thus useful as antisense constructs to control
30 sulfatase gene expression in cells, tissues, and organisms. A DNA antisense polynucleotide is designed to be complementary to a region of the gene involved in transcription, preventing transcription and hence production of sulfatase protein. An

antisense RNA or DNA polynucleotide would hybridize to the mRNA and thus block translation of mRNA into sulfatase protein.

Examples of antisense molecules useful to inhibit nucleic acid expression include antisense molecules complementary to a fragment of the 5' untranslated region of SEQ
5 ID NOS:2, 4, 6, or 8, which also includes the start codon and antisense molecules which are complementary to a fragment of the 3' untranslated region of SEQ ID NOS:2, 4, 6, or 8.

Alternatively, a class of antisense molecules can be used to inactivate mRNA in order to decrease expression of sulfatase nucleic acid. Accordingly, these molecules can
10 treat a disorder characterized by abnormal or undesired sulfatase nucleic acid expression. This technique involves cleavage by means of ribozymes containing nucleotide sequences complementary to one or more regions in the mRNA that attenuate the ability of the mRNA to be translated. Possible regions include coding regions and particularly coding regions corresponding to the catalytic and other functional activities of the
15 sulfatase protein.

Sulfatase polynucleotides also provide vectors for gene therapy in patients containing cells that are aberrant in sulfatase gene expression. Thus, recombinant cells, which include the patient's cells that have been engineered *ex vivo* and returned to the patient, are introduced into an individual where the cells produce the desired sulfatase
20 protein to treat the individual.

The invention also encompasses kits for detecting the presence of a sulfatase nucleic acid in a biological sample. For example, the kit can comprise reagents such as a labeled or labelable nucleic acid or agent capable of detecting sulfatase nucleic acid in a biological sample; means for determining the amount of sulfatase nucleic acid in the
25 sample; and means for comparing the amount of sulfatase nucleic acid in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect sulfatase mRNA or DNA.

Computer Readable Means

30 The nucleotide or amino acid sequences of the invention are also provided in a variety of mediums to facilitate use thereof. As used herein, "provided" refers to a manufacture, other than an isolated nucleic acid or amino acid molecule, which

contains a nucleotide or amino acid sequence of the present invention. Such a manufacture provides the nucleotide or amino acid sequences, or a subset thereof (e.g., a subset of open reading frames (ORFs)) in a form which allows a skilled artisan to examine the manufacture using means not directly applicable to examining the
5 nucleotide or amino acid sequences, or a subset thereof, as they exists in nature or in purified form.

In one application of this embodiment, a nucleotide or amino acid sequence of the present invention can be recorded on computer readable media. As used herein, "computer readable media" refers to any medium that can be read and accessed
10 directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. The skilled artisan will readily appreciate how any of the presently known computer readable
15 mediums can be used to create a manufacture comprising computer readable medium having recorded thereon a nucleotide or amino acid sequence of the present invention.

As used herein, "recorded" refers to a process for storing information on computer readable medium. The skilled artisan can readily adopt any of the presently known methods for recording information on computer readable medium to generate
20 manufactures comprising the nucleotide or amino acid sequence information of the present invention.

A variety of data storage structures are available to a skilled artisan for creating a computer readable medium having recorded thereon a nucleotide or amino acid sequence of the present invention. The choice of the data storage structure will
25 generally be based on the means chosen to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable medium. The sequence information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, or
30 represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. The skilled artisan can readily adapt any number of dataprocessor structuring formats (e.g., text file or database) in order to obtain

computer readable medium having recorded thereon the nucleotide sequence information of the present invention.

By providing the nucleotide or amino acid sequences of the invention in computer readable form, the skilled artisan can routinely access the sequence information for a variety of purposes. For example, one skilled in the art can use the nucleotide or amino acid sequences of the invention in computer readable form to compare a target sequence or target structural motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of the sequences of the invention which match a particular target sequence or target motif.

As used herein, a "target sequence" can be any DNA or amino acid sequence of six or more nucleotides or two or more amino acids. A skilled artisan can readily recognize that the longer a target sequence is, the less likely a target sequence will be present as a random occurrence in the database. The most preferred sequence length of a target sequence is from about 10 to 100 amino acids or from about 30 to 300 nucleotide residues. However, it is well recognized that commercially important fragments, such as sequence fragments involved in gene expression and protein processing, may be of shorter length.

As used herein, "a target structural motif," or "target motif," refers to any rationally selected sequence or combination of sequences in which the sequence(s) are chosen based on a three-dimensional configuration which is formed upon the folding of the target motif. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzyme active sites and signal sequences. Nucleic acid target motifs include, but are not limited to, promoter sequences, hairpin structures and inducible expression elements (protein binding sequences).

Computer software is publicly available which allows a skilled artisan to access sequence information provided in a computer readable medium for analysis and comparison to other sequences. A variety of known algorithms are disclosed publicly and a variety of commercially available software for conducting search means are and can be used in the computer-based systems of the present invention. Examples of such software includes, but is not limited to, MacPattern (EMBL), BLASTN and BLASTX (NCBIA).

For example, software which implements the BLAST (Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-410) and BLAZE (Brutlag *et al.* (1993) *Comp. Chem.* 17:203-207) search algorithms on a Sybase system can be used to identify open reading frames (ORFs) of the sequences of the invention which contain homology to ORFs or proteins from other libraries. Such ORFs are protein encoding fragments and are useful in producing commercially important proteins such as enzymes used in various reactions and in the production of commercially useful metabolites.

Vectors/Host Cells

The invention also provides vectors containing sulfatase polynucleotides. The term "vector" refers to a vehicle, preferably a nucleic acid molecule that can transport sulfatase polynucleotides. When the vector is a nucleic acid molecule, the sulfatase polynucleotides are covalently linked to the vector nucleic acid. With this aspect of the invention, the vector includes a plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC, PAC, YAC, OR MAC.

A vector can be maintained in the host cell as an extrachromosomal element where it replicates and produces additional copies of sulfatase polynucleotides. Alternatively, the vector may integrate into the host cell genome and produce additional copies of sulfatase polynucleotides when the host cell replicates.

The invention provides vectors for the maintenance (cloning vectors) or vectors for expression (expression vectors) of sulfatase polynucleotides. The vectors can function in prokaryotic or eukaryotic cells or in both (shuttle vectors).

Expression vectors contain cis-acting regulatory regions that are operably linked in the vector to sulfatase polynucleotides such that transcription of the polynucleotides is allowed in a host cell. The polynucleotides can be introduced into the host cell with a separate polynucleotide capable of affecting transcription. Thus, the second polynucleotide may provide a trans-acting factor interacting with the cis-regulatory control region to allow transcription of sulfatase polynucleotides from the vector. Alternatively, a trans-acting factor may be supplied by the host cell. Finally, a trans-acting factor can be produced from the vector itself.

It is understood, however, that in some embodiments, transcription and/or translation of sulfatase polynucleotides can occur in a cell-free system.

The regulatory sequence to which the polynucleotides described herein can be operably linked include promoters for directing mRNA transcription. These include, but
5 are not limited to, the left promoter from bacteriophage λ , the lac, TRP, and TAC promoters from *E. coli*, the early and late promoters from SV40, the CMV immediate early promoter, the adenovirus early and late promoters, and retrovirus long-terminal repeats.

In addition to control regions that promote transcription, expression vectors may
10 also include regions that modulate transcription, such as repressor binding sites and enhancers. Examples include the SV40 enhancer, the cytomegalovirus immediate early enhancer, polyoma enhancer, adenovirus enhancers, and retrovirus LTR enhancers.

In addition to containing sites for transcription initiation and control, expression
vectors can also contain sequences necessary for transcription termination and, in the
15 transcribed region a ribosome binding site for translation. Other regulatory control elements for expression include initiation and termination codons as well as polyadenylation signals. The person of ordinary skill in the art would be aware of the numerous regulatory sequences that are useful in expression vectors. Such regulatory sequences are described, for example, in Sambrook *et al.* (1989) *Molecular Cloning: A*
20 *Laboratory Manual 2nd. ed.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

A variety of expression vectors can be used to express a sulfatase polynucleotide. Such vectors include chromosomal, episomal, and virus-derived vectors, for example vectors derived from bacterial plasmids, from bacteriophage, from yeast episomes, from
25 yeast chromosomal elements, including yeast artificial chromosomes, from viruses such as baculoviruses, papovaviruses such as SV40, Vaccinia viruses, adenoviruses, poxviruses, pseudorabies viruses, and retroviruses. Vectors may also be derived from combinations of these sources such as those derived from plasmid and bacteriophage genetic elements, e.g. cosmids and phagemids. Appropriate cloning and expression
30 vectors for prokaryotic and eukaryotic hosts are described in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual 2nd. ed.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

The regulatory sequence may provide constitutive expression in one or more host cells (i.e. tissue specific) or may provide for inducible expression in one or more cell types such as by temperature, nutrient additive, or exogenous factor such as a hormone or other ligand. A variety of vectors providing for constitutive and inducible expression
5 in prokaryotic and eukaryotic hosts are well known to those of ordinary skill in the art.

Sulfatase polynucleotides can be inserted into the vector nucleic acid by well-known methodology. Generally, the DNA sequence that will ultimately be expressed is joined to an expression vector by cleaving the DNA sequence and the expression vector with one or more restriction enzymes and then ligating the fragments together.
10 Procedures for restriction enzyme digestion and ligation are well known to those of ordinary skill in the art.

The vector containing the appropriate polynucleotide can be introduced into an appropriate host cell for propagation or expression using well-known techniques. Bacterial cells include, but are not limited to, *E. coli*, *Streptomyces*, and *Salmonella*
15 *typhimurium*. Eukaryotic cells include, but are not limited to, yeast, insect cells such as *Drosophila*, animal cells such as COS and CHO cells, and plant cells.

As described herein, it may be desirable to express the polypeptide as a fusion protein. Accordingly, the invention provides fusion vectors that allow for the production of sulfatase polypeptides. Fusion vectors can increase the expression of a recombinant
20 protein, increase the solubility of the recombinant protein, and aid in the purification of the protein by acting for example as a ligand for affinity purification. A proteolytic cleavage site may be introduced at the junction of the fusion moiety so that the desired polypeptide can ultimately be separated from the fusion moiety. Proteolytic enzymes include, but are not limited to, factor Xa, thrombin, and enterokinase. Typical fusion
25 expression vectors include pGEX (Smith *et al.* (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.* (1988) *Gene* 69:301-315) and pET 11d
30 (Studier *et al.* (1990) *Gene Expression Technology: Methods in Enzymology* 185:60-89).

Recombinant protein expression can be maximized in a host bacteria by providing a genetic background wherein the host cell has an impaired capacity to

proteolytically cleave the recombinant protein. (Gottesman, S. (1990) *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California 119-128).

5 It is further recognized that the nucleic acid sequences of the invention can be altered to contain codons, which are preferred, or non preferred, for a particular expression system. For example, the nucleic acid can be one in which at least one altered codon, and preferably at least 10%, or 20% of the codons have been altered such that the sequence is optimized for expression in *E. coli*, yeast, human, insect, or CHO cells. Methods for determining such codon usage are well known in the art.

10 Sulfatase polynucleotides can also be expressed by expression vectors that are operative in yeast. Examples of vectors for expression in yeast e.g., *S. cerevisiae* include pYepSec1 (Baldari *et al.* (1987) *EMBO J.* 6:229-234), pMFa (Kurjan *et al.* (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.* (1987) *Gene* 54:113-123), and pYES2 (Invitrogen Corporation, San Diego, CA).

15 Sulfatase polynucleotides can also be expressed in insect cells using, for example, baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.* (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow *et al.* (1989) *Virology* 170:31-39).

20 In certain embodiments of the invention, the polynucleotides described herein are expressed in mammalian cells using mammalian expression vectors. Examples of mammalian expression vectors include pCDM8 (Seed, B. (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.* (1987) *EMBO J.* 6:187-195).

25 The expression vectors listed herein are provided by way of example only of the well-known vectors available to those of ordinary skill in the art that would be useful to express sulfatase polynucleotides. The person of ordinary skill in the art would be aware of other vectors suitable for maintenance propagation or expression of the polynucleotides described herein. These are found for example in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual 2nd, ed.*, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

30 The invention also encompasses vectors in which the nucleic acid sequences described herein are cloned into the vector in reverse orientation, but operably linked to a

regulatory sequence that permits transcription of antisense RNA. Thus, an antisense transcript can be produced to all, or to a portion, of the polynucleotide sequences described herein, including both coding and non-coding regions. Expression of this antisense RNA is subject to each of the parameters described above in relation to
5 expression of the sense RNA (regulatory sequences, constitutive or inducible expression, tissue-specific expression).

The invention also relates to recombinant host cells containing the vectors described herein. Host cells therefore include prokaryotic cells, lower eukaryotic cells such as yeast, other eukaryotic cells such as insect cells, and higher eukaryotic cells such
10 as mammalian cells.

The recombinant host cells are prepared by introducing the vector constructs described herein into the cells by techniques readily available to the person of ordinary skill in the art. These include, but are not limited to, calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic lipid-mediated transfection,
15 electroporation, transduction, infection, lipofection, and other techniques such as those found in Sambrook *et al.* (*Molecular Cloning: A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

Host cells can contain more than one vector. Thus, different nucleotide
20 sequences can be introduced on different vectors of the same cell. Similarly, sulfatase polynucleotides can be introduced either alone or with other polynucleotides that are not related to sulfatase polynucleotides such as those providing trans-acting factors for expression vectors. When more than one vector is introduced into a cell, the vectors can be introduced independently, co-introduced or joined to the sulfatase polynucleotide
25 vector.

In the case of bacteriophage and viral vectors, these can be introduced into cells as packaged or encapsulated virus by standard procedures for infection and transduction. Viral vectors can be replication-competent or replication-defective. In the case in which viral replication is defective, replication will occur in host cells providing functions that
30 complement the defects.

Vectors generally include selectable markers that enable the selection of the subpopulation of cells that contain the recombinant vector constructs. The marker can

be contained in the same vector that contains the polynucleotides described herein or may be on a separate vector. Markers include tetracycline or ampicillin-resistance genes for prokaryotic host cells and dihydrofolate reductase or neomycin resistance for eukaryotic host cells. However, any marker that provides selection for a phenotypic trait will be effective.

While the mature proteins can be produced in bacteria, yeast, mammalian cells, and other cells under the control of the appropriate regulatory sequences, cell-free transcription and translation systems can also be used to produce these proteins using RNA derived from the DNA constructs described herein.

Where secretion of the polypeptide is desired, appropriate secretion signals are incorporated into the vector. The signal sequence can be endogenous to the sulfatase polypeptides or heterologous to these polypeptides.

Where the polypeptide is not secreted into the medium, the protein can be isolated from the host cell by standard disruption procedures, including freeze thaw, sonication, mechanical disruption, use of lysing agents and the like. The polypeptide can then be recovered and purified by well-known purification methods including ammonium sulfate precipitation, acid extraction, anion or cationic exchange chromatography, phosphocellulose chromatography, hydrophobic-interaction chromatography, affinity chromatography, hydroxylapatite chromatography, lectin chromatography, or high performance liquid chromatography.

It is also understood that depending upon the host cell in recombinant production of the polypeptides described herein, the polypeptides can have various glycosylation patterns, depending upon the cell, or maybe non-glycosylated as when produced in bacteria. In addition, the polypeptides may include an initial modified methionine in some cases as a result of a host-mediated process.

Uses of Vectors and Host Cells

It is understood that "host cells" and "recombinant host cells" refer not only to the particular subject cell but also to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein. A

"purified preparation of cells", as used herein, refers to, in the case of plant or animal cells, an *in vitro* preparation of cells and not an entire intact plant or animal. In the case of cultured cells or microbial cells, it consists of a preparation of at least 10% and more preferably 50% of the subject cells.

5 The host cells expressing the polypeptides described herein, and particularly recombinant host cells, have a variety of uses. First, the cells are useful for producing sulfatase proteins or polypeptides that can be further purified to produce desired amounts of sulfatase protein or fragments. Thus, host cells containing expression vectors are useful for polypeptide production.

10 Host cells are also useful for conducting cell-based assays involving sulfatase or sulfatase fragments. Thus, a recombinant host cell expressing a native sulfatase is useful to assay for compounds that stimulate or inhibit sulfatase function, gene expression at the level of transcription or translation, and interaction with other cellular components.

 Host cells are also useful for identifying sulfatase mutants in which these
15 functions are affected. If the mutants naturally occur and give rise to a pathology, host cells containing the mutations are useful to assay compounds that have a desired effect on the mutant sulfatase (for example, stimulating or inhibiting function) which may not be indicated by their effect on the native sulfatase.

 Recombinant host cells are also useful for expressing the chimeric polypeptides
20 described herein to assess compounds that activate or suppress activation by means of a heterologous domain, segment, site, and the like, as disclosed herein.

 Further, mutant sulfatases can be designed in which one or more of the various functions is engineered to be increased or decreased and used to augment or replace sulfatase proteins in an individual. Thus, host cells can provide a therapeutic benefit by
25 replacing an aberrant sulfatase or providing an aberrant sulfatase that provides a therapeutic result. In one embodiment, the cells provide sulfatases that are abnormally active.

 In another embodiment, the cells provide sulfatases that are abnormally inactive. These sulfatases can compete with endogenous sulfatases in the individual.

30 In another embodiment, cells expressing sulfatases that cannot be activated, are introduced into an individual in order to compete with endogenous sulfatases for substrate. For example, in the case in which excessive substrate or substrate analog is

part of a treatment modality, it may be necessary to effectively inactivate the substrate or substrate analog at a specific point in treatment. Providing cells that compete for the molecule, but which cannot be affected by sulfatase activation would be beneficial.

Homologously recombinant host cells can also be produced that allow the *in situ* alteration of endogenous sulfatase polynucleotide sequences in a host cell genome. The host cell includes, but is not limited to, a stable cell line, cell *in vivo*, or cloned microorganism. This technology is more fully described in WO 93/09222, WO 91/12650, WO 91/06667, U.S. 5,272,071, and U.S. 5,641,670. Briefly, specific polynucleotide sequences corresponding to the sulfatase polynucleotides or sequences proximal or distal to a sulfatase gene are allowed to integrate into a host cell genome by homologous recombination where expression of the gene can be affected. In one embodiment, regulatory sequences are introduced that either increase or decrease expression of an endogenous sequence. Accordingly, a sulfatase protein can be produced in a cell not normally producing it. Alternatively, increased expression of sulfatase protein can be effected in a cell normally producing the protein at a specific level. Further, expression can be decreased or eliminated by introducing a specific regulatory sequence. The regulatory sequence can be heterologous to the sulfatase protein sequence or can be a homologous sequence with a desired mutation that affects expression. Alternatively, the entire gene can be deleted. The regulatory sequence can be specific to the host cell or capable of functioning in more than one cell type. Still further, specific mutations can be introduced into any desired region of the gene to produce mutant sulfatase proteins. Such mutations could be introduced, for example, into the specific functional regions such as the peptide substrate-binding site.

In one embodiment, the host cell can be a fertilized oocyte or embryonic stem cell that can be used to produce a transgenic animal containing the altered sulfatase gene. Alternatively, the host cell can be a stem cell or other early tissue precursor that gives rise to a specific subset of cells and can be used to produce transgenic tissues in an animal. See also Thomas *et al.*, *Cell* 51:503 (1987) for a description of homologous recombination vectors. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced gene has homologously recombined with the endogenous sulfatase gene is selected (see e.g., Li, E. *et al.* (1992) *Cell* 69:915). The selected cells are then injected into a blastocyst of an animal (e.g., a

mouse) to form aggregation chimeras (see e.g., Bradley, A. in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E.J. Robertson, ed. (IRL, Oxford, 1987) pp. 113-152). A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harboring the

5 homologously recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously recombined DNA by germline transmission of the transgene. Methods for constructing homologous recombination vectors and homologous recombinant animals are described further in Bradley, A. (1991) *Current Opinions in Biotechnology* 2:823-829 and in PCT International

10 Publication Nos. WO 90/11354; WO 91/01140; and WO 93/04169.

The genetically engineered host cells can be used to produce non-human transgenic animals. A transgenic animal is preferably a mammal, for example a rodent, such as a rat or mouse, in which one or more of the cells of the animal include a transgene. A transgene is exogenous DNA which is integrated into the genome of a cell

15 from which a transgenic animal develops and which remains in the genome of the mature animal in one or more cell types or tissues of the transgenic animal. These animals are useful for studying the function of a sulfatase protein and identifying and evaluating modulators of sulfatase protein activity.

Other examples of transgenic animals include non-human primates, sheep, dogs,

20 cows, goats, chickens, and amphibians.

In one embodiment, a host cell is a fertilized oocyte or an embryonic stem cell into which sulfatase polynucleotide sequences have been introduced.

A transgenic animal can be produced by introducing nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing

25 the oocyte to develop in a pseudopregnant female foster animal. Any of the sulfatase nucleotide sequences can be introduced as a transgene into the genome of a non-human animal, such as a mouse.

Any of the regulatory or other sequences useful in expression vectors can form part of the transgenic sequence. This includes intronic sequences and polyadenylation

30 signals, if not already included. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the sulfatase protein to particular cells.

Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of transgenic mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene can further be bred to other transgenic animals carrying other transgenes. A transgenic animal also includes animals in which the entire animal or tissues in the animal have been produced using the homologously recombinant host cells described herein.

In another embodiment, transgenic non-human animals can be produced which contain selected systems, which allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, e.g., Lakso *et al.* (1992) *PNAS* 89:6232-6236. Another example of a recombinase system is the FLP recombinase system of *S. cerevisiae* (O'Gorman *et al.* (1991) *Science* 251:1351-1355. If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein is required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut *et al.* (1997) *Nature* 385:810-813 and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to a pseudopregnant female

foster animal. The offspring born of this female animal will be a clone of the animal from which the cell, e.g., the somatic cell, is isolated.

Transgenic animals containing recombinant cells that express the polypeptides described herein are useful to conduct the assays described herein in an *in vivo* context.

5 Accordingly, the various physiological factors that are present *in vivo* and that could affect binding or activation, may not be evident from *in vitro* cell-free or cell-based assays. Accordingly, it is useful to provide non-human transgenic animals to assay *in vivo* sulfatase function, including peptide interaction, the effect of specific mutant sulfatases on sulfatase function and peptide interaction, and the effect of chimeric
10 sulfatases. It is also possible to assess the effect of null mutations, that is mutations that substantially or completely eliminate one or more sulfatase functions.

In general, methods for producing transgenic animals include introducing a nucleic acid sequence according to the present invention, the nucleic acid sequence capable of expressing the protein in a transgenic animal, into a cell in culture or *in vivo*.
15 When introduced *in vivo*, the nucleic acid is introduced into an intact organism such that one or more cell types and, accordingly, one or more tissue types, express the nucleic acid encoding the protein. Alternatively, the nucleic acid can be introduced into virtually all cells in an organism by transfecting a cell in culture, such as an embryonic stem cell, as described herein for the production of transgenic
20 animals, and this cell can be used to produce an entire transgenic organism. As described, in a further embodiment, the host cell can be a fertilized oocyte. Such cells are then allowed to develop in a female foster animal to produce the transgenic organism.

25 Pharmaceutical Compositions

Sulfatase nucleic acid molecules, proteins, modulators of the protein, and antibodies (also referred to herein as "active compounds") can be incorporated into pharmaceutical compositions suitable for administration to a subject, e.g., a human. Such compositions typically comprise the nucleic acid molecule, protein, modulator, or
30 antibody and a pharmaceutically acceptable carrier.

The term "administer" is used in its broadest sense and includes any method of introducing the compositions of the present invention into a subject. This includes

producing polypeptides or polynucleotides *in vivo* by *in vivo* transcription or translation of polynucleotides that have been exogenously introduced into a subject. Thus, polypeptides or nucleic acids produced in the subject from the exogenous compositions are encompassed in the term "administer."

5 As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or
10 agent is incompatible with the active compound, such media can be used in the compositions of the invention. Supplementary active compounds can also be incorporated into the compositions. A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous,
15 oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants
20 such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampules, disposable syringes or multiple dose
25 vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water,
30 Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage

and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can
5 be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents,
10 for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active
15 compound (e.g., a sulfatase protein or anti-sulfatase antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In
20 the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They
25 can be enclosed in gelatin capsules or compressed into tablets. For oral administration, the agent can be contained in enteric forms to survive the stomach or further coated or mixed to be released in a particular region of the GI tract by known methods. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions
30 can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included

as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant
5 such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser, which contains a suitable
10 propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and
15 fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with
20 conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems.
25 Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected
30 cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound
5 calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

10 The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (U.S. 5,328,470) or by stereotactic injection (see e.g., Chen *et al.* (1994) *PNAS* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an
15 acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, e.g. retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

As defined herein, a therapeutically effective amount of protein or polypeptide
20 (i.e., an effective dosage) ranges from about 0.001 to 30 mg/kg body weight, preferably about 0.01 to 25 mg/kg body weight, more preferably about 0.1 to 20 mg/kg body weight, and even more preferably about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight.

The skilled artisan will appreciate that certain factors may influence the
25 dosage required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a protein, polypeptide, or antibody can include a single treatment or, preferably, can include a series of treatments. In a preferred
30 example, a subject is treated with antibody, protein, or polypeptide in the range of between about 0.1 to 20 mg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7

weeks, and even more preferably for about 4, 5, or 6 weeks. It will also be appreciated that the effective dosage of antibody, protein, or polypeptide used for treatment may increase or decrease over the course of a particular treatment. Changes in dosage may result and become apparent from the results of diagnostic assays as described herein.

The present invention encompasses agents which modulate expression or activity. An agent may, for example, be a small molecule. For example, such small molecules include, but are not limited to, peptides, peptidomimetics, amino acids, amino acid analogs, polynucleotides, polynucleotide analogs, nucleotides, nucleotide analogs, organic or inorganic compounds (i.e., including heteroorganic and organometallic compounds) having a molecular weight less than about 10,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 5,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 1,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 500 grams per mole, and salts, esters, and other pharmaceutically acceptable forms of such compounds.

It is understood that appropriate doses of small molecule agents depends upon a number of factors within the ken of the ordinarily skilled physician, veterinarian, or researcher. The dose(s) of the small molecule will vary, for example, depending upon the identity, size, and condition of the subject or sample being treated, further depending upon the route by which the composition is to be administered, if applicable, and the effect which the practitioner desires the small molecule to have upon the nucleic acid or polypeptide of the invention. Exemplary doses include milligram or microgram amounts of the small molecule per kilogram of subject or sample weight (e.g., about 1 microgram per kilogram to about 500 milligrams per kilogram, about 100 micrograms per kilogram to about 5 milligrams per kilogram, or about 1 microgram per kilogram to about 50 micrograms per kilogram. It is furthermore understood that appropriate doses of a small molecule depend upon the potency of the small molecule with respect to the expression or activity to be modulated. Such appropriate doses may be determined using the assays described herein. When one or more of these small molecules is to be administered to an animal (e.g., a human) in order to modulate expression or activity of a polypeptide or nucleic

acid of the invention, a physician, veterinarian, or researcher may, for example, prescribe a relatively low dose at first, subsequently increasing the dose until an appropriate response is obtained. In addition, it is understood that the specific dose level for any particular animal subject will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, any drug combination, and the degree of expression or activity to be modulated.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

Other Embodiments

In another aspect, the invention features, a method of analyzing a plurality of capture probes. The method can be used, e.g., to analyze gene expression. The method includes: providing a two dimensional array having a plurality of addresses, each address of the plurality being positionally distinguishable from each other address of the plurality, and each address of the plurality having a unique capture probe, e.g., a nucleic acid or peptide sequence; contacting the array with a 22438, 23553, 25278, or 26212 nucleic acid, preferably purified, polypeptide, preferably purified, or antibody, and thereby evaluating the plurality of capture probes. Binding, e.g., in the case of a nucleic acid, hybridization with a capture probe at an address of the plurality, is detected, e.g., by signal generated from a label attached to the 22438, 23553, 25278, or 26212 nucleic acid, polypeptide, or antibody.

The capture probes can be a set of nucleic acids from a selected sample, e.g., a sample of nucleic acids derived from a control or non-stimulated tissue or cell.

The method can include contacting the 22438, 23553, 25278, or 26212 nucleic acid, polypeptide, or antibody with a first array having a plurality of capture probes and a second array having a different plurality of capture probes. The results of each hybridization can be compared, e.g., to analyze differences in expression between a first and second sample. The first plurality of capture probes can be from a control sample, e.g., a wild type, normal, or non-diseased, non-stimulated, sample, e.g., a biological fluid, tissue, or cell sample. The second plurality of capture probes can be

from an experimental sample, e.g., a mutant type, at risk, disease-state or disorder-state, or stimulated, sample, e.g., a biological fluid, tissue, or cell sample.

The plurality of capture probes can be a plurality of nucleic acid probes each of which specifically hybridizes with an allele of 22438, 23553, 25278, or 26212.

5 Such methods can be used to diagnose a subject, e.g., to evaluate risk for a disease or disorder, to evaluate suitability of a selected treatment for a subject, to evaluate whether a subject has a disease or disorder. 22438, 23553, 25278, or 26212 are associated with sulfatase activity, thus it is useful for disorders associated with abnormal sulfatase activity.

10 The method can be used to detect SNPs, as described below.

In another aspect, the invention features, a method of analyzing a plurality of probes. The method is useful, e.g., for analyzing gene expression. The method includes: providing a two dimensional array having a plurality of addresses, each address of the plurality being positionally distinguishable from each other address of
15 the plurality having a unique capture probe, e.g., wherein the capture probes are from a cell or subject which express or misexpress 22438, 23553, 25278, or 26212, or from a cell or subject in which a 22438, 23553, 25278, or 26212 mediated response has been elicited, e.g., by contact of the cell with 22438, 23553, 25278, or 26212 nucleic acid or protein, or administration to the cell or subject 22438, 23553, 25278, or 26212
20 nucleic acid or protein; contacting the array with one or more inquiry probe, wherein an inquiry probe can be a nucleic acid, polypeptide, or antibody (which is preferably other than 22438, 23553, 25278, or 26212 nucleic acid, polypeptide, or antibody); providing a two dimensional array having a plurality of addresses, each address of the plurality being positionally distinguishable from each other address of the plurality,
25 and each address of the plurality having a unique capture probe, e.g., wherein the capture probes are from a cell or subject which does not express 22438, 23553, 25278, or 26212 (or does not express as highly as in the case of the 22438, 23553, 25278, or 26212 positive plurality of capture probes) or from a cell or subject which in which a 22438, 23553, 25278, or 26212 mediated response has not been elicited (or
30 has been elicited to a lesser extent than in the first sample); contacting the array with one or more inquiry probes (which is preferably other than a 22438, 23553, 25278, or 26212 nucleic acid, polypeptide, or antibody), and thereby evaluating the plurality of

capture probes. Binding, e.g., in the case of a nucleic acid, hybridization with a capture probe at an address of the plurality, is detected, e.g., by signal generated from a label attached to the nucleic acid, polypeptide, or antibody.

In another aspect, the invention features a method of analyzing 22438, 23553, 25278, or 26212, e.g., analyzing structure, function, or relatedness to other nucleic acid or amino acid sequences. The method includes: providing a 22438, 23553, 25278, or 26212 nucleic acid or amino acid sequence; comparing the 22438, 23553, 25278, or 26212 sequence with one or more preferably a plurality of sequences from a collection of sequences, e.g., a nucleic acid or protein sequence database; to thereby analyze 22438, 23553, 25278, or 26212.

Preferred databases include GenBankTM. The method can include evaluating the sequence identity between a 22438, 23553, 25278, or 26212 sequence and a database sequence. The method can be performed by accessing the database at a second site, e.g., over the internet.

In another aspect, the invention features, a set of oligonucleotides, useful, e.g., for identifying SNP's, or identifying specific alleles of 22438, 23553, 25278, or 26212. The set includes a plurality of oligonucleotides, each of which has a different nucleotide at an interrogation position, e.g., an SNP or the site of a mutation. In a preferred embodiment, the oligonucleotides of the plurality identical in sequence with one another (except for differences in length). The oligonucleotides can be provided with different labels, such that an oligonucleotides which hybridizes to one allele provides a signal that is distinguishable from an oligonucleotides which hybridizes to a second allele.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are incorporated herein by reference.

EXAMPLES

5 Example 1: Identification and Characterization of Human 22438 cDNAs

 The human 22438 sequence (Figure 1A-B; SEQ ID NO:2), which is approximately 2175 nucleotides long including untranslated regions, contains a predicted methionine-initiated coding sequence of about 1578 nucleotides (nucleotides 248-1825 of SEQ ID NO:2; SEQ ID NO:11). The coding sequence
10 encodes a 525 amino acid protein (SEQ ID NO:1).

 PFAM analysis indicates that 22438 contains a sulfatase domain. For general information regarding PFAM identifiers, PS prefix and PF prefix domain identification numbers, refer to Sonnhammer *et al.* (1997) *Protein* 28:405-420 and <http://www.psc.edu/general/software/packages/pfam/pfam.html>.

15 As used herein, the term "sulfatase domain" includes an amino acid sequence of about 80-420 amino acid residues in length and having a bit score for the alignment of the sequence to the sulfatase domain (HMM) of at least 8. Preferably, a sulfatase domain includes at least about 100-250 amino acids, more preferably about 130-200 amino acid residues, or about 160-200 amino acids and has a bit score for the
20 alignment of the sequence to the sulfatase domain (HMM) of at least 16 or greater. The sulfatase domain (HMM) has been assigned the PFAM Accession PF00884 (<http://pfam.wustl.edu/>). An alignment of the sulfatase domain (amino acids 36-462 of SEQ ID NO:1) of human 22438 with a consensus amino acid sequence derived from a hidden Markov model is depicted in Figure 19.

25 In a preferred embodiment 22438-like polypeptide or protein has a "sulfatase domain" or a region which includes at least about 100-250, more preferably about 130-200 or 160-200, amino acid residues and has at least about 60%, 70%, 80%, 90%, 95%, 99%, or 100% sequence identity with a "sulfatase domain," e.g., the sulfatase domain of human 22438-like polypeptide or protein (e.g., amino acid residues 36-462
30 of SEQ ID NO:1).

 To identify the presence of an "sulfatase" domain in a 22438-like protein sequence, and make the determination that a polypeptide or protein of interest has a

particular profile, the amino acid sequence of the protein can be searched against a database of HMMs (e.g., the Pfam database, release 2.1) using the default parameters (http://www.sanger.ac.uk/Software/Pfam/HMM_search). For example, the hmmsf program, which is available as part of the HMMER package of search programs, is a family specific default program for MILPAT0063 and a score of 15 is the default threshold score for determining a hit. Alternatively, the threshold score for determining a hit can be lowered (e.g., to 8 bits). A description of the Pfam database can be found in Sonhammer *et al.* (1997) *Proteins* 28(3):405-420 and a detailed description of HMMs can be found, for example, in Gribskov *et al.* (1990) *Meth. Enzymol.* 183:146-159; Gribskov *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:4355-4358; Krogh *et al.* (1994) *J. Mol. Biol.* 235:1501-1531; and Stultz *et al.* (1993) *Protein Sci.* 2:305-314, the contents of which are incorporated herein by reference.

Example 2: Tissue Distribution of 22348 mRNA

Northern blot hybridizations with various RNA samples are performed under standard conditions and washed under stringent conditions, i.e., 0.2 X SSC at 65°C. A DNA probe corresponding to all or a portion of the 22348 cDNA (SEQ ID NO:2) can be used. The DNA is radioactively labeled with ³²P-dCTP using the Prime-It Kit (Stratagene, La Jolla, CA) according to the instructions of the supplier. Filters containing mRNA from mouse hematopoietic and endocrine tissues, and cancer cell lines (Clontech, Palo Alto, CA) are probed in ExpressHyb hybridization solution (Clontech) and washed at high stringency according to manufacturer's recommendations.

Example 3: Identification and Characterization of Human 23553 cDNAs

The human 23553 sequence (Figure 5A-B; SEQ ID NO:4), which is approximately 4321 nucleotides long including untranslated regions, contains a predicted methionine-initiated coding sequence of about 2616 nucleotides (nucleotides 510-3125 of SEQ ID NO:4; SEQ ID NO:12). The coding sequence encodes a 871 amino acid protein (SEQ ID NO:3).

PFAM analysis indicates that 23553 has a sulfatase domain. For general information regarding PFAM identifiers, PS prefix and PF prefix domain

identification numbers, refer to Sonnhammer *et al.* (1997) *Protein* 28:405-420 and <http://www.psc.edu/general/software/packages/pfam/pfam.html>. An alignment of the sulfatase domain (amino acids 43 to 467 of SEQ ID NO:3) of human 23553-like with a consensus amino acid sequence derived from a hidden Markov model is depicted in
5 Figure 20. For further information on sulfatase domains, see Example 1.

In one embodiment, a 23553-like protein includes at least one transmembrane domain. As used herein, the term "transmembrane domain" includes an amino acid sequence of about 15 amino acid residues in length that spans a phospholipid membrane. More preferably, a transmembrane domain includes about at least 18, 20,
10 22, or 24 amino acid residues and spans a phospholipid membrane. Transmembrane domains are rich in hydrophobic residues, and typically have an α -helical structure. In a preferred embodiment, at least 50%, 60%, 70%, 80%, 90%, 95% or more of the amino acids of a transmembrane domain are hydrophobic, e.g., leucines, isoleucines, tyrosines, or tryptophans. Transmembrane domains are described in, for example,
15 <http://pfam.wustl.edu/cgi-bin/getdesc?name=7tm-1>, and Zagotta W.N. *et al.* (1996) *Annual Rev. Neurosci.* 19:235-63, the contents of which are incorporated herein by reference.

In a preferred embodiment, a 23553-like polypeptide or protein has at least one transmembrane domain or a region which includes at least 18, 20, 22, or 24 amino
20 acid residues and has at least about 60%, 70% 80% 90% 95%, 99%, or 100% sequence identity with a "transmembrane domain," e.g., at least one transmembrane domain of human 23553 (e.g., amino acid residues 7 to 25 of SEQ ID NO:3).

In another embodiment, a 23553 protein includes at least one "non-transmembrane domain." As used herein, "non-transmembrane domains" are
25 domains that reside outside of the membrane. When referring to plasma membranes, non-transmembrane domains include extracellular domains (i.e., outside of the cell) and intracellular domains (i.e., within the cell). When referring to membrane-bound proteins found in intracellular organelles (e.g., mitochondria, endoplasmic reticulum, peroxisomes and microsomes), non-transmembrane domains include those domains of
30 the protein that reside in the cytosol (i.e., the cytoplasm), the lumen of the organelle, or the matrix or the intermembrane space (the latter two relate specifically to mitochondria organelles). The C-terminal amino acid residue of a non-transmembrane

domain is adjacent to an N-terminal amino acid residue of a transmembrane domain in a naturally occurring 23553-like protein.

In a preferred embodiment, a 23553-like polypeptide or protein has a "non-transmembrane domain" or a region which includes at least about 1-350, preferably about 200-320, more preferably about 230-300, and even more preferably about 240-280 amino acid residues, and has at least about 60%, 70% 80% 90% 95%, 99% or 100% sequence identity with a "non-transmembrane domain", e.g., a non-transmembrane domain of human 23553-like protein.

A non-transmembrane domain located at the N-terminus of a 23553-like protein or polypeptide is referred to herein as an "N-terminal non-transmembrane domain." As used herein, an "N-terminal non-transmembrane domain" includes an amino acid sequence having about 1-100. For example, an N-terminal non-transmembrane domain is located at about amino acid residues 1 to 6 of SEQ ID NO:3.

Similarly, a non-transmembrane domain located at the C-terminus of a 23553-like protein or polypeptide is referred to herein as a "C-terminal non-transmembrane domain." As used herein, a "C-terminal non-transmembrane domain" includes an amino acid sequence having about 1-800, preferably about 15-500, preferably about 20-270, more preferably about 25-255 amino acid residues in length and is located outside the boundaries of a membrane. For example, a C-terminal non-transmembrane domain is located at about amino acid residues 26-871 of SEQ ID NO:3.

The ORF analyzer predicts that 23553 has a signal peptide. Therefore, a 23553-like molecule can further include a signal sequence. As used herein, a "signal sequence" refers to a peptide of about 20-80 amino acid residues in length which occurs at the N-terminus of secretory and integral membrane proteins and which contains a majority of hydrophobic amino acid residues. For example, a signal sequence contains at least about 12-25 amino acid residues, preferably about 30-70 amino acid residues, and has at least about 40-70%, preferably about 50-65%, and more preferably about 55-60% hydrophobic amino acid residues (e.g., alanine, valine, leucine, isoleucine, phenylalanine, tyrosine, tryptophan, or proline). Such a "signal sequence", also referred to in the art as a "signal peptide", serves to direct a protein

containing such a sequence to a lipid bilayer. For example, in one embodiment, a 23553-like protein contains a signal sequence of about amino acids 1-22 of SEQ ID NO:3. The "signal sequence" is cleaved during processing of the mature protein. The mature 23553-like protein corresponds to amino acids 23-871 of SEQ ID NO:3.

5 CLUSTAL multiple sequence alignment analysis shows homology between 23553 and the following sequences (identified by GenBank accession number): P14217, *Chlamydomonas reinhardtii* arylsulfatase; Q10723, *Volvox carteri* arylsulfatase; CAB40661, human N-acetylglucosamine-6-sulfatase homolog; P15586, human N-acetylglucosamine-6-sulfatase; P50426, goat N-acetylglucosamine-6-sulfatase; AAA83618, *C. elegans* putative sulfatase; AAC02716, *Neurospora crassa* arylsulfatase; P31447, *E. coli* hypothetical sulfatase.

Example 4: Tissue Distribution of 23553 mRNA

15 In normal human tissues tested, high expression of 23553 was observed in trachea, vein, osteoblast, kidney, and testes. Significant expression of 23553 was found in adipose, colon, skeletal muscle, thyroid, prostate, and other tissues. See Figure 25. In comparisons of normal and tumor tissue, 23553 expression was detected in all samples tested, with increased expression in breast, colon, and lung tumors. See Figure 26. Further, elevated expression of 23553 was found in glioblastoma samples, as compared to normal brain tissue samples. Expression levels were determined by
20 quantitative PCR (Taqman® brand quantitative PCR kit, Applied Biosystems). The quantitative PCR reactions were performed according to the kit manufacturer's instructions.

cDNA library array analysis of 23553 revealed expression in adipose, adrenal
25 gland, bone, brain, colon, colon metastases to liver, endothelial, heart, liver, lung, muscle, osteoblast, skin, testes, thyroid, and other tissue. Reverse transcriptase polymerase chain reaction (RT-PCR) revealed 23553 expression in clinical samples of normal and tumor colon tissue, normal and metastatic liver tissue, and in lung squamous cell carcinoma tissue. *In situ* hybridization showed expression of 23553 in
30 the following tissues: 3 of 3 breast tumor; 0 of 2 normal breast; 4 of 4 lung tumor; 0 of 2 normal lung; 4 of 4 colon tumor; and 2 of 2 liver metastases. In all cases,

expression of 23553 was confined to the stromal component of tissue; no expression was detected in normal or tumor epithelium.

Angiogenic growth factors (e.g., bFGF) are present in the extracellular matrix (ECM), and can be released from the ECM by heparinase-like enzymes. This includes the glycosyl-sulfatases. The released growth factors in turn stimulate blood vessel formation. See Baird A, Ling N., "Fibroblast growth factors are present in the extracellular matrix produced by endothelial cells in vitro: implications for a role of heparinase-like enzymes in the neovascular response," *Biochem Biophys Res Commun.* (1987) 142(2):428-35.

As noted, 23553 has amino acid sequence features that place it in the class of glycosyl sulfate cleaving enzymes. Taqman results (above) show that its expression is elevated in clinical tumor samples. *In situ* hybridization shows specific, localized 23553 expression in the tumor stromal component of all tumor samples tested, whereas its expression is low or absent in normal tissues. This suggests that, through catalytic activity, 23553 promotes tumor growth or is involved in tumor maintenance by degrading the ECM and releasing growth factors.

Example 5: Identification and Characterization of Human 25278 cDNAs

The human 25278 sequence (Figure 10A-B; SEQ ID NO:6), which is approximately 2940 nucleotides long including untranslated regions, contains a predicted methionine-initiated coding sequence of about 1710 nucleotides (nucleotides 334-2043 of SEQ ID NO:6; SEQ ID NO:13). The coding sequence encodes a 569 amino acid protein (SEQ ID NO:5).

PFAM analysis indicates that 25278 has a sulfatase domain. For general information regarding PFAM identifiers, PS prefix and PF prefix domain identification numbers, refer to Sonnhammer *et al.* (1997) *Protein* 28:405-420 and <http://www.psc.edu/general/software/packages/pfam/pfam.html>. An alignment of the sulfatase domain (amino acids 47 to 471 of SEQ ID NO:5) of human 25278 with a consensus amino acid sequence derived from a hidden Markov model is depicted in Figure 27. For further information on sulfatase domains, see Example 1.

Example 6: Identification and Characterization of Human 26212 cDNAs

The human 26212 sequence (Figure 15; SEQ ID NO:8), which is approximately 2253 nucleotides long including untranslated regions, contains a predicted methionine-initiated coding sequence of about 1800 nucleotides (nucleotides 324-2123 of SEQ ID NO:8; SEQ ID NO:14). The coding sequence encodes a 599 amino acid protein (SEQ ID NO:7).

PFAM analysis indicates that 26212 has a sulfatase domain. For general information regarding PFAM identifiers, PS prefix and PF prefix domain identification numbers, refer to Sonnhammer *et al.* (1997) *Protein* 28:405-420 and <http://www.psc.edu/general/software/packages/pfam/pfam.html>. An alignment of the sulfatase domain (amino acids 76-502 of SEQ ID NO:7) of human 26212 with a consensus amino acid sequence derived from a hidden Markov model is depicted in Figure 29. For further information on sulfatase domains, see Example 1.

In one embodiment, 26212-like protein includes at least one transmembrane domain. As used herein, the term "transmembrane domain" includes an amino acid sequence of about 15 amino acid residues in length that spans a phospholipid membrane. More preferably, a transmembrane domain includes about at least 18, 20, 22, or 24 amino acid residues and spans a phospholipid membrane. For more information on transmembrane domains, see example 3.

In a preferred embodiment, a 26212-like polypeptide or protein has at least one transmembrane domain or a region which includes at least 18, 20, 22, 24, 25, or 30 amino acid residues and has at least about 60%, 70% 80% 90% 95%, 99%, or 100% sequence identity with a "transmembrane domain," e.g., at least one transmembrane domain of human 26212-like polypeptide or protein (e.g., amino acid residues 24 to 44 of SEQ ID NO:7).

In another embodiment, a 26212-like protein includes at least one "non-transmembrane domain." The C-terminal amino acid residue of a non-transmembrane domain is adjacent to an N-terminal amino acid residue of a transmembrane domain in a naturally occurring 26212-like protein. For more information on non-transmembrane domains, see Example 3.

In a preferred embodiment, a 26212-like polypeptide or protein has a “non-transmembrane domain” or a region which includes at least about 1-350, preferably about 200-320, more preferably about 230-300, and even more preferably about 240-280 amino acid residues, and has at least about 60%, 70% 80% 90% 95%, 99% or
5 100% sequence identity with a “non-transmembrane domain”, e.g., a non-transmembrane domain of human 26212-like polypeptide or protein. An N-terminal non-transmembrane domain is located at about amino acid residues 1 to 23 of SEQ ID NO:7. A C-terminal non-transmembrane domain is located at about amino acid residues 45 to 599 of SEQ ID NO:7.

10 A 26212-like molecule can further include a signal sequence. For more information on signal sequences, see Example 3.

Example 7: Tissue Distribution of 26212 mRNA

In six independent experiments, 26212 showed higher levels of expression in
15 proliferating endothelial cells as compared to arrested endothelial cells. 26212 expression was also higher in proliferating endothelial cells than in non-endothelial cells. See Figure 30. 26212 expression levels were upregulated in breast tissue cell lines treated with epidermal growth factor, as well. See Figure 34. 26212 is expressed in hemangiomas and other angiogenic tissues, including fetal heart, uterine
20 adenocarcinoma, and endometrial polyps. See Figure 35. Endothelial and glial cells showed higher levels of 26212 expression as compared to other tissues and cells. See Figure 36. 26212 also showed higher levels of expressing in some lung, breast and brain tumors as compared to normal tissues. Expression levels of 26212 were found to be higher in proliferating endothelial cells than in tumors, too. Expression levels
25 were determined by quantitative PCR (Taqman® brand quantitative PCR kit, Applied Biosystems). The quantitative PCR reactions were performed according to the kit manufacturer's instructions.

In situ hybridization analysis was also carried out. 26212 showed weak expression in ovarian tumor, and no expression in normal ovary. Similarly, colon
30 metastases showed weak expression of 26212, and normal colon tissue and primary tumors showed no expression. A subset of lung tumors tested showed expression of 26212, while no expression was revealed in normal lung.

Angiogenic growth factors (e.g., bFGF) are present in the extracellular matrix (ECM), and can be released from the ECM by heparinase-like enzymes. This includes the glycosyl-sulfatases. The released growth factors in turn stimulate blood vessel formation by, e.g., attracting endothelial cells to form new vessels. See Baird
5 A, Ling N., "Fibroblast growth factors are present in the extracellular matrix produced by endothelial cells in vitro: implications for a role of heparinase-like enzymes in the neovascular response," *Biochem Biophys Res Commun.* (1987) 142(2):428-35.

As noted, 26212 has amino acid sequence features that place it in the class of glycosyl sulfate cleaving enzymes. Taqman results (above) show that its expression
10 is elevated in proliferating endothelial cells, suggesting that 26212 is specifically involved in active angiogenic sites.

Example 8: Recombinant Expression of 22348, 23553, 25278, or 26212 in Bacterial Cells

15 In this example, 22348, 23553, 25278, or 26212 is expressed as a recombinant glutathione-S-transferase (GST) fusion polypeptide in *E. coli* and the fusion polypeptide is isolated and characterized. Specifically, 22348, 23553, 25278, or 26212 is fused to GST and this fusion polypeptide is expressed in *E. coli*, e.g., strain PEB199. Expression of the GST-26212 fusion protein in PEB199 is induced with
20 IPTG. The recombinant fusion polypeptide is purified from crude bacterial lysates of the induced PEB199 strain by affinity chromatography on glutathione beads. Using polyacrylamide gel electrophoretic analysis of the polypeptide purified from the bacterial lysates, the molecular weight of the resultant fusion polypeptide is determined.

25

Example 9: Expression of Recombinant 22348, 23553, 25278, or 26212 Protein in COS Cells

To express the 22348, 23553, 25278, or 26212 gene in COS cells, the pcDNA/Amp vector by Invitrogen Corporation (San Diego, CA) is used. This vector
30 contains an SV40 origin of replication, an ampicillin resistance gene, an *E. coli* replication origin, a CMV promoter followed by a polylinker region, and an SV40 intron and polyadenylation site. A DNA fragment encoding the entire 22348, 23553,

25278, or 26212 protein and an HA tag (Wilson *et al.* (1984) *Cell* 37:767) or a FLAG tag fused in-frame to its 3' end of the fragment is cloned into the polylinker region of the vector, thereby placing the expression of the recombinant protein under the control of the CMV promoter.

5 To construct the plasmid, the 22348, 23553, 25278, or 26212 DNA sequence is amplified by PCR using two primers. The 5' primer contains the restriction site of interest followed by approximately twenty nucleotides of the 22348, 23553, 25278, or 26212 coding sequence starting from the initiation codon; the 3' end sequence contains complementary sequences to the other restriction site of interest, a translation
10 stop codon, the HA tag or FLAG tag and the last 20 nucleotides of the 22348, 23553, 25278, or 26212 coding sequence. The PCR amplified fragment and the pCDNA/Amp vector are digested with the appropriate restriction enzymes and the vector is dephosphorylated using the CIAP enzyme (New England Biolabs, Beverly, MA). Preferably the two restriction sites chosen are different so that the 22348,
15 23553, 25278, or 26212 gene is inserted in the correct orientation. The ligation mixture is transformed into *E. coli* cells (strains HB101, DH5 α , SURE, available from Stratagene Cloning Systems, La Jolla, CA, can be used), the transformed culture is plated on ampicillin media plates, and resistant colonies are selected. Plasmid DNA is isolated from transformants and examined by restriction analysis for the
20 presence of the correct fragment.

COS cells are subsequently transfected with the 22348, 23553, 25278, or 26212-pcDNA/Amp plasmid DNA using the calcium phosphate or calcium chloride co-precipitation methods, DEAE-dextran-mediated transfection, lipofection, or electroporation. Other suitable methods for transfecting host cells can be found in
25 Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989. The expression of the 22348, 23553, 25278, or 26212 polypeptide is detected by radiolabelling (^{35}S -methionine or ^{35}S -cysteine available from NEN, Boston, MA, can be used) and immunoprecipitation (Harlow, E.
30 and Lane, D. *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1988) using an HA specific monoclonal antibody. Briefly, the cells are labeled for 8 hours with ^{35}S -methionine (or ^{35}S -cysteine). The

culture media are then collected and the cells are lysed using detergents (RIPA buffer, 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% DOC, 50 mM Tris, pH 7.5). Both the cell lysate and the culture media are precipitated with an HA specific monoclonal antibody. Precipitated polypeptides are then analyzed by SDS-PAGE.

- 5 Alternatively, DNA containing the 22348, 23553, 25278, or 26212 coding sequence is cloned directly into the polylinker of the pCDNA/Amp vector using the appropriate restriction sites. The resulting plasmid is transfected into COS cells in the manner described above, and the expression of the 22348, 23553, 25278, or 26212 polypeptide is detected by radiolabelling and immunoprecipitation using a 22348, 10 23553, 25278, or 26212 specific monoclonal antibody.

- This invention may be embodied in many different forms and should not be construed as limited to the embodiments set forth herein; rather, these embodiments are provided so that this disclosure will fully convey the invention to those skilled in the art. Many modifications and other embodiments of the invention will come to mind in one 15 skilled in the art to which this invention pertains having the benefit of the teachings presented in the foregoing description. Although specific terms are employed, they are used as in the art unless otherwise indicated.

Applicant's or agent's file reference	35800/208709	International application No.	PCT/US01/
--	--------------	-------------------------------	-----------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 5, line 31	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit	Accession Number PTA-
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Page 17, line 12; page 22, line 9; page 23, line 23; page 108, lines 7, 13, 17, 21, 24 and 29; page 109, lines 8 and 13; page 110, lines 2, 6, 13 and 22; page 111, lines 1, 6, 9 and 13.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indicators are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") Accession Number of Deposit and Date of Deposit	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer MELVIN S. BROOKS SR INTERNATIONAL DIVISION 703-305-5163 MVB

For International Bureau use only
<input type="checkbox"/> This sheet was received with the International Bureau on:
Authorized officer

Applicant's or agent's file reference	35800/208709	International application No.	PCT/US01/
--	--------------	-------------------------------	-----------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 5, line 31	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit 05 April 2000 (05.04.00)	Accession Number PTA- 1639
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Page 17, line 12; page 22, line 10; page 23, line 23; page 108, lines 7, 13, 17, 21, 24 and 29; page 109, lines 8 and 13; page 110, lines 2, 6, 13 and 22; page 111, lines 1, 6, 9 and 13.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indicators are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	
<input checked="" type="checkbox"/> This sheet was received with the international application	
Authorized officer	MELVIN S. BROOKS SR INTERNATIONAL DIVISION 703-805-9169 m53


For International Bureau use only	
<input type="checkbox"/> This sheet was received with the International Bureau on:	
Authorized officer	

Applicant's or agent's file reference	35800/208709	International application No.	PCT/US01/
--	--------------	-------------------------------	-----------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 5, line 31	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit 09 May 2000 (09.05.00)	Accession Number PTA- 1846
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Page 17, line 12; page 22, line 10; page 23, line 23; page 108, lines 7, 13, 17, 21, 24 and 29; page 109, lines 8 and 13; page 110, lines 2, 6, 13 and 22; page 111, lines 1, 6, 9 and 13.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indicators are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	
<input checked="" type="checkbox"/>	This sheet was received with the international application
Authorized officer	

For International Bureau use only	
<input type="checkbox"/>	This sheet was received with the International Bureau on:
Authorized officer	

Applicant's or agent's file reference	35800/208709	International application No.	PCT/US01/
--	--------------	-------------------------------	-----------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 5, line 32	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit	Accession Number PTA-
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Page 17, line 12; page 22, line 10; page 23, line 23; page 108, lines 8, 13, 17, 21, 24 and 29; page 109, lines 9 and 13; page 110, lines 2, 6, 13 and 22; page 111, lines 2, 6, 9 and 13.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indicators are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") Accession Number of Deposit and Date of Deposit	

For receiving Office use only	
<input checked="" type="checkbox"/>	This sheet was received with the international application
Authorized officer	MELVIN S. BROOKS SR. INTERNATIONAL DIVISION 703-805-5169 4/8

For International Bureau use only	
<input type="checkbox"/>	This sheet was received with the International Bureau on:
Authorized officer	

THAT WHICH IS CLAIMED:

1. An isolated nucleic acid molecule selected from the group consisting of:

- a) a nucleic acid molecule comprising a nucleotide sequence
5 which is at least 60% identical to the nucleotide sequence of SEQ ID NO: 2, 4, 6, 8, 11, 12, 13, or 14, or the nucleotide sequence of the cDNA insert of the plasmid deposited with ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____, wherein said nucleotide sequence encodes a polypeptide having biological activity;
- 10 b) a nucleic acid molecule comprising a fragment of at least 20 nucleotides of the nucleotide sequence of SEQ ID NO: 2, 4, 6, 8, 11, 12, 13, or 14, or the nucleotide sequence of the cDNA insert of the plasmid deposited with ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____;
- c) a nucleic acid molecule which encodes a polypeptide
15 comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____;
- d) a nucleic acid molecule which encodes a fragment of a
20 polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, 7, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:1, 3, 5, or 7, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____;
- 25 e) a nucleic acid molecule which encodes a naturally occurring allelic variant of a biologically active polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____, wherein the nucleic acid molecule hybridizes
30 to a nucleic acid molecule comprising the complement of SEQ ID NO:2, 4, 6, 8, 11, 12, 13, or 14 under stringent conditions; and

f) a nucleic acid molecule comprising the complement of a), b), c), d), or e).

2. The isolated nucleic acid molecule of claim 1, which is selected from the group consisting of:

a) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:2, 4, 6, 8, 11, 12, 13, 14, the cDNA insert of any one the plasmids deposited with ATCC as Patent Deposit Number _____, PTA-1639, PTA-1846, or _____, or a complement thereof; and

10 b) a nucleic acid molecule which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Number _____, PTA-1639, PTA-1846, or _____.

15 3. The nucleic acid molecule of claim 1 further comprising vector nucleic acid sequences.

4. The nucleic acid molecule of claim 1 further comprising nucleic acid sequences encoding a heterologous polypeptide.

20

5. A host cell which contains the nucleic acid molecule of claim 1.

6. The host cell of claim 5 which is a mammalian host cell.

25 7. A nonhuman mammalian host cell containing the nucleic acid molecule of claim 1.

8. An isolated polypeptide selected from the group consisting of:

30 a) a biological active polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 60% identical to a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 2, 4, 6, 8, 11, 12, 13,

or 14 or the nucleotide sequence of the cDNA insert of the plasmid deposited with ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____;

5 b) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising the complement of SEQ ID NO: 2, 4, 6, 8, 11, 12, 13, or 14 under stringent conditions; and,

10 c) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:1, 3, 5, or 7; and

15 d) a polypeptide having at least 60% sequence identity to the amino acid sequence SEQ ID NO:1, 3, 5, or 7, wherein the polypeptide has biological activity.

9. The isolated polypeptide of claim 8 comprising the amino acid
20 sequence of SEQ ID NO:1, 3, 5, or 7, or an amino acid sequence encoded by the cDNA insert of any of the plasmids deposited with ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____.

10. The polypeptide of claim 8 further comprising heterologous amino
25 acid sequences.

11. An antibody which selectively binds to a polypeptide of claim 8.

12. A method for producing a polypeptide selected from the group
30 consisting of:

 a) a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or the amino acid sequence encoded by the cDNA insert of the

plasmid deposited with the ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____;

5 b) a polypeptide comprising a fragment of the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:1, 3, 5, or 7, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____;

10 c) a biologically active naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC as Patent Deposit Number ____, PTA-1639, PTA-1846, or ____, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising the complement of SEQ ID NO:2, 4, 6, 8, 11, 12, 13, or 14;

15 d) a polypeptide having at least 60% sequence identity to the amino acid sequence of SEQ ID NO:1, 3, 5, or 7, wherein said polypeptide has biological activity; comprising culturing the host cell of claim 5 under conditions in which the nucleic acid molecule is expressed.

13. The method of claim 12 wherein said polypeptide comprises the amino acid sequence of SEQ ID NO:1, 3, 5, or 7.

25 14. A method for detecting the presence of a polypeptide of claim 8 in a sample, comprising:

 a) contacting the sample with a compound which selectively binds to a polypeptide of claim 8; and

30 b) determining whether the compound binds to the polypeptide in the sample.

15. The method of claim 14, wherein the compound which binds to the polypeptide is an antibody.

16. A kit comprising a compound which selectively binds to a polypeptide
5 of claim 8 and instructions for use.

17. A method for detecting the presence of a nucleic acid molecule of claim 1 in a sample, comprising the steps of:

- a) contacting the sample with a nucleic acid probe or primer
10 which selectively hybridizes to the nucleic acid molecule; and
- b) determining whether the nucleic acid probe or primer binds to a nucleic acid molecule in the sample.

18. The method of claim 17, wherein the sample comprises mRNA
15 molecules and is contacted with a nucleic acid probe.

19. A kit comprising a compound which selectively hybridizes to a nucleic acid molecule of claim 1 and instructions for use.

20. A method for identifying a compound which binds to a polypeptide of claim 8 comprising the steps of:

- a) contacting a polypeptide, or a cell expressing a polypeptide of claim 8 with a test compound; and
- b) determining whether the polypeptide binds to the test
25 compound.

21. The method of claim 20, wherein the binding of the test compound to the polypeptide is detected by a method selected from the group consisting of:

- a) detection of binding by direct detecting of test
30 compound/polypeptide binding;
- b) detection of binding using a competition binding assay;
- c) detection of binding using an assay for sulfatase activity.

22. A method for modulating the activity of a polypeptide of claim 8 comprising contacting a polypeptide or a cell expressing a polypeptide of claim 8 with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide.

23. A method for identifying a compound which modulates the activity of a polypeptide of claim 8, comprising:

- a) contacting a polypeptide of claim 8 with a test compound; and
- b) determining the effect of the test compound on the activity of the polypeptide to thereby identify a compound which modulates the activity of the polypeptide.

24. A method for identifying an agent that modulates the level of expression of a nucleic acid molecule of claim 1 in a cell, said method comprising contacting said agent with the cell expressing said nucleic acid molecule such that said level of expression of said nucleic acid molecule can be modulated in said cell by said agent and measuring said level of expression of said nucleic acid molecule.

25. A method for modulating the level of expression of a nucleic acid molecule of claim 1, said method comprising contacting said nucleic acid molecule with an agent under conditions that allow the agent to modulate the level of expression of the nucleic acid molecule.

26. A pharmaceutical composition containing any of the polypeptides in claim 8 in a pharmaceutically acceptable carrier.

Sequence length 2175

CACGCGTCGCGCAAATTTCTGATTCTTTTGAATTAGGATTCCAGATGGGGGCGCTCATTTCTACAGCCCCCAACATTCTCT
ATAGCCGTTATCACTGOCATCACCACTGCCACCAGCATCTTCTTGAGATTCCACCCCTGCTCCCCAGAGACTTCTCTGC
TTTGAAAGTGAGCAGAAAGGAAGCTCTCAGAAAAATCTCTAGTGGTGGCTGCCGTCGCTCCAGACAATCGGAATCTCTGC
M G W L F L K V L L A G V S F S G 17
CTTCACCACC ATG GGC TGG CTT TTT CTA AAG GTT TTG TTG GCG GGA GTG AGT TTC TCA GGA 51
F L Y P L V D F C I S G K T R G Q K P N 37
TTT CTT TAT OCT CTT GTG GAT TTT TGC ATC AGT GGG AAA ACA AGA GGA CAG AAG CCA AAC 111
F V I I L A D D M G W G D L G A N W A E 57
TTT GTG ATT ATT TTG GCC GAT GAC ATG GGG TGG GGT GAC CTG GGA GCA AAC TGG GCA GAA 171
T K D T A N L D K M A S E G M R F V D F 77
ACA AAG GAC ACT GCC AAC CTT GAT AAG ATG GCT TCG GAG GGA ATG AGG TTT GTG GAT TTC 231
H A A A S T C S P S R A S L L T G R L G 97
CAT GCA GCT GCC TCC ACC TGC TCA CCC TCC CGG GCT TCC TTG CTC ACC GGC CGG CTT GGC 291
L R N G V T R N F A V T S V G G L P L N 117
CTT CGC AAT GGA GTC ACA CGC AAC TTT GCA GTC ACT TCT GTG GGA GGC CTT CCG CTC AAC 351
E T T L A E V L Q Q A G Y V T G I I G K 137
GAG ACC ACC TTG GCA GAG GTG CTG CAG CAG GCG GGT TAC GTC ACT GGG ATA ATA GGC AAA 411
W H L G H H G S Y H P N F R G F D Y Y F 157
TGG CAT CTT GGA CAC CAC GGC TCT TAT CAC CCC AAC TTC CGT GGT TTT GAT TAC TAC TTT 471
G I P Y S H D M G C T D T P G Y N H P P 177
GGA ATC CCA TAT AGC CAT GAT ATG GGC TGT ACT GAT ACT CCA GGC TAC AAC CAC CCT CCT 531
C P A C P Q G D G P S R N L Q R D C Y T 197
TGT CCA GCG TGT CCA CAG GGT GAT GGA CCA TCA AGG AAC CTT CAA AGA GAC TGT TAC ACT 591
D V A L P L Y E N L N I V E Q P V N L S 217
GAC GTG GCC CTC CCT CTT TAT GAA AAC CTC AAC ATT GTG GAG CAG CCG GTG AAC TTG AGC 651
S L A Q K Y A E K A T Q F I Q R A S T S 237
AGC CTT GCC CAG AAG TAT GCT GAG AAA GCA ACC CAG TTC ATC CAG CGT GCA AGC ACC AGC 711
G R P F L L Y V A L A H M H V P L P V T 257
GGG AGG CCC TTC CTG CTC TAT GTG GCT CTG GCC CAC ATG CAC GTG CCC TTA CCC GTG ACT 771
Q L P A A P R G R S L Y G A G L W E M D 277
CAG CTA CCA GCA GCG CCA CGG GGC AGA AGC CTG TAT GGT GCA GGG CTC TGG GAG ATG GAC 831
S L V G Q I K D K V D H T V K E N T F L 297
AGT CTG GTG GGC CAG ATC AAG GAC AAA GTT GAC CAC ACA GTG AAG GAA AAC ACA TTC CTC 891
W F T G D N G P W A Q K C E L A G S V G 317
TGG TTT ACA GGA GAC AAT GGC CCG TGG GCT CAG AAG TGT GAG CTA GCG GGC AGT GTG GGT 951
P F T G F W Q T R Q G G S P A K Q T T W 337
CCC TTC ACT GGA TTT TGG CAA ACT CGT CAA GGG GGA AGT CCA GCC AAG CAG ACG ACC TGG 1011
E G G H R V P A L A Y W P G R V P V N V 357

FIGURE 1A

GAA GGA GGG CAC CGG GTC CCA GCA CTG GCT TAC TGG CCT GGC AGA GTT CCA GTT AAT GTC 1071
 T S T A L L S V L D I F P T V V A L A Q 377
 ACC AGC ACT GCC TTG TTA AGC GTG CTG GAC ATT TTT CCA ACT GTG GTA GCC CTG GCC CAG 1131
 A S L P Q G R R F D G V D V S E V L F G 397
 GCC AGC TTA CCT CAA GGA CGG CGC TTT GAT GGT GTG GAC GTC TCC GAG GTG CTC TTT GGC 1191
 R S Q P G H R V L F H P N S G / A A G E F 417
 CGG TCA CAG OCT GGG CAC AGG GTG CTG TTC CAC CCC AAC AGC GGG GCA GCT GGA GAG TTT 1251
 G A L Q T V R L E R Y K A F Y I T G G A 437
 GGA GCC CTG CAG ACT GTC CGC CTG GAG CGT TAC AAG GCC TTC TAC ATT ACC GGT GGA GCC 1311
 R A C D G S T G P E L Q H K F P L I F N 457
 AGG GCG TGT GAT GGG AGC ACG GGG CCT GAG CTG CAG CAT AAG TTT CCT CTG ATT TTC AAC 1371
 L E D D T A E A V P L E R G G A E Y Q A 477
 CTG GAA GAC GAT ACC GCA GAA GCT GTG CCC CTA GAA AGA GGT GGT GCG GAG TAC CAG GCT 1431
 V L P E V R K V L A D V L Q D I A N D N 497
 GTG CTG CCC GAG GTC AGA AAG GTT CTT GCA GAC GTC CTC CAA GAC ATT GCC AAC GAC AAC 1491
 I S S A D Y T Q D P S V T P C C N P Y Q 517
 ATC TCC AGC GCA GAT TAC ACT CAG GAC CCT TCA GTA ACT CCC TGC TGT AAT CCC TAC CAA 1551
 I A C R C Q A A * 526
 ATT GCC TGC CGC TGT CAA GCC GCA TAA 1578
 CAGACCAATTTTATTCACGAGGAGGAGTACCTGGAAATTAGGCAAGTTTGCTTCCAAATTTTCATTTTACCCCTCTTT
 ACAACACACACGCTTTAGTTTAGTCTTGGAGTTTAGTTTGGAGTTAGCCCTGCAATATCCCTTCTGTATCCTGTCCCTCC
 TOCACGCGGACCGAGAGCAGCTGAGCTGCGCTGGCTCTGGGCAGGAGTGTGCCCTTAATGGGAAGCACACGGGCTTTG
 GAGTCAGGCACAGGTGCCAGCTCCAGCTTTGAACTTGGGCAATTGTTTAACCTAACCTGCAAGTTGATTTTGAGGGTT
 AAATAAAGGCATACATGAAAAAAAAAAAAAAAAA

FIGURE 1B




FIGURE 2




FIGURE 3

Prosite Pattern Matches

Prosite version: Release 12.1 of February 1995.

>PS00001|PDOC00001|ASN_GLYCOSYLATION N-glycosylation site.

Query: 117 NETT 120

Query: 215 NLSS 218

Query: 356 NVTG 359

Query: 497 NISS 500

>PS00005|PDOC00005|PKC_PHOSPHO_SITE Protein kinase C phosphorylation site.

Query: 28 SGK 30

Query: 93 TGR 95

Query: 237 SGR 239

Query: 290 TVK 292

Query: 422 TVR 424

>PS00006|PDOC00006|CK2_PHOSPHO_SITE Casein kinase II phosphorylation site.

Query: 120 TLAE 123

Query: 290 TVKE 293

Query: 335 TTWE 338

Query: 364 SVLD 367

Query: 444 TGPE 447

Query: 499 SSAD 502

>PS00008|PDOC00008|MYRISTYL N-myristoylation site.

Query: 12 GVSFSG 17

Query: 33 GQKPNF 38

Query: 52 GANWAE 57

Query: 97 GLRNGV 102

Query: 113 GLPLNE 118

Query: 158 GIPYSH 163

Query: 328 GGSPAK 333

Query: 388 GVDVSE 393

Query: 418 GALQTV 423

Query: 435 GCARAC 440

>PS00009|PDOC00009|AMIDATION Amidation site.

Query: 382 QGRR 385

>PS00142|PDOC00117|SULFATASE_2 Sulfatases signature 2.

Query: 129 GVTGIGRM 130

FIGURE 4

Input file Fbh23553f1.seq; Output File 23553.trans
Sequence length 4321

```

CCCAOOGGTCCGGCTAATGAATCTTGGGGCCGGTGTGGGGCCGGGGGGCTTGATCGGCAACTAGGAAACCCAGGGGC
AGAGGCCAGGAGCGAGGGCAGGGATCAGAGGCCAGGCTTCCGGCTGCGGGCGCTCTCGAGGTCAGGGCAGAT
GAGGAACATGACTCTCCCCCTTGGAGGAGGAAGGAAGTCCGGCTGCCAOCCTTATCTCTGCTCTCTGCTCTCTCCCTG
TTCCAGAGCTTTTCTCTAGAGAAGATTTTGAAGCGGCTTTTGTGCTGACGGGCCACCCACCATCATCTAAAGAAGAT
AAACTTGGCAAATGACATGCAGGTTCTTCAAGGCAGAAATAATTCAGAAAACTTCAAGGACCCCTATCTGCAGATGTT
CTGAATACCTCTGAGAATAGAGATTGATTATTCAACCAGGATACCTAATTCAAGAACTCCAGAAATCAGGAGACGGAGA
                                     M K Y S C C A L V L A      11
CATTTTGTGAGTTTGTCAACATTGGACCAAATACA ATG AAG TAT TCT TGC TGT GCT CTG GTT TTG GCT      33
V L G T E L L G S L C S T V R S P R F R      31
GTC CTG GGC ACA GAA TTG CTG GGA AGC CTC TGT TCG ACT GTC AGA TCC CCG AGG TTC AGA      93
G R I Q Q E R K N I R P N I I L V L T D      51
GGA CCG ATA CAG CAG GAA CGA AAA AAC ATC CGA CCC AAC ATT ATT CTT GTG CTT ACC GAT      153
D Q D V E L G S L Q V M N K T R K I M E      71
GAT CAA GAT GTG GAG CTG GGG TCC CTG CAA GTC ATG AAC AAA ACG AGA AAG ATT ATG GAA      213
H G G A T F I N A F V T T P M C C P S R      91
CAT GGG GGG GCC ACC TTC ATC AAT GCC TTT GTG ACT ACA CCC ATG TGC TGC CCG TCA CGG      273
S S M L T G K Y V H N H N V Y T N N E N      111
TCC TCC ATG CTC ACC GGS AAG TAT GTG CAC AAT CAC AAT GTC TAC ACC AAC AAC GAG AAC      333
C S S P S W Q A M H E P R T F A V Y L N      131
TGC TCT TCC CCC TCG TGG CAG GCC ATG CAT GAG CCT CGG ACT TTT GCT GTA TAT CTT AAC      393
N T G Y R T A F F G K Y L N E Y N G S Y      151
AAC ACT GGC TAC AGA ACA GCC TTT TTT GGA AAA TAC CTC AAT GAA TAT AAT GGC AGC TAC      453
I P P G W R E W L G L I K N S R F Y N Y      171
ATC CCC CCT GGG TGG CGA GAA TGG CTT GGA TTA ATC AAG AAT TCT CCG TTC TAT AAT TAC      513
T V C R N G I K E K H G F D Y A K D Y F      191
ACT GTT TGT CGC AAT GGC ATC AAA GAA AAG CAT GGA TTT GAT TAT GCA AAG GAC TAC TTC      573
T D L I T N E S I N Y F K M S K R M Y P      211
ACA GAC TTA ATC ACT AAC GAG AGC ATT AAT TAC TTC AAA ATG TCT AAG AGA ATG TAT CCC      633
H R P V M M V I S H A A P H G P E D S A      231
CAT AGG CCC GTT ATG ATG GTG ATC AGC CAC GCT GCG CCC CAC GGC CCC GAG GAC TCA GCC      693
P Q F S K L Y P N A S Q H I T P S Y N Y      251
CCA CAG TTT TCT AAA CTG TAC CCC AAT GCT TCC CAA CAC ATA ACT CCT AGT TAT AAC TAT      753
A P N M D K H W I M Q Y T G P M L P I H      271
GCA CCA AAT ATG GAT AAA CAC TGG ATT ATG CAG TAC ACA GGA CCA ATG CTG CCC ATC CAC      813
M E F T N I L Q R K R L Q T L M S V D D      291
ATG GAA TTT ACA AAC ATT CTA CAG CGC AAA AGG CTC CAG ACT TTG ATG TCA GTG GAT GAT      873
S V E R L Y N M L V E T G E L E N T Y I      311

```

FIGURE 5A

TCT GTG GAG AGG CTG TAT AAC ATG CTC GTG GAG ACG GGG GAG CTG GAG AAT ACT TAC ATC 933
 I Y T A D H G Y H I G Q F G L V K G K S 331
 ATT TAC ACC GCC GAC CAT GGT TAC CAT ATT GGG CAG TTT GGA CTG GTC AAG GGG AAA TCC 993
 M P Y D F D I R V P F F I R G P S V E P 351
 ATG CCA TAT GAC TTT GAT ATT CGT GTG CCT TTT TTT ATT CGT GGT CCA AGT GTA GAA CCA 1053
 G S I V P Q I V L N I D L A P T I L D I 371
 GGA TCA ATA GTC CCA CAG ATC GTT CTC AAC ATT GAC TTG GCC CCC ACG ATC CTG GAT ATT 1113
A G L D T P P D V D G K S V L K L L D P 391
 GCT GGG CTC GAC ACA CCT CCT GAT GTG GAC GGC AAG TCT GTC CTC AAA CTT CTG GAC CCA 1173
 E K P G N R F R T N K K A K I W R D T F 411
 GAA AAG CCA GGT AAC AGG TTT CGA ACA AAC AAG AAG GCC AAA ATT TGG CGT GAT ACA TTC 1233
 L V E R G K F L R K K E E S S K N I Q Q 431
 CTA GTG GAA AGA GGC AAA TTT CTA CGT AAG AAG GAA GAA TCC AGC AAG AAT ATC CAA CAG 1293
 S N H L P K Y E R V K E L C Q Q A R Y Q 451
 TCA AAT CAC TTG CCC AAA TAT GAA CGG GTC AAA GAA CTA TGC CAG CAG GCC AGG TAC CAG 1353
 T A C E Q P G Q K W Q C I E D T S G K L 471
 ACA GCC TGT GAA CAA CCG GGG CAG AAG TGG CAA TGC ATT GAG GAT ACA TCT GGC AAG CTT 1413
 R I H K C K G P S D L L T V R Q S T R N 491
 CGA ATT CAC AAG TGT AAA GGA CCC AGT GAC CTG CTC ACA GTC CGG CAG AGC ACG CGG AAC 1473
 L Y A R G F H D K D K E C S C R E S G Y 511
 CTC TAC GCT CGC GGC TTC CAT GAC AAA GAC AAA GAG TGC AGT TGT AGG GAG TCT GGT TAC 1533
 R A S R S Q R K S Q R Q F L R N Q G T P 531
 CGT GCC AGC AGA AGC CAA AGA AAG AGT CAA CGG CAA TTC TTG AGA AAC CAG GGG ACT CCA 1593
 K Y K P R F V H T R Q T R S L S V E F E 551
 AAG TAC AAG CCC AGA TTT GTC CAT ACT CGG CAG ACA CGT TCC TTG TCC GTC GAA TTT GAA 1653
 G E I Y D I N L E E E E E L Q V L Q P R 571
 GGT GAA ATA TAT GAC ATA AAT CTG GAA GAA GAA GAA GAA TTG CAA GTG TTG CAA CCA AGA 1713
 N I A K R H D E G H K G P R D L Q A S S 591
 AAC ATT GCT AAG CGT CAT GAT GAA GGC CAC AAG GGG CCA AGA GAT CTC CAG GCT TCC AGT 1773
 G G N R G R M L A D S S N A V G P P T T 611
 GGT GGC AAC AGG GGC AGG ATG CTG GCA GAT AGC AGC AAC GCC GTG GGC CCA CCT ACC ACT 1833
 V R V T H K C F I L P N D S I H C E R E 631
 GTC CGA GTG ACA CAC AAG TGT TTT ATT CTT CCC AAT GAC TCT ATC CAT TGT GAG AGA GAA 1893
 L Y Q S A R A W K D H K A Y I D K E I E 651
 CTG TAC CAA TCG GCC AGA GCG TGG AAG GAC CAT AAG GCA TAC ATT GAC AAA GAG ATT GAA 1953
 A L Q D K I K N L R E V R G H L K R R K 671
 GCT CTG CAA GAT AAA ATT AAG AAT TTA AGA GAA GTG AGA GGA CAT CTG AAG AGA AGG AAG 2013
 P E E C S C S K Q S Y Y N K E K G V K K 691
 CCT GAG GAA TGT AGC TGC AGT AAA CAA AGC TAT TAC AAT AAA GAG AAA GGT GTA AAA AAG 2073
 Q E K L K S H L H P F K E A A Q E V D S 711
 CAA GAG AAA TTA AAG AGC CAT CTT CAC CCA TTC AAG GAG GCT GCT CAG GAA GTA GAT AGC 2133

FIGURE 5B

K L Q L F K E N N R R R K K E R K E K R 731
 AAA CTG CAA CTT TTC AAG GAG AAC AAC CGT AGG AGG AAG AAG GAG AGG AAG GAG AAG AGA 2193
 R Q R K G E E C S L P G L T C F T H D N 751
 CGG CAG AGG AAG GGG GAA GAG TGC AGC CTG CCT GGC CTC ACT TGC TTC ACG CAT GAC AAC 2253
 N H W Q T A P F W N L G S F C A C T S S 771
 AAC CAC TGG CAG ACA GCC CCG TTC TGG AAC CTG GGA TCT TTC TGT GCT TGC ACG AGT TCT 2313
 N N N T Y W C L R T V N E T H N F L F C 791
 AAC AAT AAC ACC TAC TGG TGT TTG CGT ACA GTT AAT GAG ACG CAT AAT TTT CTT TTC TGT 2373
 E F A T G F L E Y F D M N T D P Y Q L T 811
 GAG TTT GCT ACT GGC TTT TTG GAG TAT TTT GAT ATG AAT ACA GAT CCT TAT CAG CTC ACA 2433
 N T V H T V E R G I L N Q L H V Q L M E 831
 AAT ACA GTG CAC ACG GTA GAA CGA GGC ATT TTG AAT CAG CTA CAC GTA CAA CTA ATG GAG 2493
 L R S C Q G Y K Q C N P R P K N L D V G 851
 CTC AGA AGC TGT CAA GGA TAT AAG CAG TGC AAC CCA AGA CCT AAG AAT CTT GAT GTT GGA 2553
 N K D G G S Y D L H R G Q L W D G W E G 871
 AAT AAA GAT GGA GGA AGC TAT GAC CTA CAC AGA GGA CAG TTA TGG GAT GGA TGG GAA GGT 2613
 * 872
 TAA 2616
 TCAGCCCCGCTCTCACTGCAGACATCAACTGGCAAGGCCCTAGAGGAGCTACACAGTGTGAATGAAAACATCTATGAGTAC
 AGACAAACTACAGACTTAGTCTGGTGGACTGGACTAATTACTTTGAAGGATTTAGATAGAGTATTTGCACTGCTGAAGA
 GTCCTATGAGCAAAATAAAAACAATAAGACTCAAAGTCTCAAAGTGACGGGTCTTGGTTGTCTCTGCTGAGCACGC
 TGTGTCAATGGAGATGGCCTCTGCTGACTCAGATGAAGACCCAAAGGCATAAGGTGGGAAAACACCTCATTTGACCTTG
 CCAGCTGACCTTCAAACCTGCATTTGAACCGACCAACATTAAAGTCCAGAGAGTAAACTTGAATGGAATAACGACATTC
 CAGAAGTTAATCATTTGAATTCTGAACACTGGAGAAAAAACCAGAAAAATGGAACGGGCAATGAAGAGACTAATCATCTGGA
 AACCGATTTTCACTGGCGATGGCATGACAGAGCTAGAGCTGGGGCCAGGCCAGGCTGCAGCCCATTCGAGGCACCCG
 AAAGAACTTCCCCAGTATGGTGGTCTGGAAAGGACATTTTGAAGATCAACTATATCTTCTGTGCATTCGGATGGAA
 TTTCACTTCATCAGATGTTTCAACATGGCCACCGCAGAACACCGAAGTAATTCAGCATAGCGGGGAAGATGTTGACCAA
 GGTGGAGAAGAATCAOGAAAAGGAGAAGTCACAGCACTAGAAAGGCAGCGCTCTCTTCACTCTCTCTGATTAGATG
 AAACGTGTTAOCCTTAOCCTAATACACAGTATTTCTTTTAACTTTTATTTGTAAACTAATAAGGKAATCACAGCCACC
 AACATTCCAAGCTAOCCTGGGTACCTTTGTGCACTAGAAGCTAGTGAGCATGTGAGCAAGCGGTGTGCACACCGAGACT
 CATCGTTATAATTTACTATCTGCAAGGAGTAGAAGAAAGGCTGGGGATATTTGGGTGGCTTTGGKTTTGATTTTPT
 GCTTGGTTGGTTGGTTTGGKACTAATAACAGTATTATCTTTGAATATCGTAGGGACATAARKWWWMMWKKTWWTCMAW
 YMRKAKGSYWRRAWKGGSTYTYTSKKRKSTWAMWYKWSMOCYSKKRWAWTYWWWYWCYKYTSSTGRYKRN
 KTAATGAAGTT

FIGURE 5C

Analysis of 23553 (871 aa)




FIGURE 6




FIGURE 7

Prosit Pattern Matches for 23553

Prosit version: Release 12.2 of February 1995

>PS00001|PDOC00001|ASN_GLYCOSYLATION N-glycosylation site.

Query: 64	NKTR	67
Query: 111	NCSS	114
Query: 131	NNTG	134
Query: 148	NGSY	151
Query: 170	NYTV	173
Query: 197	NESI	200
Query: 240	NASQ	243
Query: 623	NDSI	626
Query: 773	NNTY	776
Query: 783	NETH	786

>PS00005|PDOC00005|PKC_PHOSPHO_SITE Protein kinase C phosphorylation site.

Query: 24	TVR	26
Query: 27	SPR	29
Query: 66	TRK	68
Query: 96	TGK	98
Query: 206	SKR	208
Query: 400	TNK	402
Query: 425	SSK	427
Query: 468	SGK	470
Query: 484	TVR	486
Query: 488	STR	490
Query: 505	SCR	507
Query: 516	SQR	518
Query: 520	SQR	522
Query: 530	TPK	532
Query: 611	TVR	613
Query: 615	TRK	617
Query: 635	SAR	637

>PS00006|PDOC00006|CK2_PHOSPHO_SITE Casein kinase II phosphorylation site.

Query: 107	TNNE	110
Query: 288	SVDD	291
Query: 367	TILO	370
Query: 376	TPPD	379
Query: 452	TACE	455
Query: 505	SCRE	508
Query: 781	TVNC	784

FIGURE 8A

>PS00007|PDOC00007|TYR_PHOSPHO_SITE Tyrosine kinase phosphorylation site.

Query: 637 RAKKDHKAY 645

>PS00002|PDOC00008|MYRISTYL N-myristoylation site.

Query: 19 GSLCST 24

Query: 161 GLIKNS 166

Query: 325 GLVKGK 330

Query: 592 GGNRGR 597

Query: 763 GSFCAC 768

Query: 851 GNKDGG 856

>PS00521|PDOC00117|SULFATASE_1 Sulfatases signature 1.

Query: 85 PMCCPSRSSMLTG 97

FIGURE 8B

23553 Oncology Taqman




FIGURE 9

Input file Fbh25278FL1.seq; Output File 25278.trans
Sequence length 2940

```

CCACGGCTCCGCCACCGCTCCGGCTGCCACGCCGCTCTCAGGCTGGCCGGGCTGAGCCGGGAAGAGGGAGCAAAGG
CGCGCGAGGGGCTGCGCTTAGGCAGCGGGAGGCAGCTCCGCGCGGGCCCTGACCTCCCCAGAGCGCCCGCTGCGGCCGA
CGAGATCCGGCCCGAGCGCTCCGGCAGCCAGTCCCGGACCAGACACTGGACCGTCCCCGGGGGGCGCTGAACCTCCCTCGC
AGCATCCGAGCCGGCGGGCCCGGTGGTGGCCCTGGCGCGCGAGGTGGTGAGGCCCCAGGAGCCCGCGCGCCGGGACA

      M H T L T G F S L V S L L S F      15
CGCGGGCCGGCTGGCG ATG CAC ACC CTC ACT GGC TTC TCT CTG GTC AGC CTG CTC AGC TTC      45

  G Y L S W D W A K P S F V A D G P G E A      35
GGC TAC CTG TCC TGG GAC TGG GCC AAG CCG AGC TTC GTG GCC GAC GGG CCC GGG GAG GCT      105

  G E Q P S A A P P Q P P H I I F I L T D      55
GGC GAG CAG CCC TCG GCC GCT CCG CCC CAG CCT CCC CAC ATC ATC TTC ATC CTC ACG GAC      165

  D Q G Y H D V G Y H G S D I E T P T L D      75
GAC CAA GGC TAC CAC GAC GTG GGC TAC CAT GGT TCA GAT ATC GAG ACC OCT ACG CTG GAC      225

  R L A A K G V K L E N Y Y I Q P I C T P      95
AGG CTG GCG GCC AAG GGG GTC AAG TTG GAG AAT TAT TAC ATC CAG CCC ATC TGC ACG CCT      285

  S R S Q L L T G R Y Q I H T G L Q H S I      115
TCG CGG AGC CAG CTC CTC ACT GGC AGG TAC CAG ATC CAC ACA GGA CTC CAG CAT TCC ATC      345

  I R P Q Q P N C L P L D Q V T L P Q K L      135
ATC CGC CCA CAG CAG CCC AAC TGC CTG CCC CTG GAC CAG GTG ACA CTG CCA CAG AAG CTG      405

  Q E A G Y S T H M V G K W H L G F Y R K      155
CAG GAG GCA GGT TAT TCC ACC CAT ATG GTG GGC AAG TGG CAC CTG GGC TTC TAC CGG AAG      465

  E C L P T R R G F D T F L G S L T G N V      175
GAG TGT CTG CCC ACC CGT CGG GGC TTC GAC ACC TTC CTG GGC TCG CTC ACG GGC AAT GTG      525

  D Y Y T Y D N C D G P G V C G F D L H E      195
GAC TAT TAC ACC TAT GAC AAC TGT GAT GGC CCA GGC GTG TGC GGC TTC GAC CTG CAC GAG      585

  G E N V A W G L S G Q Y S T M L Y A Q R      215
GGT GAG AAT GTG GCC TGG GGG CTC AGC GGC CAG TAC TCC ACT ATG CTT TAC GGC CAG CGC      645

  A S H I L A S H S P Q R P L F L Y V A F      235
GOC AGC CAT ATC CTG GOC AGC CAC AGC CCT CAG CGT CCC CTC TTC CTC TAT GTG GOC TTC      705

  Q A V H T P L Q S P R E Y L Y R Y R T M      255
CAG GCA GTA CAC ACA CCC CTG CAG TCC CCT CGT GAG TAC CTG TAC CGC TAC CGC ACC ATG      765

  G N V A R R K Y A A M V T C M D E A V R      275
GGC AAT GTG GCC CGG CGG AAG TAC GCG GCC ATG GTG ACC TGC ATG GAT GAG GCT GTG CGC      825

  N I T W A L K R Y G F Y N N S V I I F S      295
AAC ATC ACC TGG GCC CTC AAG CGC TAC GGT TTC TAC AAC AAC AGT GTC ATC ATC TTC TCC      885

  S D N G G Q T F S G G S N W P L R G R K      315
AGT GAC AAT GGT GGC CAG ACT TTC TCG GGG GGC AGC AAC TGG CCG CTC CGA GGA CGC AAG      945

  G T Y W E G G V R G L G F V H S P L L K      335
GGC ACT TAT TGG GAA GGT GGC GTG CGG GGC CTA GGC TTT GTC CAC AGT CCC CTG CTC AAG      1005

```

FIGURE 10A

R K Q R T S R A L M H I T D W Y P T L V 355
 CCA AAG CAA CGG ACA AGC CGG GCA CTG ATG CAC ATC ACT GAC TGG TAC CCG ACC CTG GTG 1065
 G L A G G T T S A A D G L D G Y D V W P 375
 GGT CTG GCA GGT GGT ACC ACC TCA GCA GCC GAT GGG CTA GAT GGC TAC GAC GTG TGG CCG 1125
 A I S E G R A S P R T E I L H N I D P L 395
 GCC ATC AGC GAG GGC CGG GCC TCA CCA CGC ACG GAG ATC CTG CAC AAC ATT GAC CCA CTC 1185
 Y N H A Q H G S L E G G F G I W N T A V 415
 TAC AAC CAT GCC CAG CAT GGC TCC CTG GAG GGC GGC TTT GGC ATC TGG AAC ACC GCC GTG 1245
 Q A A I R V G E W K L L T G D P G Y G D 435
 CAG GCT GCC ATC CGC GTG GGT GAG TGG AAG CTG CTG ACA GGA GAC CCC GGC TAT GGC GAT 1305
 W I P P Q T L A T P P G S W W N L E R M 455
 TGG ATC CCA CCG CAG ACA CTG GCC ACC TTC CCG GGT AGC TGG TGG AAC CTG GAA CGA ATG 1365
 A S V R Q A V W L F N I S A D P Y E R E 475
 GCC AGT GTC CGC CAG GCC GTG TGG CTC TTC AAC ATC AGT GCT GAC CCT TAT GAA CCG GAG 1425
 D L A G Q R P D V V R T L L A R L A E Y 495
 GAC CTG GCT GGC CAG CGG CCT GAT GTG GTC CGC ACC CTG CTG GCT CGC CTG GCC GAA TAT 1485
 N R T A I P V R Y P A E N P R A H P D F 515
 AAC CGC ACA GCC ATC CCG GTA CGC TAC CCA GCT GAG AAC CCC CCG GCT CAT CCT GAC TTT 1545
 N G G A W G P W A S D E E E E E E E G R 535
 AAT GGG GGT GCT TGG GGG CCC TGG GCC AGT GAT GAG GAA GAG GAG GAA GAG GAA GGG AGG 1605
 A R S F S R G R R K K K C K I C K L R S 555
 GCT CGA AGC TTC TCC CGG GGT CGT CGC AAG AAA AAA TGC AAG ATT TGC AAG CTT CGA TCC 1665
 F F R K L N T R L M S Q R I * 570
 TTT TTC CGT AAA CTC AAC ACC AGG CTA ATG TCC CAA CGG ATC TGA 1710
 TGGTGGGGAGGGAGAAAACTGTCTTTAGAGGATCTTCCCACTCCGGCTTGGCCCTGCTGTTTCTCAGGGAGAAGCCT
 GTACATCTCCATCTACAGGGAGTTGGAGGGGTAGAGTCCCTTGGTTGAACAGGGTAGGGAGCCTGGATAGGAGTGGG
 TGGGAATAAACAGACTGGGATGCTGTCTCAGTCTGCTCTCAAGGACTTGCTCTGTGACCTCAGGTGACCCAC
 ATGAGCTTTTAGCCTCAGTTTCTTCATCTGTAAAAATGAGCTCTAATGACTTTGTGACTCTTTGGTGTGGCCCTGGAGCC
 TGGGGCCAGGGTGGAGTTCTTGGCCGGCCTTGCCACTTGACAACTCTTTAAGGCTTCCCTTTAACAGGGATCCCTG
 TGGTGGTGTTTGGGAGTTGCCCTGGAGGCACTCCAAGCCTGGCCCCAGCTGAAGCATGGCAATCTGGCTGCTCTAC
 AGGGACCCCCAAGCGCTGTGGGTGGAGGGCAGGGGTGGGGGGGTGACCTTCTTGGGTCTTCACATGGCCTAGGCCAG
 TCTTCCGGTCAGACTGGTGTGAGGCAAGGTGGTGCAAAATTCCTCTTCTGGGCCCTCCAGTACCCAGAGAACTGGCTG
 GGCCATTAACTGCTGCAGCAOCAAAGGGTGGTAGAAAGAGCTGTGAAGAGCCCCCAAACAGTACCAGGACACCTGGGTT
 CTCTGTGAOCTGGGGCAGTTCTTGGCCCTTAGGCCTTGATTTCCCCACCTGCAAGTGGGGATGCCAGCCCTGGCTC
 TGCTCTCTCATGAGGCTCTGGAAGACTGGCCAAGGTGTGGAGGAGCTGTGAACTTGATTAAAGTGTGTAACATGG
 AAAAAAAAAAAAAAAAAAAAAAGGGGG

FIGURE 10B

Analysis of 25278 (569 aa)




FIGURE 11




FIGURE 12

Version: Release 11.2 of February 1995

101|PDOC000001|ASN_GLYCOSYLATION N-glycosylation site.

276	NITW	279
288	NISV	291
466	NISA	469
496	NRTA	499

104|PDOC000004|CAMP_PHOSPHO_SITE cAMP- and cGMP-dependent protein kinase phosphorylation site.

314	RKGT	317
-----	------	-----

105|PDOC000005|PKC_PHOSPHO_SITE Protein kinase C phosphorylation site.

102	TGR	104
160	TRR	162
244	SPR	246
340	TSR	342
383	SPR	385
457	SVR	459
566	SQR	568

106|PDOC000006|CK2_PHOSPHO_SITE Casein kinase II phosphorylation site.

67	SDIE	70
244	SPRE	247
268	TCMD	271
317	TYWE	320
363	SAAD	366
525	SDEE	528

107|PDOC000007|TYR_PHOSPHO_SITE Tyrosine kinase phosphorylation site.

134	KIQEAGY	140
-----	---------	-----

108|PDOC000008|MYRISTYL N-myristoylation site.

110	GLQHSI	115
169	GSLTGN	174
205	QQYSTK	210
300	QQTFSG	305
321	GGVRGL	326
356	GLAGGT	361
402	GSLEGG	407
409	GIWNTA	414
447	GSHWNL	452

109|PDOC000009|AMIDATION Amidation site.

312	RORK	315
541	RORR	544

112|PDOC001117|SULFATASE_2 Sulfatases signature 2.

139	GVSTHRVGRW	148
-----	------------	-----

21|PDOC001117|SULFATASE_1 Sulfatases signature 1.

91	PICTPSRSQULTG	103
----	---------------	-----

FIGURE 13




FIGURE 14.

26212 seqs

DNA sequence (nt 706-2118 coding)

CACGCGTCCGCCCACGCGTCCGTGGAGATATTAACCTTTTTTCTTTTTTTTTTCTTGGTGGAAGCTGCTCTAGGGAGGGGGGAGGAGGA
GGAGAAAGTGAAATGTGCTGGAGAAGAGCGAGCCCTCCTTGTTCTTCCGGAGTCCCATCCATTAAGCCATCACTTCTGGAAAGATTAAAGT
TGTCGGACATGGTGACAGCTGAGAGGAGAGGAGGATTTCTTGCCAGGTGGAGAGTCTTCACCGTCTGTTGGGTGCATGTGTGCGCCCGCA
GCGGCGCGGGGCGCGTGGTTCTCCGCGTGGAGTCTCACCTGGGACCTGAGTGAATGGCTCCCAGGGGCTGTGCGGGGCATCCGCTCCGC
CTTCTCCACAGGCCGTGTGTCTGTCTGGAAGATGCTAGCAATGGGGGCGCTGGCAGGATTCGATCCTCTGCCTCCTCACTTATGGTT
ACCTGTCTGCGGGCCAGGCCTTAGAAGAGGAGGAAGAGGGGCTTACTAGCTCAAGCTGGAGAGAACTAGAGCCAGCACAACTTCCA
CCTCCCAGCCCCATCTCATTTTCATCCTAGCGGATGATCAGGGATTTAGAGATGTGGGTTACACCGGATCTGAGATTAAACACCTACTC
TTGACAAGCTCGCTGCCGAAGGAGTTAAACTGGGAACTACTATGTCCAGCCTATTTGCACACCATCCAGGAGTCAGTTTATTACTGGAA
AGTATCAGATACACACCGGACTTCAACATTCTATCATAAGACCTACCCAACCACTGTTTACCTCTGGACAATGCCACCTACCTCAGA
AACTGAAGGAGGTTGGATATTCACGCATATGGTCGGAATGGCACTTGGGTTTTTACAGAAAAGAAATGCATGCCCACCAGAAAGAGGAT
TTGATACCTTTTTTGGTTCCTTTTGGGAAGTGGGGATTACTATACACACTACAAATGTGACAGTCTCGGGATGTGTGGCTATGACTTGT
ATGAAAACGACAATGCTGCCTGGGACTATGACAATGGCATATACTCCACACAGATGTACACTCAGAGAGTACAGCAATCTTAGCTTCCC
ATAACCCCAACAAAGCCTATATTTTATATATTGCCTATCAAGCTGTTTATTCACTGCAAGCTCCTGGCAGGTATTTGCAACACTACC
GATCCATTATCAACATAAACAGGAGGAGATATGCTGCCATGCTTTCCTGCTTAGATGAAGCAATCAACAACGTGACATTGGCTCTAAAGA
CTTATGGTTTCTATAACAACAGCATTATCATTTACTCTTCAGATAATGGTGCCAGCCTACGGCAGGAGGAGTAAGTGGCCTCTCAGAG
GTAGCAAAGGAACATATTGGGAAGGAGGGATCCGGGCTGTAGGCTTTGTGCATAGCCCACTTCTGAAAAACAAGGGAACAGTGTGTAAGG
AACTTGTGCACATCACTGACTGGTACCCCACTCTCATTTCACTGGCTGAAGGACAGATTGATGAGGACATCAACTAGATGGCTATGATA
TCTGGGAGACCATAAGTGAGGCTCTTCGCTCACCCGAGTAGATATTTGCATAACATTGACCCCATATACCAAGGCAAAAAATGGCT
CCTGGGCAGCAGGCTATGGGATCTGGAACACTGCAATCCAGTCAGCCATCAGAGTGCAGCACTGGAATTTGCTTACAGGAAATCCTGGCT
ACAGCGACTGGGTCCCCCTCAGTCTTTCAGCAACCTGGGACCGAACCAGTGGCACAATGAACGGATCACCTTGTCAACTGGCAAAAGTG
TATGGCTTTTCAACATCACAGCCGACCCATATGAGAGGGTGGACCTATCTAACAGGTATCCAGGAATCGTGAAGAAGCTCCTACGGAGGC
TCTCACAGTTCAACAAACTGCAGTGCCGGTCAGGTATCCCCCAAGACCCAGAGTAACCTTAGGCTCAATGGAGGGGTCTGGGGAC
CATGGTATAAAGAGGAAACCAAGAAAAAGCAAGCAAAATCAGGCTGAGAAAAAGCAAAAGAAAGCAAAAAAGAAAGAAAGCA
AGCAGAAAGCAGTCTCAGGTTCAACTTGCCATTGAGGTGTTACTTGTGGATAAGCACAATATTTCTGTTTGGTTAAACTTTAATCAGT
TCTTATCTTTCATCTGTTTCTAGGTAAACCAGCAAATTTGGCTCGATAATATCGCTGGCCTAAGCGTCAGGCTGTTTTTCATGCTGTGC
CAC

Protein sequence

MAPRGCAHPPPPSPQACVCPGKMLAMGALAGFWILCLLTYGYLSWGQALEEEEEEGALLAQAGEKLEPSTTSTSQPHLIFILADDQGFRD
VGYHGSEIKTPTLDKLAEGVKLENYVQPICTPSRSQFITGKYQIHTGLQHSIIIRPTQPNCLPLDNATLPQKLKEVGYSTHMGVKGWHLG
FYRKECMPTRRGFDTFFGSLLGSGDYTHYKCDSPGMCGYDLYENDNAWDYDNGIYSTQMYTORVQQILASHNPTKPIFLYIAYQAVHS
PLQAPGRYFEHYRSIININRRRYAAMLSCLDEAINNVTLALKTYGFYNNSSIIYSSDNGGQPTAGGSNWPLRSGKGTYWEGGIRAVGFVH
SPLLKNKGTVCKELVHITDWYPTLISLAEGQIDEDIQLDGYDIWETISEGLRSPRVDILHNIDPIYTKAKNGSWAAGYGIWNTAIIQSAIR
VQHWKLLTGNPGYSDWVPPQSFNSLGNRWHNERITLSTGKSVLNFNITADPYERVVLSNRYPGIVKKLLRRLSQFNKTAVPVRYPPKDP
RSNPRNLNGGVWGPWYKEETKKKKPSKNQAEKKQKSKKKKKQKQKAVSGSTCHSGVTCG

FIGURE 15




FIGURE 16




FIGURE 17

Prosite Pattern Matches for 26212

Prosite version: Release 12.2 of February 1995

>PS00001|PDOC00001|ASN_GLYCOSYLATION N-glycosylation site.

Query: 157 NATL 160

Query: 306 NVTL 309

Query: 318 NNSI 321

Query: 431 NGSW 434

Query: 497 NITA 500

Query: 527 NKTA 530

>PS00004|PDOC00004|CAMP_PHOSPHO_SITE cAMP- and cGMP-dependent protein kinase phosphorylation site.

Query: 521 RRLS 524

Query: 562 KKPS 565

>PS00005|PDOC00005|PKC_PHOSPHO_SITE Protein kinase C phosphorylation site.

Query: 131 TGK 133

Query: 189 TRR 191

Query: 243 TQR 245

Query: 413 SPR 415

Query: 489 TGK 491

Query: 509 SNR 511

Query: 559 TTK 561

Query: 576 SKK 578

>PS00006|PDOC00006|CK2_PHOSPHO_SITE Casein kinase II phosphorylation site.

FIGURE 18A

Query: 298 SCLD 301
Query: 347 TYWE 350
Query: 386 SLAE 389
Query: 406 TISE 409

>PS00007|PDOC00007|TYR_PHOSPHO_SITE Tyrosine kinase phosphorylation site.

Query: 163 KLKEVG 169

>PS00008|PDOC00008|MYRISTYL N-myristoylation site.

Query: 28 GALAGF 33
Query: 56 GALLAQ 61
Query: 139 GLQHSI 144
Query: 198 GSLLGS 203
Query: 235 GIYSTQ 240
Query: 329 GGQPTA 334
Query: 343 GSKGTY 348
Query: 351 GGIRAV 356
Query: 432 GSWAAG 437
Query: 439 GIWNTA 444

>PS00149|PDOC00117|SULFATASE_2 Sulfatases signature 2.

Query: 168 GYSTMVGKW 177

>PS00523|PDOC00117|SULFATASE_1 Sulfatases signature 1.

Query: 120 PICTPSRSQFITG 132

FIGURE 18B

Alignments of top-scoring domains:

Sulfatase: domain 1 of 1, from 36 to 462: score 323.0, E = 3.5e-93
 *->PNIllilADDIGigdlGcyGaptirtpnidRlAeeGlrFtnayvttp
 PN+++ilADD+G+gdIG+ + t t n+D +A+eG+rF ++ +++
 PNFVillADUMGWGDLGANWAETKDTANLDKMASEGMRFPVDFHAAAS 82

25277 36 lCtPSRAallTGRyphrtGmytnnragvlpftgwsleGglpldettlpe
 C+PSRA+llTGR+ r+G++ n + +s +Gglpl+ettl+e+
 25277 83 TCSPSRASLLTGRGLRNGVTRNFAV-----TS-VGGLPLNETTLAEV 124

LkeaGYaTgmVgKWHlgynessasdfahlPlgrGFdyfygnlGSEdQWY
 L+++aGY+Tg++GKWllg++ ++ +P rGFdy++g
 25277 125 LQQAGYVTGIIGKWHLGTHGSY-----HPNFRGFdyYFG----- 158

plvdallpftndtytceggvgfskdvalkplgalgvneveapdkaladyk
 +p+++ ++ c +
 25277 159 -----IPYSH-DMGCT-----D 169

tagalnvpvhvFEWadryagavdvgrpflavliFprpaacflypnatvvs
 t+g+ + p + +++++ +r + ++ + a+ ly n+ +v+
 25277 170 TPGYNHPP-----CPACFPQGDGSPRNLQORDCY--TDVALPLYENLNIVE 211

qpmphspltPrpwqladealpflerngqrkpflylsykhvhiprda
 qD s l+ q +a++a +E++r+ + +pf+ly+++h+h+p
 25277 212 QPVNLSSLA-----QKYAEKATQFIQRASTSGRPFLLYVALAHMHPV--- 253

pmfsskdfagssrrglVglilDveemDdgvgvrlnaLdelNGlildnTl
 l+ + a x lYg + + emD +vg++ + +d + +nT+
 25277 254 --LEVTLQPAAPRGRSLYG---AGLWEMDSLVGQIKDKVDHT--VKENTF 296

liFTSllDhGghlgahghlgiragGsnppfrg.....gKgtlnlye
 FT D+G+ ++ + + Gs gpf g +++++++K+T+ +e
 25277 297 LWFTG--DNGFWAQKCELA-----GSVGFFTGfwqtrqggspAKQTT--WE 338

gGtRVPlivrwPeGiiapggvsdelvslmDlFPTildLAGap1Pgvaagv
 gG+RvP++++wP G+ + + +s +l s++D+FT++LA a+1P
 25277 339 GGHrvPALAYWP-GRVFNVTSTALLSVLDIFPTVVALAQASLP----- 381

kdrilDGvsLlplLlgaagssrhetlFyesyncnegrflpavrvgkklah
 + x DGv++ ++L g+ +++h lf++ n g a++ +
 25277 382 QGRREFDGVVSEVLGR-SQPGHrvLFHP---NSG-----AAGEFGALQT 422

frtpniagwqrdfddvklfntvedfnrsgddacrhdvckclgkprrs
 +r + + k+f++++ +++++ g+ + +
 25277 423 VRLE-----RYKAFYITGGAR---ACDGSTGFELQHKF----- 452

vthhdppllydlrDP<-*
 pl ++l+ D
 25277 453 -----PLIFNLEDDT 462

//

FIGURE 19

Alignments of top-scoring domains:

Sulfatase: domain 1 of 1, from 43 to 467; score 268.9, E = 6.5e-77

```

*->PNIIlilADDIgIdlGcyGnptirtptniDrLAeeGlrftnayvttt
PNI+L+L+DD++ ++LG+ ++ ++ + +G F na+vttt
23553 43 PNIIlLVLTDDQD-VELGSLQ---VMNKRTRKIMEHGGATFINAFVITP 85

IctPSRAallTGryphrtGmytnnragvlpftgwsleGglpldettlpeI
+C+PSR++ LTG+y h+++ytann++ ++++ w+ ++ +t++++
23553 86 MCCPSRSSHLTGKYVHNHNVYTNEN--CSSPSWQ----AMHEPRTFAVY 129

LkeaGYaTgmVGKWHlgymeessasdfahIPlerG.FdyfygnlGGEdQW
L + GV+T+++GK++++yn ++ +P+g+ ++ +n
23553 130 LWNTRYRTAFFGKYLNEYNGSY-----IPPGWRWGLLIKN----- 165

Yplvdallpftndtytceggysgfskdvalkplgalgvneveapdkalady
++E+n + c++g + +++++ +++++ dy
23553 166 -----SRFYN-YTVCRNG-----IKEKHGFDYAK-----DY 190

ktagalnvphhvfEWadryagavdvgrpflavli fprpaacflypnatvv
+t++++n + y++++ p+++++ +
23553 191 FTDLITNES-----INyFKMSK-----RHYPHRPVMMV----I 219

sqcmphspltaprpwqladealpflerngqrdkpflylysykhvhiprd
s+ +ph p + + +++++ + p+ + + + + + +kh+ +++++
23553 220 SHAAFHGPEd-S---APQFSKLYPNASQH-ITPSYNYAPNMKGHWIMQYT 264

apmlfsskdfagssrrglyglilDsveemDdgvgvrvlnaLdelNGlIdnT
+pml+ + +f+ ++r++ + +++++Dd+v+r++n L e G+l+nT
23553 265 GPMLPIHMEFTNlQRKRLQ----TLMSVDDSVRLYNHLVET-GELENT 309

liiFTSllDhGghlgahghlqiragGsnppfrggKgtlnlyegGtRvPliv
+ii+T+ DhG+h+g++g+ + gK+++ y++++RvP+++
23553 310 YIIYTA--DHGYHIGQFGLV-----K-GKSHP-YDFDIRVPFFI 344

rwPeGiIapggvsdelvslmDlfpTildLAGapIpgvaagvkdriLDGvs
z+P +pg+++ ++v ++Dl+PTild+AG++ P +DG+s
23553 345 RGP--SVEPGSIVPQIVLNLIDLAPTILOIAGLDTP-----PDVDGKS 384

Llplllgaagssrbetlfyesyncnegrflpavrwgkckahfctpnagw
+l+LL+ + ++ +f + + + + + + + +f
23553 385 VLKLLDPE---KPGNRFRt-NKKAK---INRDTFLVERGKF----- 418

qrvidddvwlfnfntvedfnrsgddacrhdvckclgkprsvthhdppll
+ k + + + +s++ + + + +C +++++ ++ ++p +
23553 419 -----LRKKEESSKNIQQSNHLPKYERVKELCQQARYQTA-CEQFGQK 460

ydlrDP<+
+D
23553 461 WQCIEDT 467

```

FIGURE 20

Gene 23553 Xenograph Panel
Relative Expression (MDA435 as Reference)




FIGURE 21

23553 Expression in Breast Samples




FIGURE 22




FIGURE 23

23553 Expression in Lung




FIGURE 24

TAQMAN RESULTS

23553 expression in normal human tissues




FIGURE 25

Gene 23553 oncology Panel
Relative Expression (Normal Breast as Reference)




FIGURE 26

Alignments of top-scoring domains:

Sulfatase: domain 1 of 1, from 47 to 471: score 289.7, E = 3.6e-83
 *->PNIllilaDDlGigdlGcyGnptirtptniDrLAeeGlrFtnayvttt
 P+i++il+DD+G+ d+G +G + i+tp++DrLA+ G+++ n y+ +p
 25278 47 PHIIIFILTDDQGYHGVYHG-SDIETPTLDRLAAKGVKLEN-YYIQP 91

ICtPSRAaLLTGryphrtGmytnnragvlpftgwsleGglpldettlpe1
 +CtPSR++lLTGRy+++tG+++ + p+++ +lpld +tlp+
 25278 92 ICTPSRSQLLTGryQIHTGLQHSIIR---PQQPN---CLPLDQVTLQPK 134

LkecGYaTgmvgKWHlgynessasdfahlPlgrGfdyfygnlGGEdQWY
 L+eaGY T+mvGKWHlg +++++ lP++rGFd+E+g+
 25278 135 LQEAGYSTIMVGKWHLGfYRKEC-----LPTRRGFDTLGSG----- 170

plvdallpftndtytceggysfskdvalkplgalgvneveapdkaladyk
 l + d+yt+++ ++
 25278 171 -----LTGNVDYTYDN-----CD 184

tagalnvpvhvfeWadryagavdvgrpflavli fprpaacflypnatvvs
 +g+ + +d + + + +
 25278 185 GPGVCG-----FD-----LHEGENVAVWG 202

qmpshspltaprpwqladealpflerngqrkpf flylsykhvhiprd.
 ++s++ +a++a l ++ +p fly+++++vh+p ++
 25278 203 LSGQYSTML-----YAQRASHILASH-SFQRPLFLYVAFQAVHTPLQS 244

apmlfsskdfagssrrqlYglldSveemDdgvgvrvlnalDdelNGlldnt
 + +++ ++ g+ r+ Y+ ++v mD++v ++ aL++ G ++n
 25278 245 PREYLYRYRTMGVARRKYA---AMVTCMDEAVRNITWALKRY-GFYNN 290

liiFTSllDhGghlgahghlgiragGsnppfrrgGktnlyegGtrvPliv
 +iif+S D+Gg++ gGsn+p+rg+Kgt +cgG+R ++v
 25278 291 VIIFSS--DNGGQTF-----S-----GGSNWPLRGRKGTY-WEGGVRLGFEV 329

rwPeGiiapggvdelvslmDlfpTiildLAGaplpGvaagvkdriLDGvs
 ++P +++ ++s++l ++ D++PT++ LAG++ + lDG++
 25278 330 HSP-LLKRRQRTSRALMHITDWPYTLVGLAGGTTS-----AADGLDGYD 372

Llplllgaagssrhetl fye.....sycnegrgflpavrvqkkahfrt
 ++p++ ++ +s+r e+l++++ ++ ++ g + g ++ + +
 25278 373 VWPAISEGRASPRTEILHHITdplynhaQHGSLEG-----GFCIWNFAVQA 417

pni.agwqrvdfddvwlkifntvedfnrsgddacrghdvckclgkprrevt
 + + w + + ++d+ ++ a + g + + ++
 25278 418 AIRvGEWK-----LLTGDPGYGDWIPQTLATFPGSWNLER-----MAS 457

hhdppllydlstrDP<-*
 + l+++s+DP
 25278 458 VRQAVWLFNISADP 471

FIGURE 27

			<u>Relative Expression</u>
NDR 19	Breast T	DCIS	47.84
MDA 138	Breast N	Normal	52.89
NDR 01	Breast T	IDC	44.79
NDR 15	Breast T	DC	29.55
NDR 133	Breast T	ILC	43.26
MDA 161	Breast T	IDC	60.13
MDA 155	Breast T	IDC/DCIS	20.11
PIT 270	Lung N	Normal	36.00
CHT 427	Lung N	Normal	26.54
PIT 241	Lung N	Normal	31.45
PIT 298	Lung N	Normal	17.57
CHT 800	Lung T	AC	31.45
CHT 335	Lung T	SCC	35.02
CHT 447	Lung T	AC	27.19
CHT 752	Lung T	AC	3.89
CHT 799	Lung T	AC	5.74
CHT 813	Lung T	SCC	47.18
CHT 369	Lung T	SCC	42.37
CHT 371	Colon N	Normal	2.37
CHT 396	Colon N	Normal	16.34
CHT 398	Colon N	Normal	15.24
NDR 104	Colon N	Normal	20.89
CHT 520	Colon T	Adeno	11.71
CHT 122	Colon T	Adeno	360.79
CHT 536	Colon T	Adeno	1.00

FIGURE 28A

			<u>Relative Expression</u>
CHT 528	Colon T	Adeno	11.63
CHT 386	Colon T	Adeno	372.22
CHT 372	Colon T	Adeno	2.39
CHT 532	Colon T	Adeno	4.45
CHT 77	Liver Met	Met	23.43
CHT 321	Liver Met	Met	11.35
CHT 84	Liver Met	Met	30.38
NDR 100	Liver Met	Met	46.21
NDR 154	Liver N	Normal	7.31
CHT 322	Liver N	Normal	9.38
PIT 51	Liver N	Normal	1.77
CHT 339	Liver	Normal	1.58
PIT 265	Breast N	Normal	37.40
MDA 335	Breast N	Normal	45.57
NDR 132	Breast T	DCIS	19.56
NDR 13	Breast N	Normal	6.73
NDR 56	Breast N	Normal	20.61

FIGURE 28B

ments of top-scoring domains:

Case: domain 1 of 1, from 76 to 502: score 324.5, E = 1.3e-93

```

*->PNvllilADDlGigdlgcyghptirTPnlDrLAeeGlrFtnhytatp
P+ ++ilaDD+G+ d+g++g ++i TP+ld+LA+eG+++ n+y+ +p
26212 76 PHILIFILADDQGFrdVGYHG-SEIKTPTLDKLAEGVKLENYYV-QP 120

lCsPSRAaLlTGryphrhGmvsngrlgvlgtaksgglpldettLpelLk
+C+PSR+++ TG+y++++G + + + ++ +lpld +tLp+ Lk
26212 121 ICTPSRSQFITGKYQIHTGLQH-----SIIRPTQPNCLPLDNATLPQKLK 165

eaGYaTglvGKWHlglnensdaagdgehlPlgwrGfdyfdgflygspfty
e GY T++vGKWHlg+++ +e+ P++ rGfd f+g l+gs ++y
26212 166 EVGYSTHMGKWHLGFYR-----KECMPTR-RGFDTFFGSLLGSGDY 207

deencdngegteppeaypeqgwlpqilgyyltdlladkalglldvasaag
++ cd +p+ +++++l+ aa
26212 208 THYKCD-----SPGM-----CGYDLYENDNAA- 229

rllakalaasrPFflyisppaphfsilfrnfkevaqpyrapqltqlfvde
++++ + ++tq++++
26212 230 -----WDYD-----NGIYSTQMYTOR 245

aadfiernk.ekPfflylafllrhvhtplfspaedleskdflgrsqrgrY
++++++ kP fly a++ +vh pl++p + e+++ r+rY
26212 246 VQQILASHNpTKPIFLYIAYQ--AVHSPLQAPGRYFEHYRSIININRRRY 293

gdlveemDdlvGrvldaLedlGllDNTlviftSDnGahlegtpewygggn
+++++ D+++++v aL+ G ++N ++i++SDnG g+p+ +gg+n
26212 294 AANLSCLEAINNVTALALKTYGFYNNSIIYYSSDNG---GQPT-AGGSN 338

gplkggKgygslyeGgiRvPllvrvPggiapagrvekseivshvDlaPT
+pl+g Kg+ +eGgiR ++v++P + +g+v + elv++ D++PT
26212 339 WPLRGSKGTY--WEGGIRAVGVFVHSP-LLKNKGTVCCK--ELVHITDWYPT 383

ildlAGaplpkvanGakdrplDGvsllplllggaapsrrahetlfhyngk
+ +lA + ++ d lDG++++ + +g + s+ + +++h+
26212 384 LISLAEGQIDE-----DIQLDGYDIWBTISEGLR-SP--RVDILHN--- 421

grklravrwprksgktpklkahfftpaf.....
++ ++ +k+ + + a + ++ ++ + ++ + +++++ ++
26212 422 ---IDPIYTKAKN---GSWAAGYGIWNTaiqsairvqhkwlltgnpgysd 465

....dddtngwecvgtvsqaddiedcrcegvethhdppelyDlsrDP
++++ n+g + ++ e t+ + +l++ ++DP
26212 466 wvppQSFSNLG-----PNRWHNER-ITLSTGKSVWLFNITADP 502

<-*
26212 - -

```

FIGURE 29




FIGURE 30

26212.1 Expression in Oncology Plate I




FIGURE 31A

26212.1 Expression in Clinical Breast Samples




FIGURE 32




FIGURE 33

26212.1 Expression in MCF 10A & 3B EGF Treated Cells




FIGURE 34

26212.2 Expression in the Angiogenesis Panel




FIGURE 35




FIGURE 36

SEQUENCE LISTING

<110> Glucksman, Maria Alexandra
Williamson, Mark
Tsia, Fong-Ying
Rudolph-Owen, Laura A.

<120> 22438, 23553, 25278, and 26212 Novel
Human Sulfatases (A CIP Application)

<130> 35800/208709

<151>

<160> 14

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 525

<212> PRT

<213> homo sapiens .

<400> 1

```

Met Gly Trp Leu Phe Leu Lys Val Leu Leu Ala Gly Val Ser Phe Ser
 1           5           10           15
Gly Phe Leu Tyr Pro Leu Val Asp Phe Cys Ile Ser Gly Lys Thr Arg
          20           25           30
Gly Gln Lys Pro Asn Phe Val Ile Leu Ala Asp Asp Met Gly Trp
          35           40           45
Gly Asp Leu Gly Ala Asn Trp Ala Glu Thr Lys Asp Thr Ala Asn Leu
          50           55           60
Asp Lys Met Ala Ser Glu Gly Met Arg Phe Val Asp Phe His Ala Ala
          65           70           75           80
Ala Ser Thr Cys Ser Pro Ser Arg Ala Ser Leu Leu Thr Gly Arg Leu
          85           90           95
Gly Leu Arg Asn Gly Val Thr Arg Asn Phe Ala Val Thr Ser Val Gly
          100          105          110
Gly Leu Pro Leu Asn Glu Thr Thr Leu Ala Glu Val Leu Gln Gln Ala
          115          120          125
Gly Tyr Val Thr Gly Ile Ile Gly Lys Trp His Leu Gly His His Gly
          130          135          140
Ser Tyr His Pro Asn Phe Arg Gly Phe Asp Tyr Tyr Phe Gly Ile Pro
          145          150          155          160
Tyr Ser His Asp Met Gly Cys Thr Asp Thr Pro Gly Tyr Asn His Pro
          165          170          175
Pro Cys Pro Ala Cys Pro Gln Gly Asp Gly Pro Ser Arg Asn Leu Gln
          180          185          190
Arg Asp Cys Tyr Thr Asp Val Ala Leu Pro Leu Tyr Glu Asn Leu Asn
          195          200          205
Ile Val Glu Gln Pro Val Asn Leu Ser Ser Leu Ala Gln Lys Tyr Ala
          210          215          220
Glu Lys Ala Thr Gln Phe Ile Gln Arg Ala Ser Thr Ser Gly Arg Pro
          225          230          235          240
Phe Leu Leu Tyr Val Ala Leu Ala His Met His Val Pro Leu Pro Val
          245          250          255
Thr Gln Leu Pro Ala Ala Pro Arg Gly Arg Ser Leu Tyr Gly Ala Gly
          260          265          270
Leu Trp Glu Met Asp Ser Leu Val Gly Gln Ile Lys Asp Lys Val Asp
          275          280          285
His Thr Val Lys Glu Asn Thr Phe Leu Trp Phe Thr Gly Asp Asn Gly
          290          295          300
Pro Trp Ala Gln Lys Cys Glu Leu Ala Gly Ser Val Gly Pro Phe Thr
          305          310          315          320
Gly Phe Trp Gln Thr Arg Gln Gly Gly Ser Pro Ala Lys Gln Thr Thr

```

```

          325          330          335
Trp Glu Gly Gly His Arg Val Pro Ala Leu Ala Tyr Trp Pro Gly Arg
          340          345          350
Val Pro Val Asn Val Thr Ser Thr Ala Leu Leu Ser Val Leu Asp Ile
          355          360          365
Phe Pro Thr Val Val Ala Leu Ala Gln Ala Ser Leu Pro Gln Gly Arg
          370          375          380
Arg Phe Asp Gly Val Asp Val Ser Glu Val Leu Phe Gly Arg Ser Gln
          385          390          395          400
Pro Gly His Arg Val Leu Phe His Pro Asn Ser Gly Ala Ala Gly Glu
          405          410          415
Phe Gly Ala Leu Gln Thr Val Arg Leu Glu Arg Tyr Lys Ala Phe Tyr
          420          425          430
Ile Thr Gly Gly Ala Arg Ala Cys Asp Gly Ser Thr Gly Pro Glu Leu
          435          440          445
Gln His Lys Phe Pro Leu Ile Phe Asn Leu Glu Asp Asp Thr Ala Glu
          450          455          460
Ala Val Pro Leu Glu Arg Gly Gly Ala Glu Tyr Gln Ala Val Leu Pro
          465          470          475          480
Glu Val Arg Lys Val Leu Ala Asp Val Leu Gln Asp Ile Ala Asn Asp
          485          490          495
Asn Ile Ser Ser Ala Asp Tyr Thr Gln Asp Pro Ser Val Thr Pro Cys
          500          505          510
Cys Asn Pro Tyr Gln Ile Ala Cys Arg Cys Gln Ala Ala
          515          520          525

```

<210> 2
 <211> 2175
 <212> DNA
 <213> homo sapiens

<220>
 <221> CDS
 <222> (248)...(1825)

```

<400> 2
cacgcgtccg caaatttcct gattcttttg aattaggatt ccagatgggg gcctcatttc      60
tacagccccc aacattccta tagccgttat cactgccatc accactgcca ccagcatctt      120
cttgacagatt ccacccctgc tcccagaga cttcctgctt tgaaagtgag cagaaaggaa      180
gctctcagaa aaatctctag tgggtgctgc cgctcgtcca gacaatcgga atcctgcctt      240
caccacc atg ggc tgg ctt ttt cta aag gtt ttg ttg gcg gga gtg agt      289
      Met Gly Trp Leu Phe Leu Lys Val Leu Leu Ala Gly Val Ser
              1              5              10

ttc tca gga ttt ctt tat cct ctt gtg gat ttt tgc atc agt ggg aaa      337
Phe Ser Gly Phe Leu Tyr Pro Leu Val Asp Phe Cys Ile Ser Gly Lys
      15              20              25              30

aca aga gga cag aag cca aac ttt gtg att att ttg gcc gat gac atg      385
Thr Arg Gly Gln Lys Pro Asn Phe Val Ile Leu Ala Asp Asp Met
              35              40              45

ggg tgg ggt gac ctg gga gca aac tgg gca gaa aca aag gac act gcc      433
Gly Trp Gly Asp Leu Gly Ala Asn Trp Ala Glu Thr Lys Asp Thr Ala
              50              55              60

aac ctt gat aag atg gct tcg gag gga atg agg ttt gtg gat ttc cat      481
Asn Leu Asp Lys Met Ala Ser Glu Gly Met Arg Phe Val Asp Phe His
              65              70              75

gca gct gcc tcc acc tgc tca ccc tcc cgg gct tcc ttg ctc acc ggc      529
Ala Ala Ala Ser Thr Cys Ser Pro Ser Arg Ala Ser Leu Leu Thr Gly
              80              85              90

cgg ctt ggc ctt cgc aat gga gtc aca cgc aac ttt gca gtc act tct      577
Arg Leu Gly Leu Arg Asn Gly Val Thr Arg Asn Phe Ala Val Thr Ser
      95              100              105              110

```

gtg gga ggc ctt ccg ctc aac gag acc acc ttg gca gag gtg ctg cag Val Gly Gly Leu Pro Leu Asn Glu Thr Thr Leu Ala Glu Val Leu Gln 115 120 125	625
cag gcg ggt tac gtc act ggg ata ata ggc aaa tgg cat ctt gga cac Gln Ala Gly Tyr Val Thr Gly Ile Ile Gly Lys Trp His Leu Gly His 130 135 140	673
cac ggc tct tat cac ccc aac ttc cgt ggt ttt gat tac tac ttt gga His Gly Ser Tyr His Pro Asn Phe Arg Gly Phe Asp Tyr Tyr Phe Gly 145 150 155	721
atc cca tat agc cat gat atg ggc tgt act gat act cca ggc tac aac Ile Pro Tyr Ser His Asp Met Gly Cys Thr Asp Thr Pro Gly Tyr Asn 160 165 170	769
cac cct cct tgt cca gcg tgt cca cag ggt gat gga cca tca agg aac His Pro Pro Cys Pro Ala Cys Pro Gln Gly Asp Gly Pro Ser Arg Asn 175 180 185 190	817
ctt caa aga gac tgt tac act gac gtg gcc ctc cct ctt tat gaa aac Leu Gln Arg Asp Cys Tyr Thr Asp Val Ala Leu Pro Leu Tyr Glu Asn 195 200 205	865
ctc aac att gtg gag cag ccg gtg aac ttg agc agc ctt gcc cag aag Leu Asn Ile Val Glu Gln Pro Val Asn Leu Ser Ser Leu Ala Gln Lys 210 215 220	913
tat gct gag aaa gca acc cag ttc atc cag cgt gca agc acc agc ggg Tyr Ala Glu Lys Ala Thr Gln Phe Ile Gln Arg Ala Ser Thr Ser Gly 225 230 235	961
agg ccc ttc ctg ctc tat gtg gct ctg gcc cac atg cac gtg ccc tta Arg Pro Phe Leu Leu Tyr Val Ala Leu Ala His Met His Val Pro Leu 240 245 250	1009
ccc gtg act cag cta cca gca gcg cca cgg ggc aga agc ctg tat ggt Pro Val Thr Gln Leu Pro Ala Ala Pro Arg Gly Arg Ser Leu Tyr Gly 255 260 265 270	1057
gca ggg ctc tgg gag atg gac agt ctg gtg ggc cag atc aag gac aaa Ala Gly Leu Trp Glu Met Asp Ser Leu Val Gly Gln Ile Lys Asp Lys 275 280 285	1105
gtt gac cac aca gtg aag gaa aac aca ttc ctc tgg ttt aca gga gac Val Asp His Thr Val Lys Glu Asn Thr Phe Leu Trp Phe Thr Gly Asp 290 295 300	1153
aat ggc ccg tgg gct cag aag tgt gag cta gcg ggc agt gtg ggt ccc Asn Gly Pro Trp Ala Gln Lys Cys Glu Leu Ala Gly Ser Val Gly Pro 305 310 315	1201
ttc act gga ttt tgg caa act cgt caa ggg gga agt cca gcc aag cag Phe Thr Gly Phe Trp Gln Thr Arg Gln Gly Gly Ser Pro Ala Lys Gln 320 325 330	1249
acg acc tgg gaa gga ggg cac cgg gtc cca gca ctg gct tac tgg cct Thr Thr Trp Glu Gly Gly His Arg Val Pro Ala Leu Ala Tyr Trp Pro 335 340 345 350	1297
ggc aga gtt cca gtt aat gtc acc agc act gcc ttg tta agc gtg ctg Gly Arg Val Pro Val Asn Val Thr Ser Thr Ala Leu Leu Ser Val Leu 355 360 365	1345
gac att ttt cca act gtg gta gcc ctg gcc cag gcc agc tta cct caa Asp Ile Phe Pro Thr Val Val Ala Leu Ala Gln Ala Ser Leu Pro Gln 370 375 380 385 390	1393

```

370          375          380
gga cgg cgc ttt gat ggt gtg gac gtc tcc gag gtg ctc ttt ggc cgg      1441
Gly Arg Arg Phe Asp Gly Val Asp Val Ser Glu Val Leu Phe Gly Arg
385          390          395

tca cag cct ggg cac agg gtg ctg ttc cac ccc aac agc ggg gca gct      1489
Ser Gln Pro Gly His Arg Val Leu Phe His Pro Asn Ser Gly Ala Ala
400          405          410

gga gag ttt gga gcc ctg cag act gtc cgc ctg gag cgt tac aag gcc      1537
Gly Glu Phe Gly Ala Leu Gln Thr Val Arg Leu Glu Arg Tyr Lys Ala
415          420          425

ttc tac att acc ggt gga gcc agg gcg tgt gat ggg agc acg ggg cct      1585
Phe Tyr Ile Thr Gly Gly Ala Arg Ala Cys Asp Gly Ser Thr Gly Pro
435          440          445

gag ctg cag cat aag ttt cct ctg att ttc aac ctg gaa gac gat acc      1633
Glu Leu Gln His Lys Phe Pro Leu Ile Phe Asn Leu Glu Asp Thr
450          455          460

gca gaa gct gtg ccc cta gaa aga ggt ggt gcg gag tac cag gct gtg      1681
Ala Glu Ala Val Pro Leu Glu Arg Gly Gly Ala Glu Tyr Gln Ala Val
465          470          475

ctg ccc gag gtc aga aag gtt ctt gca gac gtc ctc caa gac att gcc      1729
Leu Pro Glu Val Arg Lys Val Leu Ala Asp Val Leu Gln Asp Ile Ala
480          485          490

aac gac aac atc tcc agc gca gat tac act cag gac cct tca gta act      1777
Asn Asp Asn Ile Ser Ser Ala Asp Tyr Thr Gln Asp Pro Ser Val Thr
495          500          505

ccc tgc tgt aat ccc tac caa att gcc tgc cgc tgt caa gcc gca taa      1825
Pro Cys Cys Asn Pro Tyr Gln Ile Ala Cys Arg Cys Gln Ala Ala *
515          520          525

cagaccaatt tttattccac gaggaggagt acctggaaat taggcaagtt tgcttccaaa      1885
tttcattttt accctcttta caaacacacg ctttagttta gtcttgagg ttagttttgg      1945
agttagcctt gcatatccct tctgtatcct gtccctcctc cacgccgacc cgagagcagc      2005
tgagctgcgc tggctctggg cagggagtgt gccttaatgg gaagcacacg ggctttggag      2065
tcaggcacag gtgccagctc cagcttttga acttgggcaa ttgtttaacc taacctgcaa      2125
gttgattttg agggttaaat aaaggcatatc atgaaaaaaaa aaaaaaaaaa      2175

<210> 3
<211> 871
<212> PRT
<213> homo sapiens

<400> 3
Met Lys Tyr Ser Cys Cys Ala Leu Val Leu Ala Val Leu Gly Thr Glu
1          5          10          15
Leu Leu Gly Ser Leu Cys Ser Thr Val Arg Ser Pro Arg Phe Arg Gly
20          25          30
Arg Ile Gln Gln Glu Arg Lys Asn Ile Arg Pro Asn Ile Ile Leu Val
35          40          45
Leu Thr Asp Asp Gln Asp Val Glu Leu Gly Ser Leu Gln Val Met Asn
50          55          60
Lys Thr Arg Lys Ile Met Glu His Gly Gly Ala Thr Phe Ile Asn Ala
65          70          75          80
Phe Val Thr Thr Pro Met Cys Cys Pro Ser Arg Ser Ser Met Leu Thr
85          90          95
Gly Lys Tyr Val His Asn His Asn Val Tyr Thr Asn Asn Glu Asn Cys
100          105          110
Ser Ser Pro Ser Trp Gln Ala Met His Glu Pro Arg Thr Phe Ala Val
115          120          125

```

Tyr Leu Asn Asn Thr Gly Tyr Arg Thr Ala Phe Phe Gly Lys Tyr Leu
 130 135 140
 Asn Glu Tyr Asn Gly Ser Tyr Ile Pro Pro Gly Trp Arg Glu Trp Leu
 145 150 155 160
 Gly Leu Ile Lys Asn Ser Arg Phe Tyr Asn Tyr Thr Val Cys Arg Asn
 165 170 175
 Gly Ile Lys Glu Lys His Gly Phe Asp Tyr Ala Lys Asp Tyr Phe Thr
 180 185 190
 Asp Leu Ile Thr Asn Glu Ser Ile Asn Tyr Phe Lys Met Ser Lys Arg
 195 200 205
 Met Tyr Pro His Arg Pro Val Met Met Val Ile Ser His Ala Ala Pro
 210 215 220
 His Gly Pro Glu Asp Ser Ala Pro Gln Phe Ser Lys Leu Tyr Pro Asn
 225 230 235 240
 Ala Ser Gln His Ile Thr Pro Ser Tyr Asn Tyr Ala Pro Asn Met Asp
 245 250 255
 Lys His Trp Ile Met Gln Tyr Thr Gly Pro Met Leu Pro Ile His Met
 260 265 270
 Glu Phe Thr Asn Ile Leu Gln Arg Lys Arg Leu Gln Thr Leu Met Ser
 275 280 285
 Val Asp Asp Ser Val Glu Arg Leu Tyr Asn Met Leu Val Glu Thr Gly
 290 295 300
 Glu Leu Glu Asn Thr Tyr Ile Ile Tyr Thr Ala Asp His Gly Tyr His
 305 310 315 320
 Ile Gly Gln Phe Gly Leu Val Lys Gly Lys Ser Met Pro Tyr Asp Phe
 325 330 335
 Asp Ile Arg Val Pro Phe Phe Ile Arg Gly Pro Ser Val Glu Pro Gly
 340 345 350
 Ser Ile Val Pro Gln Ile Val Leu Asn Ile Asp Leu Ala Pro Thr Ile
 355 360 365
 Leu Asp Ile Ala Gly Leu Asp Thr Pro Pro Asp Val Asp Gly Lys Ser
 370 375 380
 Val Leu Lys Leu Leu Asp Pro Glu Lys Pro Gly Asn Arg Phe Arg Thr
 385 390 395 400
 Asn Lys Lys Ala Lys Ile Trp Arg Asp Thr Phe Leu Val Glu Arg Gly
 405 410 415
 Lys Phe Leu Arg Lys Lys Glu Glu Ser Ser Lys Asn Ile Gln Gln Ser
 420 425 430
 Asn His Leu Pro Lys Tyr Glu Arg Val Lys Glu Leu Cys Gln Gln Ala
 435 440 445
 Arg Tyr Gln Thr Ala Cys Glu Gln Pro Gly Gln Lys Trp Gln Cys Ile
 450 455 460
 Glu Asp Thr Ser Gly Lys Leu Arg Ile His Lys Cys Lys Gly Pro Ser
 465 470 475 480
 Asp Leu Leu Thr Val Arg Gln Ser Thr Arg Asn Leu Tyr Ala Arg Gly
 485 490 495
 Phe His Asp Lys Asp Lys Glu Cys Ser Cys Arg Glu Ser Gly Tyr Arg
 500 505 510
 Ala Ser Arg Ser Gln Arg Lys Ser Gln Arg Gln Phe Leu Arg Asn Gln
 515 520 525
 Gly Thr Pro Lys Tyr Lys Pro Arg Phe Val His Thr Arg Gln Thr Arg
 530 535 540
 Ser Leu Ser Val Glu Phe Glu Gly Glu Ile Tyr Asp Ile Asn Leu Glu
 545 550 555 560
 Glu Glu Glu Glu Leu Gln Val Leu Gln Pro Arg Asn Ile Ala Lys Arg
 565 570 575
 His Asp Glu Gly His Lys Gly Pro Arg Asp Leu Gln Ala Ser Ser Gly
 580 585 590
 Gly Asn Arg Gly Arg Met Leu Ala Asp Ser Ser Asn Ala Val Gly Pro
 595 600 605
 Pro Thr Thr Val Arg Val Thr His Lys Cys Phe Ile Leu Pro Asn Asp
 610 615 620
 Ser Ile His Cys Glu Arg Glu Leu Tyr Gln Ser Ala Arg Ala Trp Lys
 625 630 635 640
 Asp His Lys Ala Tyr Ile Asp Lys Glu Ile Glu Ala Leu Gln Asp Lys
 645 650 655
 Ile Lys Asn Leu Arg Glu Val Arg Gly His Leu Lys Arg Arg Lys Pro

660 665 670
 Glu Glu Cys Ser Cys Ser Lys Gln Ser Tyr Tyr Asn Lys Glu Lys Gly
 675 680 685
 Val Lys Lys Gln Glu Lys Leu Lys Ser His Leu His Pro Phe Lys Glu
 690 695 700
 Ala Ala Gln Glu Val Asp Ser Lys Leu Gln Leu Phe Lys Glu Asn Asn
 705 710 715 720
 Arg Arg Arg Lys Lys Glu Arg Lys Glu Lys Arg Arg Gln Arg Lys Gly
 725 730 735
 Glu Glu Cys Ser Leu Pro Gly Leu Thr Cys Phe Thr His Asp Asn Asn
 740 745 750
 His Trp Gln Thr Ala Pro Phe Trp Asn Leu Gly Ser Phe Cys Ala Cys
 755 760 765
 Thr Ser Ser Asn Asn Asn Thr Tyr Trp Cys Leu Arg Thr Val Asn Glu
 770 775 780
 Thr His Asn Phe Leu Phe Cys Glu Phe Ala Thr Gly Phe Leu Glu Tyr
 785 790 795 800
 Phe Asp Met Asn Thr Asp Pro Tyr Gln Leu Thr Asn Thr Val His Thr
 805 810 815
 Val Glu Arg Gly Ile Leu Asn Gln Leu His Val Gln Leu Met Glu Leu
 820 825 830
 Arg Ser Cys Gln Gly Tyr Lys Gln Cys Asn Pro Arg Pro Lys Asn Leu
 835 840 845
 Asp Val Gly Asn Lys Asp Gly Gly Ser Tyr Asp Leu His Arg Gly Gln
 850 855 860
 Leu Trp Asp Gly Trp Glu Gly
 865 870

<210> 4
 <211> 4321
 <212> DNA
 <213> homo sapiens

<220>
 <221> CDS
 <222> (510)... (3125)

<400> 4
 cccacgcgtc cggctaataga atcttggggc cgggtgtcggg ccggggcggc ttgatcggca 60
 actaggaaac cccaggcgca gaggccagga gcgagggcag cgaggatcag aggccaggcc 120
 ttcccggctg ccggcgctcc tcggaggtca gggcagatga ggaacatgac tctcccctt 180
 cggaggagga aggaagtccc gctgccacct tatctctgct cctctgcctc ctccctgttc 240
 ccagagcttt ttctctagag aagattttga agggcgcttt tgtgctgacg gccaccacc 300
 atcatctaaa gaagataaac ttggcaaatg acatgcaggt tcttcaaggc agaataattg 360
 cagaaaatct tcaaaggacc ctatctgcag atgttctgaa tacctctgag aatagagatt 420
 gattattcaa ccaggatacc taattcaaga actccagaaa tcaggagacg gagacatttt 480
 gtcagttttg caacattgga ccaaataca atg aag tat tct tgc tgt gct ctg 533
 Met Lys Tyr Ser Cys Cys Ala Leu
 1 5
 gtt ttg gct gtc ctg ggc aca gaa ttg ctg gga agc ctc tgt tgc act 581
 Val Leu Ala Val Leu Gly Thr Glu Leu Leu Gly Ser Leu Cys Ser Thr
 10 15 20
 gtc aga tcc ccg agg ttc aga gga cgg ata cag cag gaa cga aaa aac 629
 Val Arg Ser Pro Arg Phe Arg Gly Arg Ile Gln Gln Glu Arg Lys Asn
 25 30 35 40
 atc cga ccc aac att att ctt gtg ctt acc gat gat caa gat gtg gag 677
 Ile Arg Pro Asn Ile Ile Leu Val Leu Thr Asp Asp Gln Asp Val Glu
 45 50 55
 ctg ggg tcc ctg caa gtc atg aac aaa acg aga aag att atg gaa cat 725
 Leu Gly Ser Leu Gln Val Met Asn Lys Thr Arg Lys Ile Met Glu His
 60 65 70
 ggg ggg gcc acc ttc atc aat gcc ttt gtg act aca ccc atg tgc tgc 773
 6

Gly	Gly	Ala	Thr	Phe	Ile	Asn	Ala	Phe	Val	Thr	Thr	Pro	Met	Cys	Cys		
		75					80					85					
ccg	tca	cgg	tcc	tcc	atg	ctc	acc	ggg	aag	tat	gtg	cac	aat	cac	aat	821	
Pro	Ser	Arg	Ser	Ser	Met	Leu	Thr	Gly	Lys	Tyr	Val	His	Asn	His	Asn		
	90					95				100							
gtc	tac	acc	aac	aac	gag	aac	tgc	tct	tcc	ccc	tcg	tgg	cag	gcc	atg	869	
Val	Tyr	Thr	Asn	Asn	Glu	Asn	Cys	Ser	Ser	Pro	Ser	Trp	Gln	Ala	Met		
105					110					115					120		
cat	gag	cct	cgg	act	ttt	gct	gta	tat	ctt	aac	aac	act	ggc	tac	aga	917	
His	Glu	Pro	Arg	Thr	Phe	Ala	Val	Tyr	Leu	Asn	Asn	Thr	Gly	Tyr	Arg		
				125					130					135			
aca	gcc	ttt	ttt	gga	aaa	tac	ctc	aat	gaa	tat	aat	ggc	agc	tac	atc	965	
Thr	Ala	Phe	Phe	Gly	Lys	Tyr	Leu	Asn	Glu	Tyr	Asn	Gly	Ser	Tyr	Ile		
			140					145					150				
ccc	cct	ggg	tgg	cga	gaa	tgg	ctt	gga	tta	atc	aag	aat	tct	cgc	ttc	1013	
Pro	Pro	Gly	Trp	Arg	Glu	Trp	Leu	Gly	Leu	Ile	Lys	Asn	Ser	Arg	Phe		
		155					160					165					
tat	aat	tac	act	gtt	tgt	cgc	aat	ggc	atc	aaa	gaa	aag	cat	gga	ttt	1061	
Tyr	Asn	Tyr	Thr	Val	Cys	Arg	Asn	Gly	Ile	Lys	Glu	Lys	His	Gly	Phe		
	170					175					180						
gat	tat	gca	aag	gac	tac	ttc	aca	gac	tta	atc	act	aac	gag	agc	att	1109	
Asp	Tyr	Ala	Lys	Asp	Tyr	Phe	Thr	Asp	Leu	Ile	Thr	Asn	Glu	Ser	Ile		
185					190					195					200		
aat	tac	ttc	aaa	atg	tct	aag	aga	atg	tat	ccc	cat	agg	ccc	gtt	atg	1157	
Asn	Tyr	Phe	Lys	Met	Ser	Lys	Arg	Met	Tyr	Pro	His	Arg	Pro	Val	Met		
				205					210					215			
atg	gtg	atc	agc	cac	gct	gcg	ccc	cac	ggc	ccc	gag	gac	tca	gcc	cca	1205	
Met	Val	Ile	Ser	His	Ala	Ala	Pro	His	Gly	Pro	Glu	Asp	Ser	Ala	Pro		
			220					225					230				
cag	ttt	tct	aaa	ctg	tac	ccc	aat	gct	tcc	caa	cac	ata	act	cct	agt	1253	
Gln	Phe	Ser	Lys	Leu	Tyr	Pro	Asn	Ala	Ser	Gln	His	Ile	Thr	Pro	Ser		
		235					240					245					
tat	aac	tat	gca	cca	aat	atg	gat	aaa	cac	tgg	att	atg	cag	tac	aca	1301	
Tyr	Asn	Tyr	Ala	Pro	Asn	Met	Asp	Lys	His	Trp	Ile	Met	Gln	Tyr	Thr		
	250					255					260						
gga	cca	atg	ctg	ccc	atc	cac	atg	gaa	ttt	aca	aac	att	cta	cag	cgc	1349	
Gly	Pro	Met	Leu	Pro	Ile	His	Met	Glu	Phe	Thr	Asn	Ile	Leu	Gln	Arg		
265					270					275					280		
aaa	agg	ctc	cag	act	ttg	atg	tca	gtg	gat	gat	tct	gtg	gag	agg	ctg	1397	
Lys	Arg	Leu	Gln	Thr	Leu	Met	Ser	Val	Asp	Asp	Ser	Val	Glu	Arg	Leu		
				285					290					295			
tat	aac	atg	ctc	gtg	gag	acg	ggg	gag	ctg	gag	aat	act	tac	atc	att	1445	
Tyr	Asn	Met	Leu	Val	Glu	Thr	Gly	Glu	Leu	Glu	Asn	Thr	Tyr	Ile	Ile		
			300					305					310				
tac	acc	gcc	gac	cat	ggt	tac	cat	att	ggg	cag	ttt	gga	ctg	gtc	aag	1493	
Tyr	Thr	Ala	Asp	His	Gly	Tyr	His	Ile	Gly	Gln	Phe	Gly	Leu	Val	Lys		
			315				320					325					
ggg	aaa	tcc	atg	cca	tat	gac	ttt	gat	att	cgt	gtg	cct	ttt	ttt	att	1541	
Gly	Lys	Ser	Met	Pro	Tyr	Asp	Phe	Asp	Ile	Arg	Val	Pro	Phe	Phe	Ile		
	330					335					340						

cgt ggt cca agt gta gaa cca gga tca ata gtc cca cag atc gtt ctc 1589
 Arg Gly Pro Ser Val Glu Pro Gly Ser Ile Val Pro Gln Ile Val Leu
 345 350 355 360

aac att gac ttg gcc ccc acg atc ctg gat att gct ggg ctc gac aca 1637
 Asn Ile Asp Leu Ala Pro Thr Ile Leu Asp Ile Ala Gly Leu Asp Thr
 365 370 375

cct cct gat gtg gac ggc aag tct gtc ctc aaa ctt ctg gac cca gaa 1685
 Pro Pro Asp Val Asp Gly Lys Ser Val Leu Lys Leu Leu Asp Pro Glu
 380 385 390

aag cca ggt aac agg ttt cga aca aac aag aag gcc aaa att tgg cgt 1733
 Lys Pro Gly Asn Arg Phe Arg Thr Asn Lys Lys Ala Lys Ile Trp Arg
 395 400 405

gat aca ttc cta gtg gaa aga ggc aaa ttt cta cgt aag aag gaa gaa 1781
 Asp Thr Phe Leu Val Glu Arg Gly Lys Phe Leu Arg Lys Lys Glu Glu
 410 415 420

tcc agc aag aat atc caa cag tca aat cac ttg ccc aaa tat gaa cgg 1829
 Ser Ser Lys Asn Ile Gln Gln Ser Asn His Leu Pro Lys Tyr Glu Arg
 425 430 435 440

gtc aaa gaa cta tgc cag cag gcc agg tac cag aca gcc tgt gaa caa 1877
 Val Lys Glu Leu Cys Gln Gln Ala Arg Tyr Gln Thr Ala Cys Glu Gln
 445 450 455

ccg ggg cag aag tgg caa tgc att gag gat aca tct ggc aag ctt cga 1925
 Pro Gly Gln Lys Trp Gln Cys Ile Glu Asp Thr Ser Gly Lys Leu Arg
 460 465 470

att cac aag tgt aaa gga ccc agt gac ctg ctc aca gtc cgg cag agc 1973
 Ile His Lys Cys Lys Gly Pro Ser Asp Leu Leu Thr Val Arg Gln Ser
 475 480 485

acg cgg aac ctc tac gct cgc ggc ttc cat gac aaa gac aaa gag tgc 2021
 Thr Arg Asn Leu Tyr Ala Arg Gly Phe His Asp Lys Asp Lys Glu Cys
 490 495 500

agt tgt agg gag tct ggt tac cgt gcc agc aga agc caa aga aag agt 2069
 Ser Cys Arg Glu Ser Gly Tyr Arg Ala Ser Arg Ser Gln Arg Lys Ser
 505 510 515 520

caa cgg caa ttc ttg aga aac cag ggg act cca aag tac aag ccc aga 2117
 Gln Arg Gln Phe Leu Arg Asn Gln Gly Thr Pro Lys Tyr Lys Pro Arg
 525 530 535

ttt gtc cat act cgg cag aca cgt tcc ttg tcc gtc gaa ttt gaa ggt 2165
 Phe Val His Thr Arg Gln Thr Arg Ser Leu Ser Val Glu Phe Glu Gly
 540 545 550

gaa ata tat gac ata aat ctg gaa gaa gaa gaa gaa ttg caa gtg ttg 2213
 Glu Ile Tyr Asp Ile Asn Leu Glu Glu Glu Glu Leu Gln Val Leu
 555 560 565

caa cca aga aac att gct aag cgt cat gat gaa ggc cac aag ggg cca 2261
 Gln Pro Arg Asn Ile Ala Lys Arg His Asp Glu Gly His Lys Gly Pro
 570 575 580

aga gat ctc cag gct tcc agt ggt ggc aac agg ggc agg atg ctg gca 2309
 Arg Asp Leu Gln Ala Ser Ser Gly Gly Asn Arg Gly Arg Met Leu Ala
 585 590 595 600

gat agc agc aac gcc gtg ggc cca cct acc act gtc cga gtg aca cac 2357
 Asp Ser Ser Asn Ala Val Gly Pro Pro Thr Thr Val Arg Val Thr His
 605 610 615

aag tgt ttt att ctt ccc aat gac tct atc cat tgt gag aga gaa ctg Lys Cys Phe Ile Leu Pro Asn Asp Ser Ile His Cys Glu Arg Glu Leu 620 625 630	2405
tac caa tcg gcc aga gcg tgg aag gac cat aag gca tac att gac aaa Tyr Gln Ser Ala Arg Ala Trp Lys Asp His Lys Ala Tyr Ile Asp Lys 635 640 645	2453
gag att gaa gct ctg caa gat aaa att aag aat tta aga gaa gtg aga Glu Ile Glu Ala Leu Gln Asp Lys Ile Lys Asn Leu Arg Glu Val Arg 650 655 660	2501
gga cat ctg aag aga agg aag cct gag gaa tgt agc tgc agt aaa caa Gly His Leu Lys Arg Arg Lys Pro Glu Glu Cys Ser Cys Ser Lys Gln 665 670 675 680	2549
agc tat tac aat aaa gag aaa ggt gta aaa aag caa gag aaa tta aag Ser Tyr Tyr Asn Lys Glu Lys Gly Val Lys Lys Gln Glu Lys Leu Lys 685 690 695	2597
agc cat ctt cac cca ttc aag gag gct gct cag gaa gta gat agc aaa Ser His Leu His Pro Phe Lys Glu Ala Ala Gln Glu Val Asp Ser Lys 700 705 710	2645
ctg caa ctt ttc aag gag aac aac cgt agg agg aag aag gag agg aag Leu Gln Leu Phe Lys Glu Asn Asn Arg Arg Arg Lys Lys Glu Arg Lys 715 720 725	2693
gag aag aga cgg cag agg aag ggg gaa gag tgc agc ctg cct ggc ctc Glu Lys Arg Arg Gln Arg Lys Gly Glu Glu Cys Ser Leu Pro Gly Leu 730 735 740	2741
act tgc ttc acg cat gac aac aac cac tgg cag aca gcc ccg ttc tgg Thr Cys Phe Thr His Asp Asn Asn His Trp Gln Thr Ala Pro Phe Trp 745 750 755 760	2789
aac ctg gga tct ttc tgt gct tgc acg agt tct aac aat aac acc tac Asn Leu Gly Ser Phe Cys Ala Cys Thr Ser Ser Asn Asn Asn Thr Tyr 765 770 775	2837
tgg tgt ttg cgt aca gtt aat gag acg cat aat ttt ctt ttc tgt gag Trp Cys Leu Arg Thr Val Asn Glu Thr His Asn Phe Leu Phe Cys Glu 780 785 790	2885
ttt gct act ggc ttt ttg gag tat ttt gat atg aat aca gat cct tat Phe Ala Thr Gly Phe Leu Glu Tyr Phe Asp Met Asn Thr Asp Pro Tyr 795 800 805	2933
cag ctc aca aat aca gtg cac acg gta gaa cga ggc att ttg aat cag Gln Leu Thr Asn Thr Val His Thr Val Glu Arg Gly Ile Leu Asn Gln 810 815 820	2981
cta cac gta caa cta atg gag ctc aga agc tgt caa gga tat aag cag Leu His Val Gln Leu Met Glu Leu Arg Ser Cys Gln Gly Tyr Lys Gln 825 830 835 840	3029
tgc aac cca aga cct aag aat ctt gat gtt gga aat aaa gat gga gga Cys Asn Pro Arg Pro Lys Asn Leu Asp Val Gly Asn Lys Asp Gly Gly 845 850 855	3077
agc tat gac cta cac aga gga cag tta tgg gat gga tgg gaa ggt taa Ser Tyr Asp Leu His Arg Gly Gln Leu Trp Asp Gly Trp Glu Gly * 860 865 870	3125
tcagccccgt ctactgcag acatcaactg gcaaggccta gaggagctac acagtgtgaa tgaaaacatc tatgagtaca gacaaaacta cagacttagt ctggtggact ggactaatta	3185 3245

```

cttgaaggat ttagatagag tatttgcact gctgaagagt cactatgagc aaaataaaac 3305
aaataagact caaactgctc aaagtgcagg gttcttggtt gtctctgctg agcacgctgt 3365
gtcaatggag atggcctctg ctgactcaga tgaagaccca aggcataagg ttgggaaaaac 3425
acctcatttg accttgccag ctgaccttca aacctgcgat ttgaaccgac caacattaag 3485
tccagagagt aaacttgaat ggaataacga cattccagaa gttaatcatt tgaattctga 3545
acactggaga aaaaccgaaa aatggacggg gcatgaagag actaatcatc tggaaaccga 3605
tttcagtggc gatggcatga cagagctaga gctcggggccc agccccaggc tgcagcccat 3665
tcgcaggcac ccgaaagaac ttccccagta tgggtggtcct ggaaaggaca tttttgaaga 3725
tcaactatat ctctctgtgc attccgatgg aatttcagtt catcagatgt tcaccatggc 3785
caccgcagaa caccgaagta attccagcat agcggggaag atgttgacca aggtggagaa 3845
gaatcacgaa aaggagaagt cacagcacct agaaggcagc gcctcctctt cactctcctc 3905
tgattagatg aaactgttac cttaccctaa acacagtatt tctttttaac ttttttattt 3965
gtaaactaat aaaggkaatc acagccacca acattccaag ctaccctggg tacctttgtg 4025
cagtagaagc tagtgagcat gtgagcaagc ggtgtgcaca cggagactca tcgttataat 4085
ttactatctg ccaaggagta gaaagaaagg ctggggatat ttggggttggc ttgggktttg 4145
attttttgct tggttggttg gtttgkacta aaacagtatt atcttttgaa tatcgtaggg 4205
acataarkww wwwmmwkkw wtcmawymra kakgsywrra wkgggstyty tskkrkstmw 4265
amwykwscmc cyskkrrwaw tywywmmwyc mykysstg rykrnktaat gaagtt 4321

```

<210> 5

<211> 569

<212> PRT

<213> homo sapiens

<400> 5

```

Met His Thr Leu Thr Gly Phe Ser Leu Val Ser Leu Leu Ser Phe Gly
1          5          10          15
Tyr Leu Ser Trp Asp Trp Ala Lys Pro Ser Phe Val Ala Asp Gly Pro
20          25          30
Gly Glu Ala Gly Glu Gln Pro Ser Ala Ala Pro Pro Gln Pro Pro His
35          40          45
Ile Ile Phe Ile Leu Thr Asp Asp Gln Gly Tyr His Asp Val Gly Tyr
50          55          60
His Gly Ser Asp Ile Glu Thr Pro Thr Leu Asp Arg Leu Ala Ala Lys
65          70          75          80
Gly Val Lys Leu Glu Asn Tyr Tyr Ile Gln Pro Ile Cys Thr Pro Ser
85          90          95
Arg Ser Gln Leu Leu Thr Gly Arg Tyr Gln Ile His Thr Gly Leu Gln
100         105         110
His Ser Ile Ile Arg Pro Gln Gln Pro Asn Cys Leu Pro Leu Asp Gln
115         120         125
Val Thr Leu Pro Gln Lys Leu Gln Glu Ala Gly Tyr Ser Thr His Met
130         135         140
Val Gly Lys Trp His Leu Gly Phe Tyr Arg Lys Glu Cys Leu Pro Thr
145         150         155         160
Arg Arg Gly Phe Asp Thr Phe Leu Gly Ser Leu Thr Gly Asn Val Asp
165         170         175
Tyr Tyr Thr Tyr Asp Asn Cys Asp Gly Pro Gly Val Cys Gly Phe Asp
180         185         190
Leu His Glu Gly Glu Asn Val Ala Trp Gly Leu Ser Gly Gln Tyr Ser
195         200         205
Thr Met Leu Tyr Ala Gln Arg Ala Ser His Ile Leu Ala Ser His Ser
210         215         220
Pro Gln Arg Pro Leu Phe Leu Tyr Val Ala Phe Gln Ala Val His Thr
225         230         235         240
Pro Leu Gln Ser Pro Arg Glu Tyr Leu Tyr Arg Tyr Arg Thr Met Gly
245         250         255
Asn Val Ala Arg Arg Lys Tyr Ala Ala Met Val Thr Cys Met Asp Glu
260         265         270
Ala Val Arg Asn Ile Thr Trp Ala Leu Lys Arg Tyr Gly Phe Tyr Asn
275         280         285
Asn Ser Val Ile Ile Phe Ser Ser Asp Asn Gly Gly Gln Thr Phe Ser
290         295         300
Gly Gly Ser Asn Trp Pro Leu Arg Gly Arg Lys Gly Thr Tyr Trp Glu
305         310         315         320
Gly Gly Val Arg Gly Leu Gly Phe Val His Ser Pro Leu Leu Lys Arg
325         330         335

```

Lys Gln Arg Thr Ser Arg Ala Leu Met His Ile Thr Asp Trp Tyr Pro
 340 345 350
 Thr Leu Val Gly Leu Ala Gly Gly Thr Thr Ser Ala Ala Asp Gly Leu
 355 360 365
 Asp Gly Tyr Asp Val Trp Pro Ala Ile Ser Glu Gly Arg Ala Ser Pro
 370 375 380
 Arg Thr Glu Ile Leu His Asn Ile Asp Pro Leu Tyr Asn His Ala Gln
 385 390 395 400
 His Gly Ser Leu Glu Gly Gly Phe Gly Ile Trp Asn Thr Ala Val Gln
 405 410 415
 Ala Ala Ile Arg Val Gly Glu Trp Lys Leu Leu Thr Gly Asp Pro Gly
 420 425 430
 Tyr Gly Asp Trp Ile Pro Pro Gln Thr Leu Ala Thr Phe Pro Gly Ser
 435 440 445
 Trp Trp Asn Leu Glu Arg Met Ala Ser Val Arg Gln Ala Val Trp Leu
 450 455 460
 Phe Asn Ile Ser Ala Asp Pro Tyr Glu Arg Glu Asp Leu Ala Gly Gln
 465 470 475 480
 Arg Pro Asp Val Val Arg Thr Leu Leu Ala Arg Leu Ala Glu Tyr Asn
 485 490 495
 Arg Thr Ala Ile Pro Val Arg Tyr Pro Ala Glu Asn Pro Arg Ala His
 500 505 510
 Pro Asp Phe Asn Gly Gly Ala Trp Gly Pro Trp Ala Ser Asp Glu Glu
 515 520 525
 Glu Glu Glu Glu Gly Arg Ala Arg Ser Phe Ser Arg Gly Arg Arg
 530 535 540
 Lys Lys Lys Cys Lys Ile Cys Lys Leu Arg Ser Phe Phe Arg Lys Leu
 545 550 555 560
 Asn Thr Arg Leu Met Ser Gln Arg Ile
 565

<210> 6
 <211> 2940
 <212> DNA
 <213> homo sapiens

<220>
 <221> CDS
 <222> (334)...(2043)

<400> 6
 ccacgcgtcc gccacgcgt ccggtgccca cgccgcgtct caggctggcc gggctgagcc 60
 ggggaagagg gagcaaaggc ggcgcagggc ctgcgcttag gcagcgggag gcagctcggc 120
 gcgggcctga cctcccaga gcgcccgcgt gcggccgagc agatccggcc cagccgtccg 180
 gcagccagtc ccggaccaga cactggaccg tccccggggg gcgctgaact ccctcgcagc 240
 atccgagccg gcgggcgggt ggtgcgccct gggcgcgcca ggtggtgagg cccaggagc 300
 ccggcgcgcc gggacacgcg ggccggcttg gcg atg cac acc ctc act ggc ttc 354
 Met His Thr Leu Thr Gly Phe
 1 5

 tct ctg gtc agc ctg ctc agc ttc ggc tac ctg tcc tgg gac tgg gcc 402
 Ser Leu Val Ser Leu Leu Ser Phe Gly Tyr Leu Ser Trp Asp Trp Ala
 10 15 20

 aag ccg agc ttc gtg gcc gac ggg ccc ggg gag gct ggc gag cag ccc 450
 Lys Pro Ser Phe Val Ala Asp Gly Pro Gly Glu Ala Gly Glu Gln Pro
 25 30 35

 tcg gcc gct ccg ccc cag cct ccc cac atc atc ttc atc ctc acg gac 498
 Ser Ala Ala Pro Pro Gln Pro Pro His Ile Ile Phe Ile Leu Thr Asp
 40 45 50 55

 gac caa ggc tac cac gac gtg ggc tac cat ggt tca gat atc gag acc 546
 Asp Gln Gly Tyr His Asp Val Gly Tyr His Gly Ser Asp Ile Glu Thr
 60 65 70

 cct acg ctg gac agg ctg gcg gcc aag ggg gtc aag ttg gag aat tat 594

Pro Thr Leu Asp Arg Leu Ala Ala Lys Gly Val Lys Leu Glu Asn Tyr	
75 80 85	
tac atc cag ccc atc tgc acg cct tcg cgg agc cag ctc ctc act ggc	642
Tyr Ile Gln Pro Ile Cys Thr Pro Ser Arg Ser Gln Leu Leu Thr Gly	
90 95 100	
agg tac cag atc cac aca gga ctc cag cat tcc atc atc cgc cca cag	690
Arg Tyr Gln Ile His Thr Gly Leu Gln His Ser Ile Ile Arg Pro Gln	
105 110 115	
cag ccc aac tgc ctg ccc ctg gac cag gtg aca ctg cca cag aag ctg	738
Gln Pro Asn Cys Leu Pro Leu Asp Gln Val Thr Leu Pro Gln Lys Leu	
120 125 130 135	
cag gag gca ggt tat tcc acc cat atg gtg ggc aag tgg cac ctg ggc	786
Gln Glu Ala Gly Tyr Ser Thr His Met Val Gly Lys Trp His Leu Gly	
140 145 150	
ttc tac cgg aag gag tgt ctg ccc acc cgt cgg ggc ttc gac acc ttc	834
Phe Tyr Arg Lys Glu Cys Leu Pro Thr Arg Arg Gly Phe Asp Thr Phe	
155 160 165	
ctg ggc tcg ctc acg ggc aat gtg gac tat tac acc tat gac aac tgt	882
Leu Gly Ser Leu Thr Gly Asn Val Asp Tyr Tyr Thr Tyr Asp Asn Cys	
170 175 180	
gat ggc cca ggc gtg tgc ggc ttc gac ctg cac gag ggt gag aat gtg	930
Asp Gly Pro Gly Val Cys Gly Phe Asp Leu His Glu Gly Glu Asn Val	
185 190 195	
gcc tgg ggg ctc agc ggc cag tac tcc act atg ctt tac gcc cag cgc	978
Ala Trp Gly Leu Ser Gly Gln Tyr Ser Thr Met Leu Tyr Ala Gln Arg	
200 205 210 215	
gcc agc cat atc ctg gcc agc cac agc cct cag cgt ccc ctc ttc ctc	1026
Ala Ser His Ile Leu Ala Ser His Ser Pro Gln Arg Pro Leu Phe Leu	
220 225 230	
tat gtg gcc ttc cag gca gta cac aca ccc ctg cag tcc cct cgt gag	1074
Tyr Val Ala Phe Gln Ala Val His Thr Pro Leu Gln Ser Pro Arg Glu	
235 240 245	
tac ctg tac cgc tac cgc acc atg ggc aat gtg gcc cgg cgg aag tac	1122
Tyr Leu Tyr Arg Tyr Arg Thr Met Gly Asn Val Ala Arg Arg Lys Tyr	
250 255 260	
gcg gcc atg gtg acc tgc atg gat gag gct gtg cgc aac atc acc tgg	1170
Ala Ala Met Val Thr Cys Met Asp Glu Ala Val Arg Asn Ile Thr Trp	
265 270 275	
gcc ctc aag cgc tac ggt ttc tac aac aac agt gtc atc atc ttc tcc	1218
Ala Leu Lys Arg Tyr Gly Phe Tyr Asn Asn Ser Val Ile Ile Phe Ser	
280 285 290 295	
agt gac aat ggt ggc cag act ttc tcg ggg ggc agc aac tgg ccg ctc	1266
Ser Asp Asn Gly Gly Gln Thr Phe Ser Gly Gly Ser Asn Trp Pro Leu	
300 305 310	
cga gga cgc aag ggc act tat tgg gaa ggt ggc gtg cgg ggc cta ggc	1314
Arg Gly Arg Lys Gly Thr Tyr Trp Glu Gly Gly Val Arg Gly Leu Gly	
315 320 325	
ttt gtc cac agt ccc ctg ctc aag cga aag caa cgg aca agc cgg gca	1362
Phe Val His Ser Pro Leu Leu Lys Arg Lys Gln Arg Thr Ser Arg Ala	
330 335 340	

ctg atg cac atc act gac tgg tac ccg acc ctg gtg ggt ctg gca ggt Leu Met His Ile Thr Asp Trp Tyr Pro Thr Leu Val Gly Leu Ala Gly 345 350 355	1410
ggt acc acc tca gca gcc gat ggg cta gat ggc tac gac gtg tgg ccg Gly Thr Thr Ser Ala Ala Asp Gly Leu Asp Gly Tyr Asp Val Trp Pro 360 365 370 375	1458
gcc atc agc gag ggc cgg gcc tca cca cgc acg gag atc ctg cac aac Ala Ile Ser Glu Gly Arg Ala Ser Pro Arg Thr Glu Ile Leu His Asn 380 385 390	1506
att gac cca ctc tac aac cat gcc cag cat ggc tcc ctg gag ggc ggc Ile Asp Pro Leu Tyr Asn His Ala Gln His Gly Ser Leu Glu Gly Gly 395 400 405	1554
ttt ggc atc tgg aac acc gcc gtg cag gct gcc atc cgc gtg ggt gag Phe Gly Ile Trp Asn Thr Ala Val Gln Ala Ala Ile Arg Val Gly Glu 410 415 420	1602
tgg aag ctg ctg aca gga gac ccc ggc tat ggc gat tgg atc cca ccg Trp Lys Leu Leu Thr Gly Asp Pro Gly Tyr Gly Asp Trp Ile Pro Pro 425 430 435	1650
cag aca ctg gcc acc ttc ccg ggt agc tgg tgg aac ctg gaa cga atg Gln Thr Leu Ala Thr Phe Pro Gly Ser Trp Trp Asn Leu Glu Arg Met 440 445 450 455	1698
gcc agt gtc cgc cag gcc gtg tgg ctc ttc aac atc agt gct gac cct Ala Ser Val Arg Gln Ala Val Trp Leu Phe Asn Ile Ser Ala Asp Pro 460 465 470	1746
tat gaa cgg gag gac ctg gct ggc cag cgg cct gat gtg gtc cgc acc Tyr Glu Arg Glu Asp Leu Ala Gly Gln Arg Pro Asp Val Val Arg Thr 475 480 485	1794
ctg ctg gct cgc ctg gcc gaa tat aac cgc aca gcc atc ccg gta cgc Leu Leu Ala Arg Leu Ala Glu Tyr Asn Arg Thr Ala Ile Pro Val Arg 490 495 500	1842
tac cca gct gag aac ccc cgg gct cat cct gac ttt aat ggg ggt gct Tyr Pro Ala Glu Asn Pro Arg Ala His Pro Asp Phe Asn Gly Gly Ala 505 510 515	1890
tgg ggg ccc tgg gcc agt gat gag gaa gag gag gaa gag gaa ggg agg Trp Gly Pro Trp Ala Ser Asp Glu Glu Glu Glu Glu Glu Glu Gly Arg 520 525 530 535	1938
gct cga agc ttc tcc cgg ggt cgt cgc aag aaa aaa tgc aag att tgc Ala Arg Ser Phe Ser Arg Gly Arg Arg Lys Lys Lys Cys Lys Ile Cys 540 545 550	1986
aag ctt cga tcc ttt ttc cgt aaa ctc aac acc agg cta atg tcc caa Lys Leu Arg Ser Phe Phe Arg Lys Leu Asn Thr Arg Leu Met Ser Gln 555 560 565	2034
cgg atc tga tgggtggggag ggagaaaact gtcctttaga ggatcttccc Arg Ile *	2083
caactccggct tggccctgct gtttctcagg gagaagcctg tcacatctcc atctacaggg agttggaggg tgtagagtcc cttggttgaa cagggtaggg agcctggata ggagtggtg ggaataaacc agactgggat gcctgtgtct cagtcctgcc tccctcacgga cttgctctgt gacctcaggt gacccacatg agcttttagc ctcagtttcc tcatctgtaa aatgagctct aatgactttg tgactctttg gtgtggccct ggagcctggg gccacggtgg agttcctggc cggccttgcc acttgacaac tcctttaagg cttccccctt aacacgggat ccctgtgggtg gtgtttgga gttgcctgga ggcaactcca agcctggccc ccagctgaag catggcaatc 2143 2203 2263 2323 2383 2443 2503	

```

tggctgctct ctacaggac ccccaagcgc tgtgggtgga gggcaggggt cggggggggt 2563
gaccttcttg ggtcttcaca tggcctaggc cagtcctccg gtcagactgg tgtcaggcac 2623
cgtggtgcaa aattcctctt ctggcccctc cagtaccag agaaactggc tgggccatta 2683
actgctgcag caccaagggg ggtagaaaga gctgtgaaga gccccaaac cagtaccagg 2743
acacctgggt tctcctgtga cctggggcac agttcttgcc ctctaggcct tgatttcccc 2803
acctgcaagt ggggatgcca gccctggctc tgcctccttc atgaggctct ggaagactgg 2863
ccaaggttgt ggaggagctt gtgaacttga ttaaagtgtc gtaacatgga aaaaaaaaaa 2923
aaaaaaaaaa agggcggg                                     2940

```

<210> 7

<211> 599

<212> PRT

<213> homo sapiens

<400> 7

```

Met Ala Pro Arg Gly Cys Ala Gly His Pro Pro Pro Pro Ser Pro Gln
1          5          10          15
Ala Cys Val Cys Pro Gly Lys Met Leu Ala Met Gly Ala Leu Ala Gly
20          25          30
Phe Trp Ile Leu Cys Leu Leu Thr Tyr Gly Tyr Leu Ser Trp Gly Gln
35          40          45
Ala Leu Glu Glu Glu Glu Gly Ala Leu Leu Ala Gln Ala Gly Glu
50          55          60
Lys Leu Glu Pro Ser Thr Thr Ser Thr Ser Gln Pro His Leu Ile Phe
65          70          75          80
Ile Leu Ala Asp Asp Gln Gly Phe Arg Asp Val Gly Tyr His Gly Ser
85          90          95
Glu Ile Lys Thr Pro Thr Leu Asp Lys Leu Ala Ala Glu Gly Val Lys
100         105         110
Leu Glu Asn Tyr Tyr Val Gln Pro Ile Cys Thr Pro Ser Arg Ser Gln
115         120         125
Phe Ile Thr Gly Lys Tyr Gln Ile His Thr Gly Leu Gln His Ser Ile
130         135         140
Ile Arg Pro Thr Gln Pro Asn Cys Leu Pro Leu Asp Asn Ala Thr Leu
145         150         155         160
Pro Gln Lys Leu Lys Glu Val Gly Tyr Ser Thr His Met Val Gly Lys
165         170         175
Trp His Leu Gly Phe Tyr Arg Lys Glu Cys Met Pro Thr Arg Arg Gly
180         185         190
Phe Asp Thr Phe Phe Gly Ser Leu Leu Gly Ser Gly Asp Tyr Tyr Thr
195         200         205
His Tyr Lys Cys Asp Ser Pro Gly Met Cys Gly Tyr Asp Leu Tyr Glu
210         215         220
Asn Asp Asn Ala Ala Trp Asp Tyr Asp Asn Gly Ile Tyr Ser Thr Gln
225         230         235         240
Met Tyr Thr Gln Arg Val Gln Gln Ile Leu Ala Ser His Asn Pro Thr
245         250         255
Lys Pro Ile Phe Leu Tyr Ile Ala Tyr Gln Ala Val His Ser Pro Leu
260         265         270
Gln Ala Pro Gly Arg Tyr Phe Glu His Tyr Arg Ser Ile Ile Asn Ile
275         280         285
Asn Arg Arg Arg Tyr Ala Ala Met Leu Ser Cys Leu Asp Glu Ala Ile
290         295         300
Asn Asn Val Thr Leu Ala Leu Lys Thr Tyr Gly Phe Tyr Asn Asn Ser
305         310         315         320
Ile Ile Ile Tyr Ser Ser Asp Asn Gly Gly Gln Pro Thr Ala Gly Gly
325         330         335
Ser Asn Trp Pro Leu Arg Gly Ser Lys Gly Thr Tyr Trp Glu Gly Gly
340         345         350
Ile Arg Ala Val Gly Phe Val His Ser Pro Leu Leu Lys Asn Lys Gly
355         360         365
Thr Val Cys Lys Glu Leu Val His Ile Thr Asp Trp Tyr Pro Thr Leu
370         375         380
Ile Ser Leu Ala Glu Gly Gln Ile Asp Glu Asp Ile Gln Leu Asp Gly
385         390         395         400
Tyr Asp Ile Trp Glu Thr Ile Ser Glu Gly Leu Arg Ser Pro Arg Val
405         410         415

```


Asp Ile Leu His Asn Ile Asp Pro Ile Tyr Thr Lys Ala Lys Asn Gly
 420 425 430
 Ser Trp Ala Ala Gly Tyr Gly Ile Trp Asn Thr Ala Ile Gln Ser Ala
 435 440 445
 Ile Arg Val Gln His Trp Lys Leu Leu Thr Gly Asn Pro Gly Tyr Ser
 450 455 460
 Asp Trp Val Pro Pro Gln Ser Phe Ser Asn Leu Gly Pro Asn Arg Trp
 465 470 475 480
 His Asn Glu Arg Ile Thr Leu Ser Thr Gly Lys Ser Val Trp Leu Phe
 485 490 495
 Asn Ile Thr Ala Asp Pro Tyr Glu Arg Val Asp Leu Ser Asn Arg Tyr
 500 505 510
 Pro Gly Ile Val Lys Lys Leu Leu Arg Arg Leu Ser Gln Phe Asn Lys
 515 520 525
 Thr Ala Val Pro Val Arg Tyr Pro Pro Lys Asp Pro Arg Ser Asn Pro
 530 535 540
 Arg Leu Asn Gly Gly Val Trp Gly Pro Trp Tyr Lys Glu Glu Thr Lys
 545 550 555 560
 Lys Lys Lys Pro Ser Lys Asn Gln Ala Glu Lys Lys Gln Lys Lys Ser
 565 570 575
 Lys Lys Lys Lys Lys Gln Gln Lys Ala Val Ser Gly Ser Thr Cys
 580 585 590
 His Ser Gly Val Thr Cys Gly
 595

<210> 8

<211> 2253

<212> DNA

<213> homo sapiens

<220>

<221> CDS

<222> (324)... (2123)

<400> 8

cacgcgctccg cccacgcgctc cgtggagata ttaacttttt tctttttttt tttccttggt 60
 ggaagctgct ctaggaggagg gggaggagga ggagaaagt aaatgtgctg gagaagagcg 120
 agccctcctt gttcttccgg agtcccatcc attaaagcc cacttctgga agattaaagt 180
 tgtcggacat ggtgacagct gagaggagag gaggatttct tgccagggtg agagtcttca 240
 ccgtctgttg ggtgcatgtg tgcgccgca gcggcgcggg gcgcgtggtt ctccgcgtgg 300
 agtctcacct gggacctgag tga atg gct ccc agg ggc tgt gcg ggg cat ccg 353
 Met Ala Pro Arg Gly Cys Ala Gly His Pro
 1 5 10

 cct ccg cct tct cca cag gcc tgt gtc tgt cct gga aag atg cta gca 401
 Pro Pro Pro Ser Pro Gln Ala Cys Val Cys Pro Gly Lys Met Leu Ala
 15 20 25

 atg ggg gcg ctg gca gga ttc tgg atc ctc tgc ctc ctc act tat ggt 449
 Met Gly Ala Leu Ala Gly Phe Trp Ile Leu Cys Leu Leu Thr Tyr Gly
 30 35 40

 tac ctg tcc tgg ggc cag gcc tta gaa gag gag gaa gaa ggg gcc tta 497
 Tyr Leu Ser Trp Gly Gln Ala Leu Glu Glu Glu Glu Glu Gly Ala Leu
 45 50 55

 cta gct caa gct gga gag aaa cta gag ccc agc aca act tcc acc tcc 545
 Leu Ala Gln Ala Gly Glu Lys Leu Glu Pro Ser Thr Thr Ser Thr Ser
 60 65 70

 cag ccc cat ctc att ttc atc cta gcg gat gat cag gga ttt aga gat 593
 Gln Pro His Leu Ile Phe Ile Leu Ala Asp Asp Gln Gly Phe Arg Asp
 75 80 85 90

 gtg ggt tac cac gga tct gag att aaa aca cct act ctt gac aag ctc 641
 Val Gly Tyr His Gly Ser Glu Ile Lys Thr Pro Thr Leu Asp Lys Leu
 95 100 105

gct gcc gaa gga gtt aaa ctg gag aac tac tat gtc cag cct att tgc 689
Ala Ala Glu Gly Val Lys Leu Glu Asn Tyr Tyr Val Gln Pro Ile Cys
110 115 120

aca cca tcc agg agt cag ttt att act gga aag tat cag ata cac acc 737
Thr Pro Ser Arg Ser Gln Phe Ile Thr Gly Lys Tyr Gln Ile His Thr
125 130 135

gga ctt caa cat tct atc ata aga cct acc caa ccc aac tgt tta cct 785
Gly Leu Gln His Ser Ile Ile Arg Pro Thr Gln Pro Asn Cys Leu Pro
140 145 150

ctg gac aat gcc acc cta cct cag aaa ctg aag gag gtt gga tat tca 833
Leu Asp Asn Ala Thr Leu Pro Gln Lys Leu Lys Glu Val Gly Tyr Ser
155 160 165 170

acg cat atg gtc gga aaa tgg cac ttg ggt ttt tac aga aaa gaa tgc 881
Thr His Met Val Gly Lys Trp His Leu Gly Phe Tyr Arg Lys Glu Cys
175 180 185

atg ccc acc aga aga gga ttt gat acc ttt ttt ggt tcc ctt ttg gga 929
Met Pro Thr Arg Arg Gly Phe Asp Thr Phe Phe Gly Ser Leu Leu Gly
190 195 200

agt ggg gat tac tat aca cac tac aaa tgt gac agt cct ggg atg tgt 977
Ser Gly Asp Tyr Tyr Thr His Tyr Lys Cys Asp Ser Pro Gly Met Cys
205 210 215

ggc tat gac ttg tat gaa aac gac aat gct gcc tgg gac tat gac aat 1025
Gly Tyr Asp Leu Tyr Glu Asn Asp Asn Ala Ala Trp Asp Tyr Asp Asn
220 225 230

ggc ata tac tcc aca cag atg tac act cag aga gta cag caa atc tta 1073
Gly Ile Tyr Ser Thr Gln Met Tyr Thr Gln Arg Val Gln Gln Ile Leu
235 240 245 250

gct tcc cat aac ccc aca aag cct ata ttt tta tat att gcc tat caa 1121
Ala Ser His Asn Pro Thr Lys Pro Ile Phe Leu Tyr Ile Ala Tyr Gln
255 260 265

gct gtt cat tca cca ctg caa gct cct ggc agg tat ttc gaa cac tac 1169
Ala Val His Ser Pro Leu Gln Ala Pro Gly Arg Tyr Phe Glu His Tyr
270 275 280

cga tcc att atc aac ata aac agg agg aga tat gct gcc atg ctt tcc 1217
Arg Ser Ile Ile Asn Ile Asn Arg Arg Arg Tyr Ala Ala Met Leu Ser
285 290 295

tgc tta gat gaa gca atc aac aac gtg aca ttg gct cta aag act tat 1265
Cys Leu Asp Glu Ala Ile Asn Asn Val Thr Leu Ala Leu Lys Thr Tyr
300 305 310

ggt ttc tat aac aac agc att atc att tac tct tca gat aat ggt ggc 1313
Gly Phe Tyr Asn Asn Ser Ile Ile Ile Tyr Ser Ser Asp Asn Gly Gly
315 320 325 330

cag cct acg gca gga ggg agt aac tgg cct ctc aga ggt agc aaa gga 1361
Gln Pro Thr Ala Gly Gly Ser Asn Trp Pro Leu Arg Gly Ser Lys Gly
335 340 345

aca tat tgg gaa gga ggg atc cgg gct gta ggc ttt gtg cat agc cca 1409
Thr Tyr Trp Glu Gly Gly Ile Arg Ala Val Gly Phe Val His Ser Pro
350 355 360

ctt ctg aaa aac aag gga aca gtg tgt aag gaa ctt gtg cac atc act 1457
Leu Leu Lys Asn Lys Gly Thr Val Cys Lys Glu Leu Val His Ile Thr

365	370	375	
gac tgg tac ccc act ctc att tca ctg gct gaa gga cag att gat gag			1505
Asp Trp Tyr Pro Thr Leu Ile Ser Leu Ala Glu Gly Gln Ile Asp Glu			
380	385	390	
gac att caa cta gat ggc tat gat atc tgg gag acc ata agt gag ggt			1553
Asp Ile Gln Leu Asp Gly Tyr Asp Ile Trp Glu Thr Ile Ser Glu Gly			
395	400	405	410
ctt cgc tca ccc cga gta gat att ttg cat aac att gac ccc ata tac			1601
Leu Arg Ser Pro Arg Val Asp Ile Leu His Asn Ile Asp Pro Ile Tyr			
	415	420	425
acc aag gca aaa aat ggc tcc tgg gca gca ggc tat ggg atc tgg aac			1649
Thr Lys Ala Lys Asn Gly Ser Trp Ala Ala Gly Tyr Gly Ile Trp Asn			
	430	435	440
act gca atc cag tca gcc atc aga gtg cag cac tgg aaa ttg ctt aca			1697
Thr Ala Ile Gln Ser Ala Ile Arg Val Gln His Trp Lys Leu Leu Thr			
	445	450	455
gga aat cct ggc tac agc gac tgg gtc ccc cct cag tct ttc agc aac			1745
Gly Asn Pro Gly Tyr Ser Asp Trp Val Pro Pro Gln Ser Phe Ser Asn			
	460	465	470
ctg gga ccg aac cgg tgg cac aat gaa cgg atc acc ttg tca act ggc			1793
Leu Gly Pro Asn Arg Trp His Asn Glu Arg Ile Thr Leu Ser Thr Gly			
	475	480	485
aaa agt gta tgg ctt ttc aac atc aca gcc gac cca tat gag agg gtg			1841
Lys Ser Val Trp Leu Phe Asn Ile Thr Ala Asp Pro Tyr Glu Arg Val			
	495	500	505
gac cta tct aac agg tat cca gga atc gtg aag aag ctc cta cgg agg			1889
Asp Leu Ser Asn Arg Tyr Pro Gly Ile Val Lys Lys Leu Leu Arg Arg			
	510	515	520
ctc tca cag ttc aac aaa act gca gtg ccg gtc agg tat ccc ccc aaa			1937
Leu Ser Gln Phe Asn Lys Thr Ala Val Pro Val Arg Tyr Pro Pro Lys			
	525	530	535
gac ccc aga agt aac cct agg ctc aat gga ggg gtc tgg gga cca tgg			1985
Asp Pro Arg Ser Asn Pro Arg Leu Asn Gly Gly Val Trp Gly Pro Trp			
	540	545	550
tat aaa gag gaa acc aag aaa aag aag cca agc aaa aat cag gct gag			2033
Tyr Lys Glu Glu Thr Lys Lys Lys Lys Pro Ser Lys Asn Gln Ala Glu			
	555	560	565
aaa aag caa aag aaa agc aaa aaa aag aag aag aaa cag cag aaa gca			2081
Lys Lys Gln Lys Lys Ser Lys Lys Lys Lys Lys Lys Gln Gln Lys Ala			
	575	580	585
gtc tca ggt tca act tgc cat tca ggt gtt act tgt gga taa			2123
Val Ser Gly Ser Thr Cys His Ser Gly Val Thr Cys Gly *			
	590	595	
gcacaaatat ttctgttttg gttaaacttt aatcagttct tatctttcat ctgtttccta			2183
ggtaaacacag caaatttggc tcgataatat cgctggccta agcgtcaggc ttgttttcat			2243
gctgtgccac			2253

<210> 9

<211> 552

<212> PRT

<213> Artificial Sequence

<220>

<223> Pfam consensus sequence for human sulfatase

<400> 9

```

Pro Asn Ile Leu Leu Ile Leu Ala Asp Asp Leu Gly Ile Gly Asp Leu
1      5      10      15
Gly Cys Tyr Gly Asn Pro Thr Ile Arg Thr Pro Asn Ile Asp Arg Leu
20      25      30
Ala Glu Glu Gly Leu Arg Phe Thr Asn Ala Tyr Val Thr Thr Pro Leu
35      40      45
Cys Thr Pro Ser Arg Ala Ala Leu Leu Thr Gly Arg Tyr Pro His Arg
50      55      60
Thr Gly Met Tyr Thr Asn Asn Arg Ala Gly Val Leu Pro Phe Thr Gly
65      70      75      80
Trp Ser Leu Glu Gly Gly Leu Pro Leu Asp Glu Thr Thr Leu Pro Glu
85      90      95
Leu Leu Lys Glu Ala Gly Tyr Ala Thr Gly Met Val Gly Lys Trp His
100     105     110
Gly Tyr Asn Glu Glu Ser Ser Ala Ser Asp Phe Ala His Leu Pro Leu
115     120     125
Gly Arg Gly Phe Asp Tyr Phe Tyr Gly Asn Leu Gly Gly Glu Asp Gln
130     135     140
Trp Tyr Pro Leu Val Asp Ala Leu Leu Pro Phe Thr Asn Asp Thr Tyr
145     150     155     160
Thr Cys Glu Gly Gly Tyr Gly Phe Ser Lys Asp Val Ala Leu Lys Pro
165     170     175
Leu Gly Ala Leu Gly Val Asn Glu Val Glu Ala Pro Asp Lys Ala Leu
180     185     190
Ala Asp Tyr Lys Thr Ala Gly Ala Leu Asn Val Pro His His Val Phe
195     200     205
Glu Trp Ala Asp Arg Tyr Ala Gly Ala Val Asp Val Gly Arg Pro Phe
210     215     220
Leu Ala Val Leu Ile Phe Pro Arg Pro Ala Ala Cys Phe Leu Tyr Pro
225     230     235     240
Asn Ala Thr Val Val Ser Gln Pro Met Pro His Ser Pro Leu Thr Ala
245     250     255
Pro Arg Pro Trp Gln Leu Leu Ala Asp Glu Ala Leu Pro Phe Leu Glu
260     265     270
Arg Asn Gly Gln Arg Asp Lys Pro Phe Phe Leu Tyr Leu Ser Tyr Lys
275     280     285
His Val His Ile Pro Arg Asp Ala Pro Met Leu Phe Ser Ser Lys Asp
290     295     300
Phe Ala Gly Ser Ser Arg Arg Gly Leu Tyr Gly Leu Ile Leu Asp Ser
305     310     315     320
Val Glu Glu Met Asp Asp Gly Val Gly Arg Val Leu Asn Ala Leu Asp
325     330     335
Glu Leu Asn Gly Leu Leu Asp Asn Thr Leu Ile Ile Phe Thr Ser Leu
340     345     350
Leu Asp His Gly Gly His Leu Gly Ala His Gly His Leu Gly Ile Arg
355     360     365
Ala Gly Gly Ser Asn Gly Pro Phe Arg Gly Gly Lys Gly Thr Asn Leu
370     375     380
Tyr Glu Gly Gly Thr Arg Val Pro Leu Ile Val Arg Trp Pro Glu Gly
385     390     395     400
Ile Ile Ala Pro Gly Gln Val Ser Asp Glu Leu Val Ser Leu Met Asp
405     410     415
Leu Phe Pro Thr Ile Leu Asp Leu Ala Gly Ala Pro Leu Pro Gly Val
420     425     430
Ala Ala Gly Val Lys Asp Arg Ile Leu Asp Gly Val Ser Leu Leu Pro
435     440     445
Leu Leu Leu Gly Ala Ala Gly Ser Ser Arg His Glu Thr Leu Phe Tyr
450     455     460
Glu Ser Tyr Cys Asn Glu Gly Arg Gly Phe Leu Pro Ala Val Arg Trp
465     470     475     480
Gly Lys Lys Lys Ala His Phe Arg Thr Pro Asn Ile Ala Gly Trp Gln
485     490     495
Arg Val Asp Phe Asp Asp Val Trp Lys Leu Phe Asn Thr Val Glu Asp

```

			500					505					510				
Phe	Asn	Arg	Ser	Gly	Asp	Asp	Ala	Cys	Arg	His	Gly	Asp	Val	Cys	Lys		
		515					520					525					
Cys	Leu	Gly	Lys	Pro	Arg	Arg	Ser	Val	Thr	His	His	Asp	Pro	Pro	Leu		
	530					535					540						
Leu	Tyr	Asp	Leu	Ser	Arg	Asp	Pro										
545					550												

<210> 10

<211> 520

<212> PRT

<213> Artificial Sequence

<220>

<223> Pfam consensus sequence for human sulfatase

<400> 10

Pro	Asn	Val	Leu	Leu	Ile	Leu	Ala	Asp	Asp	Leu	Gly	Ile	Gly	Asp	Leu		
1			5					10					15				
Gly	Cys	Tyr	Gly	His	Pro	Thr	Ile	Arg	Thr	Pro	Asn	Leu	Asp	Arg	Leu		
		20						25					30				
Ala	Glu	Glu	Gly	Leu	Arg	Phe	Thr	Asn	His	Tyr	Thr	Ala	Thr	Pro	Leu		
	35					40						45					
Cys	Ser	Pro	Ser	Arg	Ala	Ala	Leu	Leu	Thr	Gly	Arg	Tyr	Pro	His	Arg		
	50				55						60						
His	Gly	Met	Val	Ser	Asn	Gly	Arg	Leu	Gly	Val	Leu	Gly	Phe	Thr	Ala		
65				70					75					80			
Lys	Ser	Gly	Gly	Leu	Pro	Leu	Asp	Glu	Thr	Thr	Leu	Pro	Glu	Leu	Leu		
		85						90					95				
Lys	Glu	Ala	Gly	Tyr	Ala	Thr	Gly	Leu	Val	Gly	Lys	Trp	His	Leu	Gly		
	100						105						110				
Leu	Asn	Glu	Asn	Ser	Asp	Ala	Ala	Gly	Asp	Gly	Glu	His	Leu	Pro	Leu		
	115					120						125					
Gly	Trp	Arg	Gly	Phe	Asp	Tyr	Phe	Asp	Gly	Phe	Leu	Tyr	Gly	Ser	Pro		
	130				135						140						
Phe	Thr	Tyr	Asp	Glu	Glu	Asn	Cys	Asp	Asn	Gly	Glu	Gly	Thr	Glu	Pro		
145				150					155					160			
Pro	Glu	Ala	Tyr	Pro	Glu	Gln	Gly	Trp	Leu	Pro	Gln	Ile	Leu	Gly	Tyr		
		165						170					175				
Tyr	Leu	Thr	Asp	Leu	Leu	Ala	Asp	Lys	Ala	Leu	Gly	Leu	Leu	Asp	Val		
	180							185					190				
Ala	Ser	Ala	Ala	Gly	Arg	Leu	Leu	Ala	Lys	Ala	Leu	Ala	Ala	Ser	Arg		
	195					200						205					
Pro	Phe	Phe	Leu	Tyr	Ile	Ser	Pro	Pro	Ala	Pro	His	Phe	Ser	Ile	Leu		
	210				215						220						
Phe	Arg	Asn	Phe	Lys	Glu	Val	Ala	Gln	Pro	Tyr	Arg	Ala	Pro	Gln	Leu		
225				230					235					240			
Thr	Gln	Leu	Phe	Val	Asp	Glu	Ala	Ala	Asp	Phe	Ile	Glu	Arg	Asn	Lys		
		245						250					255				
Glu	Lys	Pro	Phe	Phe	Leu	Tyr	Leu	Ala	Phe	Leu	Arg	Leu	His	Val	His		
	260							265					270				
Thr	Pro	Leu	Phe	Ser	Pro	Ala	Glu	Asp	Leu	Glu	Ser	Lys	Asp	Phe	Leu		
	275					280						285					
Gly	Arg	Ser	Gln	Arg	Gly	Arg	Tyr	Gly	Asp	Leu	Val	Glu	Glu	Met	Asp		
	290				295						300						
Asp	Leu	Val	Gly	Arg	Val	Leu	Asp	Ala	Leu	Glu	Asp	Leu	Gly	Leu	Leu		
305				310					315					320			
Asp	Asn	Thr	Leu	Val	Ile	Phe	Thr	Ser	Asp	Asn	Gly	Ala	His	Leu	Glu		
		325						330					335				
Gly	Thr	Pro	Glu	Trp	Tyr	Gly	Gly	Gly	Asn	Gly	Pro	Leu	Lys	Gly	Gly		
	340							345					350				
Lys	Gly	Tyr	Gly	Ser	Leu	Tyr	Glu	Gly	Gly	Ile	Arg	Val	Pro	Leu	Leu		
	355					360						365					
Val	Arg	Trp	Pro	Gly	Gly	Ile	Ala	Pro	Ala	Gly	Arg	Val	Lys	Glu	Lys		
	370				375						380						
Ser	Glu	Leu	Val	Ser	His	Val	Asp	Leu	Ala	Pro	Thr	Ile	Leu	Asp	Leu		
385				390					395					400			

Ala Gly Ala Pro Leu Pro Lys Val Ala Asn Gly Ala Lys Asp Arg Pro
 405 410 415
 Leu Asp Gly Val Ser Leu Leu Pro Leu Leu Leu Gly Gly Ala Ala Pro
 420 425 430
 Ser Arg Arg Ala His Glu Thr Leu Phe His Tyr Asn Gly Lys Gly Arg
 435 440 445
 Lys Leu Arg Ala Val Arg Trp Pro Arg Lys Ser Gly Lys Thr Pro Lys
 450 455 460
 Leu Lys Ala His Phe Phe Thr Pro Ala Phe Asp Asp Asp Thr Asn Asn
 465 470 475 480
 Gly Trp Glu Cys Val Gly Thr Val Ser Gln Ala Asp Asp Ile Glu Asp
 485 490 495
 Cys Arg Cys Glu Gly Val Glu Thr Val Thr His His Asp Pro Pro Glu
 500 505 510
 Leu Tyr Asp Leu Ser Arg Asp Pro
 515 520

<210> 11
 <211> 1578
 <212> DNA
 <213> homo sapiens

<400> 11
 atgggctggc tttttctaaa ggttttgttg gcgggagtga gtttctcagg atttctttat 60
 cctcttgtagg atttttgcac cagtgggaaa acaagaggac agaagccaaa ctttgtgatt 120
 attttggcgg atgacatggg gtgggggtgac ctgggagcaa actgggcaga aacaaaggac 180
 actgccaacc ttgataagat ggcttcggag ggaatgaggt ttgtggattt ccatgcagct 240
 gcctccacct gctcacctc ccgggcttcc ttgctcaccg gccggcttgg ccttcgcaat 300
 ggagtcacac gcaactttgc agtcacttct gtgggaggcc ttccgctcaa cgagaccacc 360
 ttggcagagg tgctgcagca ggcggtttac gtcactggga taataggcaa atggcatctt 420
 ggacaccacg gctcttatca cccaacttc cgtggttttg attactactt tggaatccca 480
 tatagccatg atatgggctg tactgatact ccagggtaca accaccctcc ttgtccagcg 540
 tgtccacagg gtgatggacc atcaaggaac cttcaaagag actgttacac tgacgtggcc 600
 ctccctcttt atgaaaacct caacattgtg gagcagccgg tgaacttgag cagccttgcc 660
 cagaagtatg ctgagaaagc aaccagttc atccagcgtg caagcaccag cgggaggccc 720
 ttctgtctct atgtggctct ggcccacatg cacgtgccct taccctgac tcagctacca 780
 gcagcgccac ggggcagaag cctgtatggt gcagggtctt gggagatgga cagtctgggtg 840
 ggccagatca aggacaaagt tgaccacaca gtgaaggaaa acacattcct ctggtttaca 900
 ggagacaatg gcccggtggc tcagaagtgt gagctagcgg gcagtgtggg tcccttctact 960
 ggattttggc aaactcgtca agggggaagt ccagccaagc agacgacctg ggaaggaggg 1020
 caccgggtcc cagcactggc ttactggcct ggagagttc cagttaatgt caccagcact 1080
 gccttggttaa gcgtgctgga catttttcca actgtggtag ccctggccca ggccagctta 1140
 cctcaaggac ggcgctttga tgggtgtggac gtctccgagg tgctcttttg ccggtcacag 1200
 cctgggcaca ggggtgctgt ccacccaac agcggggcag ctggagagtt tggagccctg 1260
 cagactgtcc gcctggagcg ttacaaggcc ttctacatta ccggtggagc cagggcgctgt 1320
 gatgggagca cggggcctga gctgcagcat aagtttctc tgattttcaa cctggaagac 1380
 gataccgcag aagctgtgcc cctagaaaga ggtgtgctg agtaccaggc tgtgtgtgcc 1440
 gaggtcagaa aggttcttgc agacgtctc caagacattg ccaacgacaa catctccagc 1500
 gcagattaca ctgaggaccc ttcagtaact ccctgctgta atccctacca aattgcctgc 1560
 cgctgtcaag ccgcataa 1578

<210> 12
 <211> 2616
 <212> DNA
 <213> homo sapiens

<400> 12
 atgaagtatt cttgctgtgc tctggttttg gctgtcctgg gcacagaatt gctgggaagc 60
 ctctgttcga ctgtcagatc cccgaggttc agaggacgga tacagcagga acgaaaaaac 120
 atccgacca acattattct tgtgcttacc gatgatcaag atgtggagct ggggtccctg 180
 caagtcatga acaaaacgag aaagattatg gaacatggg gggccacctt catcaatgcc 240
 tttgtgacta caccatgtg ctgcccgtca cgttcctcca tgctcaccgg gaagtatgtg 300
 cacaatcaca atgtctacac caacaacgag aactgctctt cccctcctg gcaggccatg 360
 catgagcctc ggacttttgc tgtatatctt aacaacactg gctacagaac agcctttttt 420
 ggaaaatacc tcaatgaata taatggcagc tacatcccc ctgggtggcg agaattggctt 480
 ggattaatca agaattctcg cttctataat tacactgttt gtcgcaatgg catcaaagaa 540
 aagcatggat ttgattatgc aaaggactac ttcacagact taatcactaa cgagagcatt 600

aattacttca	aaatgtctaa	gagaatgtat	ccccataggg	ccgttatgat	ggtgatcagc	660
cacgctgcgc	cccacggccc	cgaggactca	gccccacagt	tttctaaact	gtaccccaat	720
gcttcccaac	acataactcc	tagttataac	tatgcaccaa	atatggataa	acactggatt	780
atgcagtaca	caggaccaat	gctgcccac	cacatggaat	ttacaaacat	tctacagcgc	840
aaaaggctcc	agactttgat	gtcagtggtg	gattctgtgg	agaggctgta	taacatgctc	900
gtggagacgg	gggagctgga	gaatacttac	atcatttaca	ccgccgacca	tggttaccat	960
attgggcagt	ttggactggg	caaggggaaa	tccatgccat	atgactttga	tattcgtgtg	1020
ccttttttta	ttcgtggtcc	aagtgtagaa	ccaggatcaa	tagtcccaca	gatcgttctc	1080
aacattgact	tggccccac	gacactggat	attgctgggc	tcgacacacc	tcctgatgtg	1140
gacggcaagt	ctgtcctcaa	acttctggac	ccagaaaagc	caggtaacag	gtttcgaaca	1200
aacaagaagg	ccaaaatttg	gcgtgatata	ttcctagtgg	aaagaggcaa	atcttctacgt	1260
aagaagggaag	aatccagcaa	gaatatccaa	cagtcaaata	acttgcccaa	atatgaacgg	1320
gtcaaagaac	tatgccagca	ggccaggtac	cagacagcct	gtgaacaacc	ggggcagaag	1380
tggcaatgca	ttgaggatag	atctggcaag	cttcgaattc	acaagtgtaa	aggaccacgt	1440
gacctgtca	cagtccggca	gagcacgcgg	aacctctacg	ctcgcggctt	ccatgacaaa	1500
gacaaagagt	gcagttgtag	ggagtctggt	taccgtgcca	gcagaagcca	aagaaagagt	1560
caacggcaat	tcttgagaaa	ccaggggact	ccaaagtaca	agcccagatt	tgtccatact	1620
cggcagacac	gttccttgct	cgtcgaattt	gaagggtgaa	tatatgacat	aaatctggaa	1680
gaagaagaag	aattgcaagt	gttgcaacca	agaaacattg	ctaagcgtca	tgatgaaggc	1740
cacaaggggc	caagagatct	ccaggcttcc	agtgggtggc	acaggggcag	gatgctggca	1800
gatagcagca	acgccgtggg	cccacctacc	actgtccgag	tgacacacaa	gtgttttatt	1860
cttcccaatg	actctatcca	ttgtgagaga	gaactgtacc	aatcggccag	agcgtggaag	1920
gaccataagg	catacattga	caaagagatt	gaagctctgc	aagataaaat	taagaattta	1980
agagaagtga	gaggacatct	gaagagaagg	aagcctgagg	aatgtagctg	cagtaaacaa	2040
agctattaca	ataaagagaa	aggtgtaaaa	aagcaagaga	aattaagag	ccatcttcac	2100
ccattcaagg	aggctgctca	ggaagtagat	agcaaactgc	aacttttcaa	ggagaacaac	2160
cgtagaggaga	agaaggagag	gaaggagaag	agacggcaga	ggaaggggga	agagtgcagc	2220
ctgcctggcc	tcacttgctt	cacgcgatgac	aacaaccact	ggcagacagc	cccgttctgg	2280
aacctgggat	ctttctgtgc	ttgcacgagt	tctaacaata	acacctactg	gtgtttgcgt	2340
acagttaatg	agacgcataa	ttttcttttc	tgtgagtttg	ctactggctt	tttgaggtat	2400
tttgatatga	atacagatcc	ttatcagctc	acaaatacag	tgcacacggg	agaacgaggc	2460
attttgatc	agctacacgt	acaactaatg	gagctcagaa	gctgtcaagg	atataagcag	2520
tgcaacccaa	gacctaaaga	tcttgatgtt	ggaaataaag	atggaggaag	ctatgacct	2580
cacagaggac	agttatggga	tggatgggaa	ggttaa			2616

<210> 13

<211> 1710

<212> DNA

<213> homo sapiens

<400> 13

atgcacaccc	tcactggctt	ctctctggtc	agcctgctca	gcttcggcta	cctgtcctgg	60
gactgggcca	agccgagctt	cgtggccgac	gggcccgggg	aggctggcga	gcagccctcg	120
gccgctccgc	cccagcctcc	ccacatcatc	ttcatcctca	cggacgacca	aggctaccac	180
gacgtgggct	accatggttc	agatatacag	acccctacgc	tggaacaggct	ggcggccaag	240
ggggctcaagt	tggagaatta	ttacatccag	cccatctgca	cgccttcgcg	gagccagctc	300
ctcactggca	ggtaccagat	ccacacagga	ctccagcatt	ccatcatccg	cccacagcag	360
cccaactgcc	tgcccctgga	ccagggtgaca	ctgccacaga	agctgcagga	ggcaggttat	420
tccacccata	tggtagggcaa	gtggcacctg	ggcttctacc	ggaaggagtg	tctgcccacc	480
cgtaggggct	tcgacacact	cctgggctcg	ctcacgggca	atgtggacta	ttacacctat	540
gacaactgtg	atggcccagg	cgtgtgcggc	ttcgacctgc	acgagggtga	gaatgtggcc	600
tgggggctca	gcggccagta	ctccactatg	ctttacgccc	agcgcgccag	ccatatcctg	660
gccagccaca	gccctcagcg	tcccctcttc	ctctatgtgg	ccttcagggc	agtacacaca	720
cccctgcagt	cccctcgtga	gtacctgtac	cgctaccgca	ccatgggcaa	tgtggcccgg	780
cggaagtacg	cggccatggg	gacctgcatg	gatgaggctg	tgcgcaacat	cacctggggc	840
ctcaagcgct	acggtttcta	caacaacagt	gtcatcatct	tctccagtga	caatgggtggc	900
cagactttct	cggggggcag	caactggccg	ctccgaggac	gcaagggcac	ttattgggaa	960
ggtggcgtgc	ggggcctagg	ctttgtccac	agtcccctgc	tcaagcgaaa	gcaacggaca	1020
agccgggcac	tgatgcacat	cactgactgg	tacccgaccc	tggtaggtct	ggcagggtgg	1080
accacctcag	cagccgatgg	gctagatggc	tacgacgtgt	ggccggccat	cagcgagggc	1140
cgggcctcac	cacgcacgga	gatcctgcac	aacattgacc	cactctacaa	ccatgcccag	1200
catggctccc	tggaggcgcg	ctttggcate	tggaacaccc	cogtgcaggc	tgccatccgc	1260
gtgggtgagt	ggaagctgct	gacaggagac	ccgggctatg	gcgattggat	cccaccgcag	1320
acactggcca	ccttccgggg	tagctggtgg	aacctggaac	gaatggccag	tgtccgcccag	1380
gccgtgtggc	tcttcaacat	cagtgtgtac	ccttatgaac	gggaggacct	ggctggccag	1440
cggcctgatg	tggtagggc	cctgctggct	cgcctggccg	aatataaccg	cacagccatc	1500
ccggtacgct	acccagctga	gaacccccgg	gctcctcctg	actttaatgg	gggtgcttgg	1560

gggccctggg	ccagtgatga	ggaagaggag	gaagaggaag	ggagggctcg	aagcttctcc	1620
cggggtcgtc	gcaagaaaaa	atgcaagatt	tgcaagcttc	gatccttttt	ccgtaaactc	1680
aacaccaggc	taatgtccca	acggatctga				1710

<210> 14

<211> 1800

<212> DNA

<213> homo sapiens

<400> 14

atggctccca	ggggctgtgc	ggggcatccg	cctccgcctt	ctccacaggc	ctgtgtctgt	60
cctggaaaga	tgctagcaat	gggggcgctg	gcaggattct	ggatcctctg	cctcctcaact	120
tatggttacc	tgctctgggg	ccaggcctta	gaagaggagg	aagaaggggc	cttactagct	180
caagctggag	agaaactaga	gcccagcaca	acttccacct	cccagcccca	tctcatthttc	240
atcctagcgg	atgatcaggg	atthtagagat	gtgggttacc	acggatctga	gattaaaaaca	300
cctactcttg	acaagctcgc	tgccgaagga	gttaaactgg	agaactacta	tgtccagcct	360
atthgcacac	catccaggag	tcagthttatt	actggaaagt	atcagataca	caccggactt	420
caacattcta	tcataagacc	tacccaaccc	aactgthttac	ctctggacaa	tgccacccta	480
cctcagaaac	tgaaggagg	tggatattca	acgcatatgg	tcggaaaatg	gcacttgggt	540
thttacagaa	aagaatgcat	gcccaccaga	agaggatttg	atacctthtt	tggttccctt	600
ttgggaagt	gggattacta	tacacactac	aatagtga	gtcctgggat	gtgtggctat	660
gacttgtag	aaaacgacaa	tgctgcctgg	gactatgaca	atggcatata	ctccacacag	720
atgtacactc	agagagtaca	gcaaacttta	gcttcccata	acccacaaa	gcctatattt	780
ttatatattg	cctatcaagc	tgthtcattca	ccactgcaag	ctcctggcag	gtatttcgaa	840
cactaccgat	ccattatcaa	cataaacagg	aggagatatg	ctgccatgct	ttcctgctta	900
gatgaagcaa	tcaacaacgt	gacattggct	ctaaagactt	atggthttcta	taacaacagc	960
attatcattt	actcttcaga	taatgggtgc	cagcctacgg	caggagggag	taactggcct	1020
ctcagaggta	gcaaaggaac	atattgggaa	ggagggatcc	gggctgtagg	ctthgtgcat	1080
agcccacttc	tgaaaaacaa	gggaacagtg	tgtaaggaac	ttgtgcacat	cactgactgg	1140
taccccactc	tcatttcact	ggctgaagga	cagattgatg	aggacattca	actagatggc	1200
tatgatattc	gggagaccat	aagtgagggg	cttcgctcac	cccagtaga	tattthtgc	1260
aacattgacc	ccatatacac	caaggcaaaa	aatggctcct	gggcagcagg	ctatgggatc	1320
tggaacactg	caatccagtc	agccatcaga	gtgcagcact	ggaaattgct	tacaggaaat	1380
cctggctaca	gcgactgggt	ccccctcag	tctthtcagca	acctgggacc	gaaccgggtg	1440
cacaatgaac	ggatcacctt	gtcaactggc	aaaagtgtat	ggctthttcaa	catcacagcc	1500
gaccatattg	agagggtgga	cctatctaac	aggtatccag	gaatcgtgaa	gaagctccta	1560
cggaggctct	cacagttcaa	caaaactgca	gtgccggtca	ggtatcccc	caaagacccc	1620
agaagtaacc	ctaggctcaa	tgagggggtc	tggggaccat	ggtataaaga	ggaaaccaag	1680
aaaaagaagc	caagcaaaaa	tcaggctgag	aaaaagcaaa	agaaaagcaa	aaaaaagaag	1740
aagaaacagc	agaaagcagt	ctcaggttca	acttgccatt	caggtgttac	thgtggataa	1800

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 August 2001 (02.08.2001)

PCT

(10) International Publication Number
WO 01/55411 A3

(51) International Patent Classification⁷: **C12N 15/55**,
5/10, 9/16, C07K 16/40, G01N 33/53, C12Q 1/68, A61K
38/46

(21) International Application Number: PCT/US01/03266

(22) International Filing Date: 31 January 2001 (31.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/495,823 31 January 2000 (31.01.2000) US

(71) Applicant (for all designated States except US): **MIL-
LENNIUM PHARMACEUTICALS, INC.** [US/US]; 75
Sidney Street, Cambridge, MA 02139 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GLUCKSMANN, Maria, Alexandra** [AR/US]; 33 Summit Road, Lexington, MA 02173 (US). **WILLIAMSON, Mark** [US/US]; 15 Stonecrest Drive, Saugus, MA 01906 (US). **RUDOLPH-OWEN, Laura, A.** [US/US]; 186 Arborway, #1, Jamaica Plain, MA 02130 (US). **TSAI, Fong-Ying** [—/US]; 15 Montclair Road, Newton, MA 02468 (US).

(74) Agents: **KRON, Eric, J. et al.**; Alston & Bird LLP, Bank of America Plaza, 101 South Tryon Street, Suite 4000, Charlotte, NC 28280-4000 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
21 March 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/55411 A3

(54) Title: HUMAN SULFATASES

(57) Abstract: The present invention relates to newly identified human sulfatases. In particular, the invention relates to sulfatase polypeptides and polynucleotides, methods of detecting the sulfatase polypeptides and polynucleotides, and methods of diagnosing and treating sulfatase-related disorders. Also provided are vectors, host cells, and recombinant methods for making and using the novel molecules.

INTERNATIONAL SEARCH REPORT

International Application No
PC, US 01/03266

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/55 C12N5/10 C12N9/16 C07K16/40 G01N33/53
C12Q1/68 A61K38/46

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EMBL, SEQUENCE SEARCH, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL [Online] ACCESSION NO: AB023218, 9 April 1999 (1999-04-09) OHARA T ET AL: "Homo sapiens mRNA for KIAA1001 protein, complete cds" XP002181669 nucleotides 221-2373 ---	1,2,8-13
X	DATABASE EMBL [Online] ACCESSION NO: AI423178, 15 March 1999 (1999-03-15) NATIONAL CANCER INSTITUTE: "Homo sapiens cDNA clone IMAGE:2097922 3', mRNA sequence" XP002181670 nucleotides 9-440 --- -/--	1,2

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

31 October 2001

Date of mailing of the international search report

08.01.2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

CUPIDO, M

INTERNATIONAL SEARCH REPORT

International Application No

PC, / US 01/03266

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	STEIN C ET AL: "CLONING AND EXPRESSION OF HUMAN ARYLSULFATASE A" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 264, no. 2, 1989, pages 1252-1259, XP002181668 ISSN: 0021-9258 the whole document -----	1-13

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/03266

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1, 2, 8, 9, 12, 16, 19 and 22
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-26 (partly)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-26 (all partly)

An isolated nucleic acid molecule which comprises a nucleotide sequence which is at least 60% identical to SEQ ID NO:2, or comprises a fragment of at least 20 nucleotides of SEQ ID NO:2 or 11, or encodes a polypeptide with SEQ ID NO:1 a fragment or an allelic variant thereof, its complementing strand, host cells containing this sequence, the encoded polypeptide and antibodies binding thereto, as well as a various diagnostic and therapeutic applications thereof.

2. Claims: 1-26 (all partly)

An isolated nucleic acid molecule which comprises a nucleotide sequence which is at least 60% identical to SEQ ID NO:4, or comprises a fragment of at least 20 nucleotides of SEQ ID NO:4 or 12, or encodes a polypeptide with SEQ ID NO:3 a fragment or an allelic variant thereof, its complementing strand, host cells containing this sequence, the encoded polypeptide and antibodies binding thereto, as well as a various diagnostic and therapeutic applications thereof.

3. Claims: 1-26 (all partly)

An isolated nucleic acid molecule which comprises a nucleotide sequence which is at least 60% identical to SEQ ID NO:6, or comprises a fragment of at least 20 nucleotides of SEQ ID NO:6 or 13, or encodes a polypeptide with SEQ ID NO:5 a fragment or an allelic variant thereof, its complementing strand, host cells containing this sequence, the encoded polypeptide and antibodies binding thereto, as well as a various diagnostic and therapeutic applications thereof.

4. Claims: 1-26 (all partly)

An isolated nucleic acid molecule which comprises a nucleotide sequence which is at least 60% identical to SEQ ID NO:8, or comprises a fragment of at least 20 nucleotides of SEQ ID NO:8 or 14, or encodes a polypeptide with SEQ ID NO:7 a fragment or an allelic variant thereof, its complementing strand, host cells containing this sequence, the encoded polypeptide and antibodies binding thereto, as well as a various diagnostic and therapeutic applications thereof.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box 1.2

Claims Nos.: 1, 2, 8, 9, 12, 16, 19 and 22

1. The ATCC deposit numbers of the plasmids referred to in claims 1, 2, 8, 9 and 12 are not provided and a search concerning this subject-matter was therefore not possible.

2. Claim 16 and 22 refer to a compound which binds to a polypeptide of claim 8 without giving a true technical characterisation. Moreover, no such compounds are defined in the application, and the search was therefore limited to antibodies that bind the polypeptide with SEQ ID NO:1.

3. Claim 19 concerns a kit containing a compound which selectively hybridizes to a nucleic acid molecule of claim 1. The search was limited to nucleotide sequences that are complementary to SEQ ID NO:2 or 11.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

THIS PAGE BLANK (USPTO)